Peer Community In Archaeology

RESEARCH ARTICLE

Open Access

Open Peer-Review

Posted
2022-04-26

Cite as

Evin, A., Bouby, L.,
Bonhomme, V., Jeanty, A.,
Jeanjean, M. and Terral, J.-F.
(2022) Archaeophenomics of
ancient domestic plants and
animals using geometric
morphometrics : a review. OSF
Preprints, skeu5, ver. 3 peer-
reviewed and recommended
by Peer community in
Archaeology.
<https://doi.org/10.31219/osf.io/skeu5>

Correspondence

allowen.evin@umontpellier.fr

Recommender

Anneke H. van Heteren

Reviewers

Stefan Schlager and one
anonymous reviewer

Archaeophenomics of ancient domestic plants and animals using geometric morphometrics : a review

Allowen Evin^{*1}, Laurent Bouby¹, Vincent
Bonhomme¹, Angèle Jeanty¹, Marine
Jeanjean¹, Jean-Frédéric Terral¹

¹ ISEM, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France. 2 place Eugène Bataillon, 34095
Montpellier, cedex 5, France.

^{*}Corresponding author

This article has been peer-reviewed and recommended by
Peer Community in Archaeology

<https://doi.org/10.24072/pci.archaeo.100014>

ABSTRACT

Geometric morphometrics revolutionized domestication studies through the precise quantification of the phenotype of ancient plant and animal remains. Geometric morphometrics allow for an increasingly detailed understanding of the past agrobiodiversity and our ability to characterize large scale ancient phenotypes has led to what can be named archaeophenomics : the large scale phenotyping of ancient remains. This review describes advances in the bioarchaeological study of domesticated species and their wild relatives where their phenomes are quantified through geometric morphometrics. The two main questions addressed by archaeophenomics are i) taxonomic identification, including domestication signature, and ii) the inference of the spatio-temporal agrobiodiversity dynamics. Archaeophenomics is a growing field in bioarchaeology of domestic species that will benefit in the near future from advances in artificial intelligence and from an increasing interest in multiproxy approaches combining morphometric data with e.g. isotopes or archaeogenomics.

Keywords: bioarchaeology, phenotypic evolution, artificial selection, domestication, diversification, -omics

Contents

Contents	2
Introduction.....	2
From phenomics to archaeophenomics	2
Aim and scope of the review	3
Geometric morphometrics in archaeophenomics.....	3
Domestic species studied with geometric morphometrics	4
Main bioarchaeological research themes based on GMM data	6
Multi-proxy approaches and future methodological developments	8
Conclusion/perspectives	8
Acknowledgements	8
Funding.....	9
Conflict of interest disclosure.....	9
Supplementary information	9
References.....	9

Introduction

Domesticated species have played a major role in the development of Charles Darwin's work, being the subject of the first chapter of the 'Origin of Species' (Darwin, 1859) and later of a dedicated book, 'The variation of animals and plants under domestication' (Darwin, 1868) where he described the mechanisms of variation in domestic species. With his work, Darwin contributed to the understanding of the morphological changes that occurred during the long process of domestication. For most domesticated species, modern breeds and varieties today present a huge morphological diversity reflecting millennia of human selection for many purposes (*e.g.* food production, work, aestheticism) in various environmental conditions. While the study of the current domestic diversity is mainly carried out in agronomic research, with breed and varietal improvement using molecular breeding programs, a large amount of research has been done to explore the past diversity of domestic species whose remains are found in increasing numbers in archaeological deposits. The methodological development in morphometrics have revolutionized, qualitatively and quantitatively, the study of the phenotype of those remains. Today we have reached a state where the use of several tools, including morphometrics, have allowed phenome (*i.e.* the full set of observable traits) quantification of a large number of archaeological specimens leading to a renew in the study of archaeological remains of domestic species. We here coin the word *archaeophenomics* for such large-scale quantification of phenotypic data from archaeological specimens. Archaeophenomics, *i.e.* phenomics of the past, is an emerging field that will likely become a standard for future bioarchaeological studies. This neologism fulfils the needs to express the new realities of bioarchaeological domestication studies.

From phenomics to archaeophenomics

Phenomics, the analysis of high-dimensional phenotypic data, is part of the 'omics' revolution as genomics or proteomics. Phenomics is the new generation of acquisition and analysis of phenotypic data based on techniques which allow a very large amount of quantitative characters to be acquired and processed with minimal handling time. Assessing the full phenome of an organism is illusional (Houle et al., 2010) and this is even more true in bioarchaeology. Archaeological remains of plants and animals are often altered by

taphonomical processes (*e.g.* preservation, fragmentation). Despite these inherent constraints associated with studying archaeological material, the large quantity of remains allows for large-scale morphometric analyses.

Morphometrics is one of the many tools of *phenomics*. The ‘morphometric revolution’ corresponding to the development of *geometric morphometrics* (GMM), *i.e.* the study of forms in multi-dimensional spaces, allow more in-depth investigation of morphological changes (James Rohlf & Marcus, 1993; Adams et al., 2004). The main improvement of geometric morphometrics compared to the so-called ‘traditional morphometrics’ is that biological forms are no longer captured by sets of independent measurements of lengths or angles, but by sets of point coordinates, improving dramatically the capture of the geometric complexity of these objects (Kuhl & Giardina, 1982; Bookstein, 1991; James Rohlf & Marcus, 1993). Morphometric analyses are often the only available approach for studying the morphology of ancient remains with a fine-scale resolution, while suffering less from preservation limitations than *e.g.* ancient DNA and offering much better possibilities for being carried out on a large scale at a limited cost, in both time and money. This is especially true for plants whose remains are often found charred, a condition strongly detrimental to DNA preservation and which generally prevents the analysis of these remains, at least with current aDNA techniques (Nistelberger et al., 2016). Most of the time, once a specimen is recovered a morphometric analysis can be performed, as long as the structures grabbing the geometrical features analysed are present. While it is always better to analyse complete specimens, fragmented remains can even be studied using a restricted version of the initial protocol since fragmentation does not necessary prevent taxonomical identification (*e.g.* Cornette et al., 2015; Owen et al., 2014, Durocher in press). Geometric morphometric techniques are therefore particularly well adapted to bioarchaeological studies and are a growing field in the discipline.

Archaeophenomics appears to be as a major breakthrough in bioarchaeology, with a drastic quantitative change in the scale of the number of specimens and populations that can be analysed, and a qualitative improvement provided by an increased resolution of the analyses paired with a better description of the morphometric variation with improved detection and visualisation of the shape variation.

Aim and scope of the review

Here we provide an exhaustive review of bioarchaeological studies using archaeophenomics, through geometric morphometrics, to study archaeological remains of domestic species. We restricted our review to studies published in international journals (*i.e.* excluding grey literature), only those focusing on domestic species and that include archaeological specimens. This therefore explicitly excludes: studies of commensal species (*e.g.* rodents (Cucchi, 2008; Valenzuela-Lamas et al., 2011; Cucchi et al., 2014)); studies focusing only on the ancestors and modern relatives of the domestic populations (*e.g.* Late Glacial horse (Bignon et al., 2005), rabbit (Pelletier, 2019)); and finally the numerous studies focusing only on modern domestic specimens (*e.g.* Battesti et al., 2018; Bonhomme et al., 2017; Evin et al., 2022; Evin, Dobney, et al., 2015; Gros-Balthazard et al., 2016; Hanot et al., 2021; Harbers et al., 2020; Neaux et al., 2020; Pelletier, Kotiaho, Niinimäki, & Salmi, 2020, 2021). Yet these studies are of prime interest for the understanding of ancient populations, they fall out of the scope of this review.

Geometric morphometrics in archaeophenomics

The development of geometric modern morphometrics (GMM) (Bookstein, 1991; James Rohlf & Marcus, 1993) came as a response to the conceptual and methodological limits of traditional morphometric methods, such as a better ability to efficiently partition the size and the shape components of the form variation and the possibility of visualizing shape variation. Shape analysis, through geometric morphometrics, allows analysing microscale variation that could not otherwise be identified using traditional techniques. Two main geometric morphometric approaches are currently used in bioarchaeology (fig. 1): **Procrustes approaches** through the acquisition of landmarks and sliding semi-landmark coordinates **and outline analyses** using various methodologies (see below). In bioarchaeology, as in biology, objects are studied in two- or three-dimensions depending on the geometry and size of the remains. Until recently, all archaeobotanical remains appear to have been studied in 2D from digital images, though one recent publication used 3D X-ray-computed tomography to quantify watermelon seeds (Wolcott et al., 2021) (table 1). In addition, archaeobotanical remains are nearly exclusively studied through their outlines geometries using mainly either Elliptic Fourier transforms (EFT) (Giardina & Kuhl, 1977; Kuhl & Giardina, 1982) or natural/orthogonal polynomial equations

(Rohlf, 1990) (table 1) and only few studies use landmarks and sliding semi-landmarks coordinates (Ros et al. 2014; García-Granero et al. 2016; Wolcott et al. 2021) (table 1).

For animals, teeth are studied in 2D and the same applies to some postcranial bones (*e.g.* phalanges, and talus) although it is quite possible to study them in 3D (Hanot et al., 2017; Haruda, 2017) (table 1). The same applies to mandible that are studied either in 2D (cat: (Vigne et al., 2016), dog: (Ameen et al., 2019)) or 3D (dog: (Drake et al., 2017)). Skulls, which are geometrically more complex objects, are studied in 3D, either directly on the specimens using *e.g.* a Microscribe digitizer (Drake & Klingenberg, 2008; Hanot et al., 2017; Geiger et al., 2017) or through 3D model reconstruction obtained by CT-scanning (Schoenebeck et al., 2021) or photogrammetry (dog: (Ameen et al., 2019)). So far, a single study really takes advantage of a CT-scanning technology to analyze the internal structure of the skull that is the inner ear morphometry (Clavel et al., 2021).

Figure 1: Example of geometric morphometric protocols applied to bioarchaeological remains. A Example of Procrustes approaches used to quantify the morphometric variation of canid skulls with 3D landmarks. B Example of protocols for 2D outline analyses used to quantified the morphometric variation of barley grains.

Domestic species studied with geometric morphometrics

We identified a total of 71 studies among which 38 focus on animals and 33 on plants (fig. 2.A, table 1, SI table 1).

For animals, only mammals have been studied (though a PhD thesis should be mentioned on chicken (Foster, 2018)), which also represent the large majority of domesticated animals. Ten species are listed and the most represented taxa are pig (N=16), followed by dog (N=9), equids (N=5, horse and donkey), caprines (sheep and goat) (N=5), camelids (N=3, guanaco and llama), and finally cat, and cattle with a single mention (table 1, fig. 2). Animal studies focus primarily on teeth (N=21), skull (N=10, cranium and mandible), in a much larger majority than postcranial bones (*e.g.* talus, phalanges or calcaneus) (SI-table 1).

For plants, only angiosperms are concerned and 14 species (or group of closely related species) are listed with grapevine that largely outnumber other species in terms of publications number (N=15), followed by olive (N=6), date palm (N=3) while all other species are only represented by a single mention (table 1). A higher number of studies is dedicated to dicotyledons (N=27) than to monocotyledons (N=6) and focus exclusively on fruits and seeds (SI table 1).

Table 1: List of the reviewed publications. An extended version of the table, including research themes, employed methodology, and combination with biomolecular markers can be found in SI-table 1.

Group	Taxa	References
Mammals	Pig	(Cucchi et al., 2008, 2011, 2016, 2021; Dobney et al., 2008; Bartosiewicz et al., 2013; Krause-Kyora et al., 2013; Ottoni et al., 2013; Duval et al., 2015, 2018; Evin, Flink, et al., 2015; Balasse et al., 2016; Frémondeau et al., 2017; Price & Evin, 2017; Bopp-Ito et al., 2018; Marom et al., 2019)
	Dog	(Drake & Klingenberg, 2008; Drake et al., 2015, 2017; Daza Perea, 2017; Geiger et al., 2017; Fisher, 2019; Ameen et al., 2019; Manin & Evin, 2020; Schoenebeck et al., 2021)
	Caprines (sheep and goat)	(Haruda, 2017; Pöllath et al., 2018, 2019; Haruda et al., 2019; Colominas et al., 2019)
	Equids (horse and donkey)	(Seetah et al., 2016; Cucchi et al., 2017; Hanot et al., 2017; Chuang & Bonhomme, 2019; Clavel et al., 2021)
	Camelids (guanaco and llama)	(Hernández et al., 2021)
	Cat	(Vigne et al., 2016)
	Cattle	(Cucchi et al., 2019)
Monocotyledons	Date palm	(Terral et al., 2012; Gros-Balthazard et al., 2017; Sallon et al., 2020)
	Barley	(Ros et al., 2014)
	Millet	(García-Granero et al., 2016)
	Wheat	(Bonhomme et al., 2016)
Dicotyledons	Lemon	(Grasso et al., 2018)
	Melon	(Sabato et al., 2019)
	Grapevine	(Bacilieri et al., 2017; Bonhomme et al., 2020; Bonhomme, Terral, et al., 2021; Bouby et al., 2018, 2021; Figueiral et al., 2015; Mariotti Lippi et al., 2020; Margaritis et al., 2021; Moricca et al., 2021; Orrù, Grillo, Lovicu, Venora, & Bacchetta, 2013; Pagnoux et al., 2015, 2021; Terral et al., 2010; Uccesu et al., 2015, 2016; Valamoti et al., 2020)
	Olive	(Terral et al., 2004, 2021; Newton et al., 2006, 2014; Bourgeon et al., 2018; Margaritis et al., 2021)
	Opium poppy	(Jesus et al., 2021)
	Cherry	(Burger et al., 2011)
	Pulses (grass pea, lentil, broad bean)	(Tarongi et al., 2020)
	Watermelon	(Wolcott et al., 2021)

Since 2004, year of the first publication included in this review (Terral et al., 2004), the yearly number of published bioarchaeological studies on domestic animal and plant species using geometric morphometrics is steadily increasing (Fig. 1.B).

Figure 2: Bioarchaeological archaeophenomic studies, concerning domestic species and using geometric morphometrics, represented by taxa (A) and year (B). A: relative frequencies of the different groups studied. B: Evolution of the number of studies per year. Details of the references can be found in table 1 and SI table 1.

The number of phenomes now available greatly differs between species with grapevine coming out on top with a maximum of 2223 archaeological seeds having been quantified in one single study (Pagnoux et al., 2021). The number of papers published per year is still low compared to *e.g.* palaeogenomic studies (~1480 references obtained for a quick online search of the term “palaeogenomics” in google scholar the 11/02/2022). Numbers of studied individuals are usually lower for animals which are mostly represented by fewer individuals per archaeological assemblage. It should be noted, however, that the concept of ‘individual’ differs here between archaeozoology and archaeobotany, since an animal will be represented by single elements such as a cranium or a lower right third molar, while numerous studied individuals (*e.g.* seeds or fruit-stones) may come from a single plant individual.

Main bioarchaeological research themes based on GMM data

Archaeophenomics through geometric morphometrics is increasingly used for bio-archaeological studies for two main purposes that are the taxonomic identification of the archaeological remains including the domestication signature, and to assess the agrobiodiversity variation in both time and space primarily related to processes of colonisation-dispersal, adaptation, diversification and changes in husbandry or cultivation practices or cultural choices.

Taxonomic identification and domestication signature

A prerequisite of many bioarchaeological studies is to perform a taxonomical identification of the remains, either by identifying the taxa to which the specimen belongs and/or to identify its wild or domestic status. Archaeological remains are often fragmented or altered due to either taphonomic due to various anthropic and taphonomic processes (*e.g.* charring, butchery or culinary preparation) processes rendering their identification potentially delicate. Geometric morphometric protocols are now available to distinguish (more or less effectively depending on the study and model) morphologically close mammalian species of equids, bovids, sheep-goats, camelids, canids, cats and pigs (table 1). Taxonomic identification of plant remains is even more challenging due to the larger number of species (or sub-species) potentially occurring at an archaeological site. Botanical studies dealing with taxonomic identification include cereals, such as barley, millet, einkorn and emmer, opium poppy, pulses, citrus, melon, watermelon, date palm and prunus species (SI table 1). For other remains whose species identification is unambiguous, the question of the wild and domestic

status distinction and identification may arise. This is especially true for species that have a wild ancestor with a wide geographical range and for which the wild and domestic populations have coexisted for millennia. This is the case for nearly all species with the exception of those having an ancestor leaving in a restricted geographic area (*e.g.* sheep, goat and most cereals). This geographic proximity can also be source of hybridization between wild and domestic individuals as already documented from genomic and palaeogenomic data. for *e.g.* pig (Frantz et al., 2019), dog (Pilot et al., 2018), grapevine (Myles et al., 2011; Riaz et al., 2018) or date palm (Gros-Balthazard et al., 2017) or among individuals of distinct species as evidenced for north African date palm (Flowers et al., 2019) which can render their morphometric identification even more challenging. In addition, studies looking at bridging archaeological samples to modern breeds or varieties, or groups of them, are more often found for perennial clonal plants, that show an extended varietal diversity and for which varieties can theoretically persist substantially unchanged for centuries or even millennia through vegetative multiplication (*e.g.* grapevine, olive, *Prunus* species and date palm (table 1)). For non-perennial clonal organisms, such as animals, it seems that direct comparison to specific modern breeds has been done, so far, only for dogs (Geiger et al., 2017; Schoenebeck et al., 2021), and that such comparison can be questioned as the intensification of selective pressures during the last centuries likely dramatically altered ancient morphologies.

Another specificity of archaeobotanical remains compared to zooarchaeological ones is that remains are often found charred and that charring can affect their size and shape and therefore their taxonomic identification. An important effort has been made for multiple taxa to understand the effect of charring on the morphometric results and their interpretations (*e.g.* cereals (Bonhomme et al., 2017; Ros et al., 2014), grapevine (Ucchesu et al., 2016; Bouby et al., 2018), and olive (Terral et al., 2004)).

A significant proportion of the domestic species whose remains are found during archaeological excavations have been the subject of geometric morphometrics and such approaches have been found effective for taxonomic identification of the remains. The many available protocols can now be adapted to nearby species not yet subjected to such studies.

Documenting spatio-temporal variation of past agrobiodiversity

A large number of archaeophenomic studies explore the morphometric spatio-temporal variation of domestic populations. Such studies span either long periods of time of several millennia (*e.g.* Pagnoux et al., 2021; Price & Evin, 2017; Terral et al., 2004), or a much shorter period of no more than a century (*e.g.* Drake & Klingenberg, 2008). Time and space are intertwined components of bioarchaeological studies. It is however possible to study them separately by comparing either diachronic populations of a single locality or synchronous populations of various geographic origins.

The studies that explore the geographic variation between populations from a single chrono-cultural period (SI-table 1) evidenced that both geographically near and far populations can show morphometric differences. In term of interpretations, a geographic structuration of synchronous populations may correspond not exclusively to local environmental adaptation, different husbandry/cultivation practices, cultural choices or distinct genetic lineages.

Similarly, diachronic differences between populations originating from the same geographic area can reveal either changes in human practices, spread of new genetic stock, environmental variation, or drift. Such comparison between diachronic populations (SI-table 1) can reveal both long term variation, but also more abrupt morphological shift between periods.

In comparison, fewer studies focus both on time and space (SI-Table 1). Generally, while absence of differences between assemblages cannot lead to the conclusion that they belong to similar populations, on the other hand, the existence of differences allows to hypothesize the existence of distinct populations with at least limited cultural or genetic exchanges. As a consequence, morphometric analyses can make only limited contribution to mobility studies, that can be better explored using *e.g.* biomolecular markers such as ancient DNA or isotopes. In the same way, phenotypic proximity does not necessarily reflect genetic proximity (*i.e.* phylogeny) due to natural or anthropic selection. During domestication, human populations have selected certain traits such as larger quantity of meat or of fruit size. In archaeophenomics studies, the target of human selection during domestication and subsequent diversification is not necessarily the target of the morphometric analysis. For cereals, grain size was likely intentionally selected, but not their shape, while for fruit stones neither size or shape were likely directly targeted, even if in some cases (*e.g.* at least for grapevine) seed and fruit measurements covariate (Bonhomme et al., 2020). Similarly, in mammals, it is unlikely that

teeth, that are commonly studied and considered as a phenotypic marker of adaptation to natural or anthropic environment, were not likely the aim of human selection that primarily targeted primary (*e.g.* meat) or secondary products (*e.g.* milk, wool). In all these cases, where domestic and wild populations differ in size and shape of anatomical elements non targeted by selection, other evolutionary pressures and mechanisms such as drift, genetic hitchhiking or indirect selection (*e.g.* morpho-functional constraint) can be invoked. In addition, many anatomical structures are polygenic (*e.g.* Harjunmaa et al., 2012), or the genes involved are not known. Moreover, not all anatomical elements necessarily evolve in parallel, at the same rate or following the same direction (*e.g.* Geiger & Sánchez-Villagra, 2018).

Multi-proxy approaches and future methodological developments

Archaeobotanical remains are often conserved through charring which is detrimental to DNA preservation (Nistelberger et al., 2016) rendering the combination of such approaches with morphometric data impossible. This is however possible to the less frequently found waterlogged remains where DNA can be preserved and the results compared to morphometric data (Bacilieri et al., 2017; Bouby et al., 2021).

For animals, several studies combined geometric morphometrics with ancient DNA (*e.g.* pig: (Evin, Flink, et al. 2015), dog (Ameen et al., 2019)), geometric morphometrics and isotopes (pig: (Cucchi et al., 2016; Frémontdeau et al., 2017)), or the combination of the three approaches geometric morphometrics, ancient DNA and isotopes (pig: (Balasse et al., 2016)). Isotopic analyses are increasingly used in archaeobotany (*e.g.* Fiorentino, Ferrio, Bogaard, Araus, & Riehl, 2015), but not yet in combination with other approaches.

Artificial intelligence is increasingly used in biology (*e.g.* Ching et al., 2018; Hassoun et al., 2021) and archaeology (*e.g.* Bickler, 2021; Horn et al., 2021), but only few bioarchaeological studies use yet such approaches (*e.g.* Miele, Dussert, Cucchi, & Renaud, 2020). Machine learning in general, and deep learning using convolutional neural networks in particular, will certainly help in the future for automatic data acquisition such as landmark coordinates (*e.g.* Devine et al., 2020), image post-treatment prior to outline analyses, and/or directly for binary (status) or multiple (species identification) classification tasks. In addition, the ever-growing motivation to share data and knowledge should drastically extent the chrono-cultural and geographic scopes of the studies allowing comparisons not possible before. This would be possible only after careful inter-operator and methodological comparisons (Evin, Bonhomme, and Claude 2021). Other future lines of research will also certainly focus on evo-devo perspectives (Bonhomme et al., 2020), form-function interactions (Harbers et al., 2020; Neaux et al., 2020) as well as further exploration of the genotype-phenotype relationships (Schoenebeck & Ostrander, 2013).

Conclusion/perspectives

Archaeophenomics through geometric morphometrics allows addressing questions regarding the micro-evolutionary processes that accompanied the long history of domestic species in an unprecedented way. Such approaches are increasingly used in bioarchaeology and are becoming one of the many approaches now available to us to study past populations. The future of phenotypic studies will require carefully thought, managed and open large-scale databases, precisely contextualised archaeologically, and combining the whole set of available approaches, if possible carried out on the exact same specimens. As for the relatively recent research fields of palaeoproteomics or palaeogenomics, this review shows that archaeophenomics definitely corresponds to a renew of domestication studies deserving a new terminology. This review attempts to list all the studies in the scope of archaeophenomics where the phenomes of domestic species are quantified through geometric morphometrics. The many approaches now available pave the way to future research expanding the diversity of studied species and the archaeological questions that can be addressed through archaeophenomics.

Acknowledgements

We thank Stefan Schlager and one anonymous reviewer for their comments on the earlier draft of the manuscript. Version 3 of this preprint has been peer-reviewed and recommended by Peer Community In Archaeology (<https://doi.org/10.24072/pci.archaeo.100014>).

Funding

This work has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (Grant Agreement 852573), the French National Agency (ANR-16-CE27-0013) and the International Research Program EVOLEA (France - Morocco) (CNRS-INEE).

Conflict of interest disclosure

The authors declare they have no conflict of interest relating to the content of this article.

Supplementary information

SI-Table 1 is available in the open repository OSF with the DOI : 10.17605/OSF.IO/T3Q96.

References

- Adams DC, Rohlf J, Slice D (2004) Geometric morphometrics: Ten years of progress following the 'revolution.' *Italian Journal of Zoology*, 71, 5–16. <https://doi.org/10.1080/11250000409356545>
- Ameen C, Feuerborn TR, Brown SK, Linderholm A, Hulme-Beaman A, Lebrasseur O, Sinding MHS, Lounsberry ZT, Lin AT, Appelt M, Bachmann L, Betts M, Britton K, Darwent J, Dietz R, Fredholm M, Gopalakrishnan S, Goriunova OI, Grønnow B, Haile J, Hallsson JH, Harrison R, Heide-Jørgensen MP, Knecht R, Losey RJ, Masson-MacLean E, McGovern TH, McManus-Fry E, Meldgaard M, Midtdal Å, Moss ML, Nikitin IG, Nomokonova T, Pálsdóttir AH, Perri A, Popov AN, Rankin L, Reuther JD, Sablin M, Schmidt AL, Shirar S, Smiarowski K, Sonne C, Stiner MC, Vasyukov M, West CF, Ween GB, Wennerberg SE, Wiig Ø, Woollett J, Dalén L, Hansen AJ, Gilbert MTP, Sacks BN, Frantz L, Larson G, Dobney K, Darwent CM, Evin A (2019) Specialized sledge dogs accompanied Inuit dispersal across the North American Arctic. *Proceedings of the Royal Society B: Biological Sciences*, 286, 20191929. <https://doi.org/10.1098/rspb.2019.1929>
- Bacilieri R, Bouby L, Figueiral I, Schaal C, Terral J-F, Breton C, Picq S, Weber A, Schlumbaum A (2017) Potential of combining morphometry and ancient DNA information to investigate grapevine domestication. *Vegetation History and Archaeobotany*, 26, 345–356. <https://doi.org/10.1007/s00334-016-0597-4>
- Balasse M, Evin A, Tornero C, Radu V, Fiorillo D, Popovici D, Andreescu R, Dobney K, Cucchi T, Bălăşescu A (2016) Wild, domestic and feral? Investigating the status of suids in the Romanian Gumelnița (5th mil. cal BC) with biogeochemistry and geometric morphometrics. *Journal of Anthropological Archaeology*, 42, 27–36. <https://doi.org/10.1016/j.jaa.2016.02.002>
- Bartosiewicz L, Gillis R, Girdland Flink L, Evin A, Cucchi T, Hoelzel R, Vidarsdottir U, Dobney K, Larson G, Schoop U-D (2013) Chalcolithic pig remains from Çamlıbel Tarlasi, Central Anatolia. *Archaeozoology of the Near East X. Proceedings of the Tenth International Symposium on the Archaeozoology of South-Western Asia and Adjacent Areas.*, 101–120.
- Battesti V, Gros-Balthazard M, Ogéron C, Ivorra S, Terral JF, Newton C (2018) Date Palm Agrobiodiversity (Phoenix dactylifera L.) in Siwa Oasis, Egypt: Combining Ethnography, Morphometry, and Genetics. *Human Ecology*, 46, 529–546. <https://doi.org/10.1007/s10745-018-0006-y>
- Bickler SH (2021) Machine Learning Arrives in Archaeology. *Advances in Archaeological Practice*, 9, 186–191. <https://doi.org/10.1017/aap.2021.6>
- Bignon O, Baylac M, Vigne J, Eisenmann V (2005) Geometric morphometrics and the population diversity of Late Glacial horses in Western Europe (I): phylogeographic and anthropological implications. *Journal of Archaeological Science*, 32, 375–391. <https://doi.org/10.1016/j.jas.2004.02.016>
- Bonhomme V, Forster E, Wallace M, Stillman E, Charles M, Jones G (2016) The first shoots of a modern morphometrics approach to the origins of agriculture. *Web Ecology*, 16, 1–2. <https://doi.org/10.5194/we-16-1-2016>
- Bonhomme V, Forster E, Wallace M, Stillman E, Charles M, Jones G (2017) Identification of inter- and intra-species variation in cereal grains through geometric morphometric analysis, and its resilience under experimental charring. *Journal of Archaeological Science*, 86, 60–67. <https://doi.org/10.1016/j.jas.2017.09.010>

- Bonhomme V, Picq S, Ivorra S, Evin A, Pastor T, Bacilieri R, Lacombe T, Figueiral I, Terral J-F, Bouby L (2020) Eco-evo-devo implications and archaeobiological perspectives of trait covariance in fruits of wild and domesticated grapevines (M Singh, Ed.). *PLOS ONE*, 15, e0239863. <https://doi.org/10.1371/journal.pone.0239863>
- Bonhomme V, Terral J-F, Zech-Matterne V, Ivorra S, Lacombe T, Deborde G, Kuchler P, Limier B, Pastor T, Rollet P, Bouby L (2021) Seed morphology uncovers 1500 years of vine agrobiodiversity before the advent of the Champagne wine. *Scientific Reports*, 11, 1–14. <https://doi.org/10.1038/s41598-021-81787-3>
- Bookstein FL (1991) *Morphometric tools for land- mark data: Geometry and biology*. Cambridge University Press, New York.
- Bopp-Ito M, Cucchi T, Evin A, Stopp B, Schibler J (2018) Phenotypic diversity in Bronze Age pigs from the Alpine and Central Plateau regions of Switzerland. *Journal of Archaeological Science: Reports*, 21, 38–46. <https://doi.org/10.1016/j.jasrep.2018.07.002>
- Bouby L, Bonhomme V, Ivorra S, Pastor T, Rovira N, Tillier M, Pagnoux C, Terral J-F (2018) Back from burn out: are experimentally charred grapevine pips too distorted to be characterized using morphometrics? *Archaeological and Anthropological Sciences*, 10, 943–954. <https://doi.org/10.1007/s12520-016-0425-x>
- Bouby L, Wales N, Jalabadze M, Rusishvili N, Bonhomme V, Ramos-Madrigal J, Evin A, Ivorra S, Lacombe T, Pagnoux C, Boaretto E, Gilbert MTP, Bacilieri R, Lordkipanidze D, Maghradze D (2021) Tracking the history of grapevine cultivation in Georgia by combining geometric morphometrics and ancient DNA. *Vegetation History and Archaeobotany*, 30, 63–76. <https://doi.org/10.1007/s00334-020-00803-0>
- Bourgeon O, Pagnoux C, Mauné S, Vargas EG, Ivorra S, Bonhomme V, Ater M, Moukhli A, Terral JF (2018) Olive tree varieties cultivated for the great Baetic oil trade between the 1st and the 4th centuries ad: morphometric analysis of olive stones from Las Delicias (Ecija, Province of Seville, Spain). *Vegetation History and Archaeobotany*, 27, 463–476. <https://doi.org/10.1007/s00334-017-0648-5>
- Burger P, Terral JF, Ruas MP, Ivorra S, Picq S (2011) Assessing past agrobiodiversity of *Prunus avium* L. (Rosaceae): A morphometric approach focussed on the stones from the archaeological site Hôtel-Dieu (16th century, Tours, France). *Vegetation History and Archaeobotany*, 20, 447–458. <https://doi.org/10.1007/s00334-011-0310-6>
- Ching T, Himmelstein DS, Beaulieu-Jones BK, Kalinin AA, Do BT, Way GP, Ferrero E, Agapow PM, Zietz M, Hoffman MM, Xie W, Rosen GL, Lengerich BJ, Israeli J, Lanchantin J, Woloszynek S, Carpenter AE, Shrikumar A, Xu J, Cofer EM, Lavender CA, Turaga SC, Alexandari AM, Lu Z, Harris DJ, Decaprio D, Qi Y, Kundaje A, Peng Y, Wiley LK, Segler MHS, Boca SM, Swamidass SJ, Huang A, Gitter A, Greene CS (2018) *Opportunities and obstacles for deep learning in biology and medicine*. <https://doi.org/10.1098/rsif.2017.0387>
- Chuang R, Bonhomme V (2019) Rethinking the dental morphological differences between domestic equids. *Journal of Archaeological Science*, 101, 140–148. <https://doi.org/10.1016/j.jas.2018.02.020>
- Clavel P, Dumoncel J, Der Sarkissian C, Seguin-Orlando A, Calvière-Tonasso L, Schiavinato S, Chauvey L, Perdureau A, Aury J-M, Wincker P, Onar V, Clavel B, Lepetz S, Braga J, Orlando L (2021) Assessing the predictive taxonomic power of the bony labyrinth 3D shape in horses, donkeys and their F1-hybrids. *Journal of Archaeological Science*, 131, 105383. <https://doi.org/10.1016/j.jas.2021.105383>
- Colominas L, Evin A, Burch J, Campmajó P, Casas J, Castanyer P, Carreras C, Guardia J, Olesti O, Pons E, Tremoleda J, Palet J-M (2019) Behind the steps of ancient sheep mobility in Iberia: new insights from a geometric morphometric approach. *Archaeological and Anthropological Sciences*, 11, 4971–4982. <https://doi.org/10.1007/s12520-019-00837-0>
- Cornette R, Herrel A, Stoetzel E, Moulin S, Hutterer R, Denys C, Baylac M (2015) Specific information levels in relation to fragmentation patterns of shrew mandibles: Do fragments tell the same story? *Journal of Archaeological Science*, 53, 323–330. <https://doi.org/10.1016/j.jas.2014.10.020>
- Cucchi T (2008) Uluburun shipwreck stowaway house mouse: molar shape analysis and indirect clues about the vessel's last journey. *Journal of Archaeological Science*, 35, 2953–2959. <https://doi.org/10.1016/j.jas.2008.06.016>
- Cucchi T, Barnett R, Martínková N, Renaud S, Renvoisé E, Evin A, Sheridan A, Mainland I, Wickham-Jones C, Tougaard C, Quéré JJP, Pascal MM, Heckel G, O'Higgins P, Searle J, Dobney K (2014) The changing pace of insular life: 5000 years of microevolution in the orkney vole (*Microtus arvalis orcadensis*). *Evolution*, 68, 2804–2820. <https://doi.org/10.1111/evo.12476>

- Cucchi T, Dai L, Balasse M, Zhao C, Gao J, Hu Y, Yuan J, Vigne JD (2016) Social complexification and pig (*Sus scrofa*) husbandry in ancient China: A combined geometric morphometric and isotopic approach. *PLoS ONE*, 11, 1–20. <https://doi.org/10.1371/journal.pone.0158523>
- Cucchi T, Domont A, Harbers H, Leduc C, Guidez A, Bridault A, Hongo H, Price M, Peters J, Briois F, Guilaine J, Vigne J-D (2021) Bones geometric morphometrics illustrate 10th millennium cal. BP domestication of autochthonous Cypriot wild boar (*Sus scrofa circeus nov. ssp.*). *Scientific Reports*, 11, 10–11. <https://doi.org/10.1038/s41598-021-90933-w>
- Cucchi T, Fujita M, Dobney K (2008) New insights into pig taxonomy, domestication and human dispersal in Island South East Asia: molar shape analysis of *Sus* remains from Niah Caves, Sarawak. *International Journal of Osteoarchaeology*, 19, 508–530. <https://doi.org/10.1002/oa>
- Cucchi T, Hulme-Beaman A, Yuan J, Dobney K (2011) Early Neolithic pig domestication at Jiahu, Henan Province, China: Clues from molar shape analyses using geometric morphometric approaches. *Journal of Archaeological Science*, 38, 11–22. <https://doi.org/10.1016/j.jas.2010.07.024>
- Cucchi T, Mohaseb A, Peigné S, Debue K, Orlando L, Mashkour M (2017) Detecting taxonomic and phylogenetic signals in equid cheek teeth: Towards new palaeontological and archaeological proxies. *Royal Society Open Science*, 4. <https://doi.org/10.1098/rsos.160997>
- Cucchi T, Stopp B, Schafberg R, Lesur J, Hassanin A, Schibler J (2019) Taxonomic and phylogenetic signals in bovine cheek teeth: Towards new biosystematic markers to explore the history of wild and domestic cattle. *Journal of Archaeological Science*, 109, 104993. <https://doi.org/10.1016/j.jas.2019.104993>
- Darwin C (1859) *The origin of species by means of natural selection, or, The preservation of favoured races in the struggle for life*. John Murray, London. <https://doi.org/10.5962/bhl.title.87916>
- Darwin C (1868) *The Variation of Animals and Plants under Domestication*. Cambridge University Press, Cambridge. <https://doi.org/10.1017/CBO9780511709517>
- Daza Perea A (2017) Preliminary Studies of Late Prehistoric Dog (*Canis lupus f. Familiaris* Linnaeus, 1758) Remains from the Iberian Peninsula: Osteometric and 2D Geometric Morphometric Approaches. *Papers from the Institute of Archaeology*, 27, 1–21. <https://doi.org/10.5334/pia-487>
- Devine J, Aponte JD, Katz DC, Liu W, Vercio LD Lo, Forkert ND, Marcucio R, Percival CJ, Hallgrímsson B (2020) A Registration and Deep Learning Approach to Automated Landmark Detection for Geometric Morphometrics. *Evolutionary Biology*, 47, 246–259. <https://doi.org/10.1007/s11692-020-09508-8>
- Dobney K, Cucchi T, Larson G (2008) The pigs of Island Southeast Asia and the Pacific: New evidence for taxonomic status and human-mediated dispersal. *Asian Perspectives*, 47, 59–74. <https://doi.org/10.1353/asi.2008.0009>
- Drake AG, Coquerelle M, Colombeau G (2015) 3D morphometric analysis of fossil canid skulls contradicts the suggested domestication of dogs during the late Paleolithic. *Scientific Reports*, 5, 8299. <https://doi.org/10.1038/srep08299>
- Drake AG, Coquerelle M, Kosintsev PA, Bachura OP, Sablin M, Gusev A V., Fleming LS, Losey RJ (2017) Three-Dimensional Geometric Morphometric Analysis of Fossil Canid Mandibles and Skulls. *Scientific Reports*, 7, 1–8. <https://doi.org/10.1038/s41598-017-10232-1>
- Drake AG, Klingenberg CP (2008) The pace of morphological change: historical transformation of skull shape in St Bernard dogs. *Proceedings. Biological sciences / The Royal Society*, 275, 71–76. <https://doi.org/10.1098/rspb.2007.1169>
- Duval C, Cucchi T, Horard-Herbin M-P, Lepetz S (2018) The development of new husbandry and economic models in Gaul between the Iron Age and the Roman Period: New insights from pig bones and teeth morphometrics. *Journal of Archaeological Science*, 99, 10–18. <https://doi.org/10.1016/j.jas.2018.08.016>
- Duval C, Lepetz S, Horard-Herbin MP, Cucchi T (2015) Did Romanization impact Gallic pig morphology? New insights from molar geometric morphometrics. *Journal of Archaeological Science*, 57, 345–354. <https://doi.org/10.1016/j.jas.2015.03.004>
- Evin A, Bonhomme V, Claude J (2020) Optimizing digitalization effort in morphometrics. *Biology Methods and Protocols*, 5, 1–10. <https://doi.org/10.1093/biomethods/bpaa023>
- Evin A, David L, Souron A, Mennecart B, Orliac M, Lebrun R (2022) Size and shape of the semicircular canal of the inner ear: A new marker of pig domestication? *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution*. <https://doi.org/10.1002/jez.b.23127>

- Evin A, Dobney K, Schafberg R, Owen J, Strand Vidarsdottir U, Larson G, Cucchi T (2015) Phenotype and animal domestication: A study of dental variation between domestic, wild, captive, hybrid and insular *Sus scrofa*. *BMC Evolutionary Biology*, 15, 6. <https://doi.org/10.1186/s12862-014-0269-x>
- Evin A, Flink LG, Bălăşescu A, Popovici D, Andreescu R, Bailey D, Mirea P, Lazăr C, Boroneanţ A, Bonsall C, Vidarsdottir US, Brehard S, Tresset A, Cucchi T, Larson G, Dobney K (2015) Unravelling the complexity of domestication: A case study using morphometrics and ancient DNA analyses of archaeological pigs from Romania. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 370, 20130616. <https://doi.org/10.1098/rstb.2013.0616>
- Figueiral I, Pomarède H, Court-Picon M, Bouby L, Tardy C, Terral JF (2015) *New insights into Mediterranean Gallo-Roman farming: a closer look at archaeological wells in Southern France*. <https://doi.org/10.1007/s12520-014-0181-8>
- Fiorentino G, Ferrio JP, Bogaard A, Araus JL, Riehl S (2015) Stable isotopes in archaeobotanical research. *Vegetation History and Archaeobotany*, 24, 215–227. <https://doi.org/10.1007/s00334-014-0492-9>
- Fisher AE (2019) When is a wolf a dog? Combined geometric morphometrics and stable isotope analyses for differentiating wild from domestic canids on the Northern Plains. *Plains Anthropologist*, 64, 316–349. <https://doi.org/10.1080/00320447.2018.1548064>
- Flowers JM, Hazzouri KM, Gros-Balthazard M, Mo Z, Koutroumpa K, Perrakis A, Ferrand S, Khierallah HSM, Fuller DQ, Aberlenc F, Fournaraki C, Purugganan MD (2019) Cross-species hybridization and the origin of North African date palms. *Proceedings of the National Academy of Sciences of the United States of America*, 116, 1651–1658. <https://doi.org/10.1073/pnas.1817453116>
- Foster A (2018) Identifying chicken breeds in the archaeological record: a geometric and linear morphometric approach. Cambridge University Press, Cambridge. <https://doi.org/10.1017/CBO9781107415324.004>
- Frantz L, Haile J, Lin A, Scheu A, Geörg C, Benecke N, Alexander M, Linderholm A, Mullin V, Daly KG, Battista V, Price M, Gron KJ, Alexandri P, Arbogast RM, Arbuckle B, Bălăşescu A, Barnett R, Bartosiewicz L, Baryshnikov G, Bonsall C, Borić D, Boroneanţ A, Bulatović J, Çakırlar C, Carretero JM, Chapman J, Church M, Crooijmans R, De Cupere B, Detry C, Dimitrijevic V, Dumitrascu V, Du Plessis L, Edwards CJ, Ereik CM, Erim-Özdoğan A, Ervynck A, Fulgione D, Gligor M, Götherström A, Gourichon L, Groenen MAM, Helmer D, Hongo H, Horwitz LK, Irving-Pease EK, Lebrasseur O, Lesur J, Malone C, Manaseryan N, Marciniak A, Martlew H, Mashkour M, Matthews R, Matuzeviciute GM, Maziar S, Meijaard E, McGovern T, Megens HJ, Miller R, Mohaseb AF, Orschiedt J, Orton D, Papathanasiou A, Pearson MP, Pinhasi R, Radmanović D, Ricaut F-X, Richards M, Sabin R, Sarti L, Schier W, Sheikhi S, Stephan E, Stewart JR, Stoddart S, Tagliacozzo A, Tasić N, Trantalidou K, Tresset A, Valdiosera C, Van Den Hurk Y, Van Poucke S, Vigne J-D, Yanevich A, Zeeb-Lanz A, Triantafyllidis A, Gilbert MTP, Schibler J, Rowley-Conwy P, Zeder M, Peters J, Cucchi T, Bradley DG, Dobney K, Burger J, Evin A, Girdland-Flink L, Larson G (2019) Ancient pigs reveal a near-complete genomic turnover following their introduction to Europe. *Proceedings of the National Academy of Sciences of the United States of America*, 116, 17231–17238. <https://doi.org/10.1073/pnas.1901169116>
- Frémondeau D, De Cupere B, Evin A, Van Neer W (2017) Diversity in pig husbandry from the Classical-Hellenistic to the Byzantine periods: An integrated dental analysis of Düzen Tepe and Sagalassos assemblages (Turkey). *Journal of Archaeological Science: Reports*, 11, 38–52. <https://doi.org/10.1016/j.jasrep.2016.11.030>
- García-Granero JJ, Arias-Martorell J, Madella M, Lancelotti C (2016) Geometric morphometric analysis of *Setaria italica* (L.) P. Beauv. (foxtail millet) and *Brachiaria ramosa* (L.) Stapf. (browntop millet) and its implications for understanding the biogeography of small millets. *Vegetation History and Archaeobotany*, 25, 303–310. <https://doi.org/10.1007/s00334-015-0541-z>
- Geiger M, Evin A, Sánchez-Villagra MR, Gascho D, Mainini C, Zollikofer CPE (2017) Neomorphosis and heterochrony of skull shape in dog domestication. *Scientific Reports*, 7, 13443. <https://doi.org/10.1038/s41598-017-12582-2>
- Geiger M, Sánchez-Villagra MR (2018) Similar rates of morphological evolution in domesticated and wild pigs and dogs. *Frontiers in Zoology*, 15, 23. <https://doi.org/10.1186/s12983-018-0265-x>
- Giardina CR, Kuhl FP (1977) Accuracy of Curve Approximation By Harmonically Related Vectors With Elliptical Loci. *Comput Graphics Image Process*, 6, 277–285. [https://doi.org/10.1016/S0146-664X\(77\)80029-4](https://doi.org/10.1016/S0146-664X(77)80029-4)
- Grasso AM, Mavelli F, Fiorentino G (2018) Quantitative evaluation of modern Citrus seed shape and comparison with archaeological remains discovered in Pompeii and Rome. In: *AGRUMED: Archaeology and history of citrus fruit in the Mediterranean*. <https://doi.org/10.4000/books.pcbj.2211>

- Gros-Balthazard M, Galimberti M, Kousathanas A, Newton C, Ivorra S, Paradis L, Vigouroux Y, Carter R, Tengberg M, Battesti V, Santoni S, Falquet L, Pintaud JC, Terral JF, Wegmann D (2017) The Discovery of Wild Date Palms in Oman Reveals a Complex Domestication History Involving Centers in the Middle East and Africa. *Current Biology*, 27, 2211–2218.e8. <https://doi.org/10.1016/j.cub.2017.06.045>
- Gros-Balthazard M, Newton C, Ivorra S, Pierre MH, Pintaud JC, Terral JF (2016) The domestication syndrome in Phoenix dactylifera seeds: Toward the identification of wild date palm populations. *PLoS ONE*, 11, 1–21. <https://doi.org/10.1371/journal.pone.0152394>
- Hanot P, Bayarsaikhan J, Guintard C, Haruda A, Mijiddorj E, Schafberg R, Taylor W (2021) Cranial shape diversification in horses: variation and covariation patterns under the impact of artificial selection. *BMC Ecology and Evolution*, 21, 1–19. <https://doi.org/10.1186/s12862-021-01907-5>
- Hanot P, Guintard C, Lepetz S, Cornette R (2017) Identifying domestic horses, donkeys and hybrids from archaeological deposits: A 3D morphological investigation on skeletons. *Journal of Archaeological Science*, 78, 88–98. <https://doi.org/10.1016/j.jas.2016.12.002>
- Harbers H, Zanolli C, Cazenave M, Theil JC, Ortiz K, Blanc B, Locatelli Y, Schafberg R, Lecompte F, Baly I, Laurens F, Callou C, Herrel A, Puymeraul L, Cucchi T (2020) Investigating the impact of captivity and domestication on limb bone cortical morphology: an experimental approach using a wild boar model. *Scientific Reports*, 10, 1–13. <https://doi.org/10.1038/s41598-020-75496-6>
- Harjunmaa E, Kallonen A, Voutilainen M, Hämäläinen K, Mikkola ML, Jernvall J (2012) On the difficulty of increasing dental complexity. *Nature*, 483, 324–327. <https://doi.org/10.1038/nature10876>
- Haruda A (2017) Separating Sheep (Ovis aries L.) and Goats (Capra hircus L.) Using Geometric Morphometric Methods: An Investigation of Astragalus Morphology from Late and Final Bronze Age Central Asian Contexts. *International Journal of Osteoarchaeology*, 27, 551–562. <https://doi.org/10.1002/oa.2576>
- Haruda A, Varfolomeev V, Goriachev A, Yermolayeva A, Outram AK (2019) A new zooarchaeological application for geometric morphometric methods: Distinguishing Ovis aries morphotypes to address connectivity and mobility of prehistoric Central Asian pastoralists. *Journal of Archaeological Science*, 107, 50–57. <https://doi.org/10.1016/j.jas.2019.05.002>
- Hassoun S, Jefferson F, Shi X, Stucky B, Wang J, Rosa E (2021) Artificial Intelligence for Biology. *Integrative and Comparative Biology*. <https://doi.org/10.1093/icb/icab188>
- Hernández A, L'Heureux GL, Leoni JB (2021) Guanaco hunting and Llama herding in the South-Central Andes (3000–900 BP): An osteomorphometrical approach. *Journal of Archaeological Science: Reports*, 37. <https://doi.org/10.1016/j.jasrep.2021.102952>
- Horn C, Ivarsson O, Lindhé C, Potter R, Green A, Ling J (2021) Artificial Intelligence, 3D Documentation, and Rock Art—Approaching and Reflecting on the Automation of Identification and Classification of Rock Art Images. *Journal of Archaeological Method and Theory*. <https://doi.org/10.1007/s10816-021-09518-6>
- Houle D, Govindaraju DR, Omholt S (2010) Phenomics: The next challenge. *Nature Reviews Genetics*, 11, 855–866. <https://doi.org/10.1038/nrg2897>
- James Rohlf F, Marcus LF (1993) A revolution morphometrics. *Trends in Ecology & Evolution*, 8, 129–132. [https://doi.org/10.1016/0169-5347\(93\)90024-J](https://doi.org/10.1016/0169-5347(93)90024-J)
- Jesus A, Bonhomme V, Evin A, Ivorra S, Soteras R, Salavert A, Antolín F, Bouby L (2021) A morphometric approach to track opium poppy domestication. *Scientific Reports*, 11, 1–11. <https://doi.org/10.1038/s41598-021-88964-4>
- Krause-Kyora B, Makarewicz C, Evin A, Flink LGLG, Dobney K, Larson G, Hartz S, Schreiber S, Von Carnap-Bornheim C, Wurmb-Schwark N, Nebel A (2013) Use of domesticated pigs by Mesolithic hunter-gatherers in northwestern Europe. *Nature Communications*, 4, 1–7. <https://doi.org/10.1038/ncomms3348>
- Kuhl FP, Giardina CR (1982) Elliptic Fourier features of a closed contour. *Computer Graphics and Image Processing*, 18, 236–258. [https://doi.org/10.1016/0146-664X\(82\)90034-X](https://doi.org/10.1016/0146-664X(82)90034-X)
- Manin A, Evin A (2020) Canis spp. identification in central Mexico and its archaeological implications : toward a better understanding of the ecology and the cultural role of canids in ancient Mesoamerica. In: *Relations hommes – canidés de la Préhistoire aux périodes modernes* , pp. 95–114. UN@ Éditions. <https://doi.org/10.46608/DANA3.9782381490120.6>
- Margaritis E, Pagnoux C, Bouby L, Bonhomme V, Ivorra S, Tsirtsis K, Terral JF (2021) Hellenistic grape and olive diversity: A case study from rural estates in Greece. *Journal of Archaeological Science: Reports*, 38, 102842. <https://doi.org/10.1016/j.jasrep.2021.102842>

- Mariotti Lippi M, Mori Secci M, Giachi G, Bouby L, Terral J-F, Castiglioni E, Cottini M, Rottoli M, de Grummond NT (2020) Plant remains in an Etruscan-Roman well at Cetamura del Chianti, Italy. *Archaeological and Anthropological Sciences*, 12, 35. <https://doi.org/10.1007/s12520-019-00992-4>
- Marom N, Meiri M, Tepper Y, Erickson-Gini T, Reshef H, Weissbrod L, Bar-Oz G (2019) Zooarchaeology of the social and economic upheavals in the Late Antique-Early Islamic sequence of the Negev Desert. *Scientific Reports*, 9, 6702. <https://doi.org/10.1038/s41598-019-43169-8>
- Miele V, Dussert G, Cucchi T, Renaud S (2020) Deep learning for species identification of modern and fossil rodent molars. In: *bioRxiv*, pp. 1–27. <https://doi.org/10.1101/2020.08.20.259176>
- Moricca C, Bouby L, Bonhomme V, Ivorra S, Pérez-Jordà G, Nigro L, Spagnoli F, Peña-Chocarro L, van Dommelen P, Sadori L (2021) Grapes and vines of the Phoenicians: Morphometric analyses of pips from modern varieties and Iron Age archaeological sites in the Western Mediterranean. *Journal of Archaeological Science: Reports*, 37. <https://doi.org/10.1016/j.jasrep.2021.102991>
- Myles S, Boyko AR, Owens CL, Brown PJ, Grassi F, Aradhya MK, Prins B, Reynolds A, Chia J-M, Ware D, Bustamante CD, Buckler ES (2011) Genetic structure and domestication history of the grape. *Proceedings of the National Academy of Sciences*, 108, 3530–3535. <https://doi.org/10.1073/pnas.1009363108>
- Neaux D, Blanc B, Ortiz K, Locatelli Y, Laurens F, Baly I, Callou C, Lecompte F, Cornette R, Sansalone G, Haruda A, Schafberg R, Vigne JD, Debat V, Herrel A, Cucchi T (2020) How Changes in Functional Demands Associated with Captivity Affect the Skull Shape of a Wild Boar (*Sus scrofa*). *Evolutionary Biology*. <https://doi.org/10.1007/s11692-020-09521-x>
- Newton C, Lorre C, Sauvage C, Ivorra S, Terral JF (2014) On the origins and spread of *Olea europaea* L. (olive) domestication: Evidence for shape variation of olive stones at Ugarit, Late Bronze Age, Syria—a window on the Mediterranean Basin and on the westward diffusion of olive varieties. *Vegetation History and Archaeobotany*, 23, 567–575. <https://doi.org/10.1007/s00334-013-0412-4>
- Newton C, Terral JF, Ivorra S (2006) The Egyptian olive (*Olea europaea* subsp. *europaea*) in the later first millennium BC: Origins and history using the morphometric analysis of olive stones. *Antiquity*, 80, 405–414. <https://doi.org/10.1017/S0003598X00093716>
- Nistelberger HM, Smith O, Wales N, Star B, Boessenkool S (2016) The efficacy of high-throughput sequencing and target enrichment on charred archaeobotanical remains. *Scientific Reports*, 6, 37347. <https://doi.org/10.1038/srep37347>
- Orrù M, Grillo O, Lovicu G, Venora G, Bacchetta G (2013) Morphological characterisation of *Vitis vinifera* L. seeds by image analysis and comparison with archaeological remains. *Vegetation History and Archaeobotany*, 22, 231–242. <https://doi.org/10.1007/s00334-012-0362-2>
- Ottoni C, Girdland Flink L, Evin A, Georg C, De Cupere B, Van Neer W, Bartosiewicz L, Linderholm A, Barnett R, Peters J, Decorte R, Waelkens M, Vanderheyden N, Ricaut FX, Çakırlar C, Çevik Ö, Hoelzel AR, Mashkour M, Mohaseb Karimlu AF, Sheikhi Seno S, Daujat J, Brock F, Pinhasi R, Hongo H, Perez-Enciso M, Rasmussen M, Frantz L, Megens HJ, Crooijmans R, Groenen M, Arbuckle B, Benecke N, Strand Vidarsdottir U, Burger J, Cucchi T, Dobney K, Larson G (2013) Pig domestication and human-mediated dispersal in western Eurasia revealed through ancient DNA and geometric morphometrics. *Molecular Biology and Evolution*, 30, 824–832. <https://doi.org/10.1093/molbev/mss261>
- Owen J, Dobney K, Evin A, Cucchi T, Larson G, Strand Vidarsdottir U (2014) The zooarchaeological application of quantifying cranial shape differences in wild boar and domestic pigs (*Sus scrofa*) using 3D geometric morphometrics. *Journal of Archaeological Science*, 43, 159–167. <https://doi.org/10.1016/j.jas.2013.12.010>
- Pagnoux C, Bouby L, Ivorra S, Petit C, Valamoti S-M, Pastor T, Picq S, Terral J-F (2015) Inferring the agrobiodiversity of *Vitis vinifera* L. (grapevine) in ancient Greece by comparative shape analysis of archaeological and modern seeds. *Vegetation History and Archaeobotany*, 24, 75–84. <https://doi.org/10.1007/s00334-014-0482-y>
- Pagnoux C, Bouby L, Valamoti SM, Bonhomme V, Ivorra S, Gkatzogia E, Karathanou A, Kotsachristou D, Kroll H, Terral JF (2021) Local domestication or diffusion? Insights into viticulture in Greece from Neolithic to Archaic times, using geometric morphometric analyses of archaeological grape seeds. *Journal of Archaeological Science*, 125. <https://doi.org/10.1016/j.jas.2020.105263>
- Pelletier M (2019) Morphological diversity of wild rabbit populations: implications for archaeology and palaeontology. *Biological Journal of the Linnean Society*, 128, 211–224. <https://doi.org/10.1093/biolinnean/blz074>

- Pelletier M, Kotiaho A, Niinimäki S, Salmi AK (2020) Identifying early stages of reindeer domestication in the archaeological record: a 3D morphological investigation on forelimb bones of modern populations from Fennoscandia. *Archaeological and Anthropological Sciences*, 12. <https://doi.org/10.1007/s12520-020-01123-0>
- Pelletier M, Kotiaho A, Niinimäki S, Salmi AK (2021) Impact of selection and domestication on hindlimb bones of modern reindeer populations: Archaeological implications for early reindeer management by Sámi in Fennoscandia. *Historical Biology*, 00, 1–19. <https://doi.org/10.1080/08912963.2021.1947268>
- Pilot M, Greco C, vonHoldt BM, Randi E, Jędrzejewski W, Sidorovich VE, Konopiński MK, Ostrander EA, Wayne RK (2018) Widespread, long-term admixture between grey wolves and domestic dogs across Eurasia and its implications for the conservation status of hybrids. *Evolutionary Applications*, 11, 662–680. <https://doi.org/10.1111/eva.12595>
- Pöllath N, Alibert P, Schafberg R, Peters J (2018) Striking new paths – Distinguishing ancient *Ovis orientalis* from its modern domestic descendant (Karakul breed) applying Geometric and traditional Morphometric approaches to the astragalus. In: *Archaeozoology of the Near East XII. Proceedings of the 12th International Symposium of the ICAZ Archaeozoology of Southwest Asia and Adjacent Areas Working Group, Groningen Institute of Archaeology, June 14-15 2015, University of Groningen, the Netherlands* (eds Çakırlar C, Chahoud J, Berthon R, Pilaar Birch SE), pp. 207–225. Barkhuis Publishing & University of Groningen. <https://doi.org/10.2307/j.ctvggx2m4>
- Pöllath N, Schafberg R, Peters J (2019) Astragalar morphology: Approaching the cultural trajectories of wild and domestic sheep applying Geometric Morphometrics. *Journal of Archaeological Science: Reports*, 23, 810–821. <https://doi.org/10.1016/j.jasrep.2018.12.004>
- Price M, Evin A (2017) Long-term morphological changes and evolving human-pig relations in the northern Fertile Crescent from 11,000 to 2000 cal. bc. *Archaeological and Anthropological Sciences*, 11, 237–251. <https://doi.org/10.1007/s12520-017-0536-z>
- Riaz S, De Lorenzis G, Velasco D, Koehmstedt A, Maghradze D, Bobokashvili Z, Musayev M, Zdunic G, Laucou V, Andrew Walker M, Failla O, Preece JE, Aradhya M, Arroyo-Garcia R (2018) Genetic diversity analysis of cultivated and wild grapevine (*Vitis vinifera* L.) accessions around the Mediterranean basin and Central Asia. *BMC Plant Biology*, 18, 1–14. <https://doi.org/10.1186/s12870-018-1351-0>
- Rohlf J (1990) Fitting curves to outlines. In: *Proceedings of the Michigan Morphometrics Workshop* (eds Rohlf JF, Bookstein FL), pp. 167–177. University of Michigan Museum of Zoology Special Publication 2.
- Ros J, Evin A, Bouby L, Ruas M-P (2014) Geometric morphometric analysis of grain shape and the identification of two-rowed barley (*Hordeum vulgare* subsp. *distichum* L.) in southern France. *Journal of Archaeological Science*, 41, 568–575. <https://doi.org/10.1016/j.jas.2013.09.015>
- Sabato D, Esteras C, Grillo O, Peña-Chocarro L, Leida C, Uccchesu M, Usai A, Bacchetta G, Picó B (2019) Molecular and morphological characterisation of the oldest *Cucumis melo* L. seeds found in the Western Mediterranean Basin. *Archaeological and Anthropological Sciences*, 11, 789–810. <https://doi.org/10.1007/s12520-017-0560-z>
- Sallon S, Cherif E, Chabrilange N, Solowey E, Gros-Balthazard M, Ivorra S, Terral JF, Egli M, Aberlenc F (2020) Origins and insights into the historic Judean date palm based on genetic analysis of germinated ancient seeds and morphometric studies. *Science Advances*, 6, 1–11. <https://doi.org/10.1126/sciadv.aax0384>
- Schoenebeck JJ, Hamilton-Dyer S, Baxter IL, Schwarz T, Nussbaumer M (2021) From head to hind: Elucidating function through contrasting morphometrics of ancient and modern pedigree dogs. *Anatomical Record*, 304, 63–77. <https://doi.org/10.1002/ar.24412>
- Schoenebeck JJ, Ostrander EA (2013) The Genetics of Canine Skull Shape Variation. *Genetics*, 193, 317–325. <https://doi.org/10.1534/genetics.112.145284>
- Seetah K, Cardini A, Barker G (2016) A ‘long-fuse domestication’ of the horse? Tooth shape suggests explosive change in modern breeds compared with extinct populations and living Przewalski’s horses. *The Holocene*, 26, 1326–1333. <https://doi.org/10.1177/0959683616638436>
- Tarongi M, Bonhomme V, Evin A, Ivorra S, López D, Alonso N, Bouby L (2020) A new way of seeing pulses: preliminary results of geometric morphometric analyses of Iron Age seeds from the site of La Font de la Canya (Catalonia, Spain). *Vegetation History and Archaeobotany*. <https://doi.org/10.1007/s00334-020-00801-2>
- Terral J-F, Alonso N, Buxó I, Capdevila R, Chatti N, Fabre L, Fiorentino G, Marínval P, Jordá GP, Pradat B, Rovira N, Alibert P (2004) Historical biogeography of olive domestication (*Olea europaea* L.) as revealed by

- geometrical morphometry applied to biological and archaeological material. *Journal of Biogeography*, 31, 63–77. <https://doi.org/10.1046/j.0305-0270.2003.01019.x>
- Terral J-F, Bonhomme V, Pagnoux C, Ivorra S, Newton C, Paradis L, Ater M, Kassoul J, Limier B, Bouby L, Cornet F, Barazani O, Galili E (2021) Shape Diversity of Olive Stone, Resulting from Domestication and Diversification, Unveils Traits of the Oldest Known, 7000-Years-Old Table Olives from Hishuley Carmel Site (Israel). *agronomy*, 11, 2187. <https://doi.org/10.3390/agronomy11112187>
- Terral J-F, Newton C, Ivorra S, Gros-Balthazard M, de Morais CT, Picq S, Tengberg M, Pintaud JC (2012) Insights into the historical biogeography of the date palm (*Phoenix dactylifera* L.) using geometric morphometry of modern and ancient seeds. *Journal of Biogeography*, 39, 929–941. <https://doi.org/10.1111/j.1365-2699.2011.02649.x>
- Terral J-F, Tabard E, Bouby L, Ivorra S, Pastor T, Figueiral I, Picq S, Chevance J-B, Jung C, Fabre L, Tardy C, Compan M, Bacilieri R, Lacombe T, This P (2010) Evolution and history of grapevine (*Vitis vinifera*) under domestication: new morphometric perspectives to understand seed domestication syndrome and reveal origins of ancient European cultivars. *Annals of botany*, 105, 443–455. <https://doi.org/10.1093/aob/mcp298>
- Ucchesu M, Orrù M, Grillo O, Venora G, Paglietti G, Ardu A, Bacchetta G (2016) Predictive method for correct identification of archaeological charred grape seeds: Support for advances in knowledge of grape domestication process. *PLoS ONE*, 11, 1–18. <https://doi.org/10.1371/journal.pone.0149814>
- Ucchesu M, Orrù M, Grillo O, Venora G, Usai A, Serreli PF, Bacchetta G (2015) Earliest evidence of a primitive cultivar of *Vitis vinifera* L. during the Bronze Age in Sardinia (Italy). *Vegetation History and Archaeobotany*, 24, 587–600. <https://doi.org/10.1007/s00334-014-0512-9>
- Valamoti SM, Pagnoux C, Ntinou M, Bouby L, Bonhomme V, Terral JF (2020) More than meets the eye: new archaeobotanical evidence on Bronze Age viticulture and wine making in the Peloponnese, Greece. *Vegetation History and Archaeobotany*, 29, 35–50. <https://doi.org/10.1007/s00334-019-00733-6>
- Valenzuela-Lamas S, Baylac M, Cucchi T, Vigne J-D (2011) House mouse dispersal in Iron Age Spain: A geometric morphometrics appraisal. *Biological Journal of the Linnean Society*, 102, 483–497. <https://doi.org/10.1111/j.1095-8312.2010.01603.x>
- Vigne J-D, Evin A, Cucchi T, Dai L, Yu C, Hu S, Soulages N, Wang W, Sun Z, Gao J, Dobney K, Yuan J (2016) Earliest “domestic” cats in China identified as leopard cat (*Prionailurus bengalensis*). *PLoS ONE*, 11. <https://doi.org/10.1371/journal.pone.0147295>
- Wolcott KA, Chomicki G, Staedler YM, Wasylukowa K, Nesbitt M, Schönenberger J, Renner SS (2021) Three-dimensional X-ray-computed tomography of 3300- to 6000-year-old *Citrullus* seeds from Libya and Egypt compared to extant seeds throws doubts on species assignments. *Plants People Planet*, 3, 694–702. <https://doi.org/10.1002/ppp3.10220>