DSCAML1 is differentially expressed in ovarian cancers.

Shahan Mamoor, MS¹

¹Thomas Jefferson School of Law

San Diego, CA 92901

mamoorsk@tjisl.edu

Ovarian cancer is the fifth leading cause of cancer death for women in the United States and the leading cause of death from a gynecologic cancer for women in the developed world (1, 2). We compared the transcriptional profiles of benign ovarian tissues to that ovarian tumors isolated from women diagnosed with ovarian cancer using published datasets (3, 4) and found that the Down Syndrome cell adhesion molecule-like 1, or DSCAML1 (5) was one of the genes whose expression was most different between ovarian tumor and healthy ovarian tissue using separate datasets. This was consistent between two independent datasets analyzed. This is the first report of differential expression of DSCAML1 in ovarian cancers. DSCAML1 should be evaluated for its ability to initiate, or maintain the tumorigenic state in human ovarian cancer.

Keywords: ovarian cancer, differential gene expression, tumor transcriptome analysis, systems oncology, DSCAML1, targeted therapeutics in ovarian cancer.

Introduction

Ovarian cancer was diagnosed in over 24,000 women in the United States in 2018 and in the same year greater than 17,000 women expired from this disease (2). It is the most common reason why women die from a gynecologic cancer in the developed world (1). There are zero targeted treatments available for women diagnosed with ovarian cancer.

Understanding the most significant differences between the gene expression profiles of ovarian cancers as compared to the normal ovary can provide critical insights into the genes that play an essential role in the process of becoming and staying an ovarian tumor.

In this study, we compared the transcriptomes of ovarian tumors with that of normal ovarian tissue using two separate datasets (3, 4). We found that the Down Syndrome cell adhesion molecule-like 1 DSCAML1 (5) was among the genes most differentially expressed in ovarian cancers. DSCAML1 should be evaluated for its ability to support ovarian tumor growth in xenograft mouse models of ovarian cancer *in vivo* (6).

Methods

We utilized datasets GSE119056 (3) ($n=3$ for control and $n=6$ for tumor) and GSE54388 (4) ($n=6$ for control and $n=16$ for tumor) to perform this analysis in conjunction with GEO2R. The authors of GSE119056 used Agilent 067406 CBC lncRNA + mRNA microarray V4.0 technology to generate the microarray dataset utilized here. The authors of GSE54388 used Affymetrix Human Genome U133 Plus 2.0 array technology to generate the microarray dataset also utilized here. The Benjamini and Hochberg method of p-value adjustment was utilized, and p-value adjustment was set to “Auto-detect”. The NCBI generated category of platform annotation was utilized. To evaluate the statistical significance of differences in RNA expression values in control (normal, benign) versus tumor groups, a two-tailed, unpaired t test with Welch’s correction was performed to obtain a p-value. P-values less than 0.05 were considered statistically significant.

Results

We performed whole transcriptome analysis of ovarian cancers by comparing the transcriptome of ovarian tumors isolated from patients diagnosed with ovarian cancer with that of the transcriptome of the normal ovary with non-affected, control ovarian tissue samples using two independent datasets (3, 4).

DSCAML1 is differentially expressed in ovarian tumors compared to healthy, non-affected ovarian tissue.

We identified DSCAML1 as one of the most differentially expressed genes in the ovarian tumor transcriptome. DSCAML1 was ranked 143 with respect to its change in expression between tumor and benign ovarian tissue groups, out of 65534 total transcripts. This was statistically significant (Table 1; $p=2.75E-06$).

We queried a separate dataset that possessed transcriptome information on both ovarian tumors and control, non-affected ovarian tissue. In this dataset, DSCAML1 was ranked 141 in terms of differential expression between ovarian tumor and normal ovarian tissue groups, out of 54675 total transcripts. This was also statistically significant (Table 1; $p=1.02E-09$).

DSCAML1 is expressed at significantly lower levels in ovarian tumors than in healthy ovarian tissue

Next we extracted the exact mRNA expression values of DSCAML1 for both the ovarian tumors and benign ovarian tissue, from both datasets. We found that across both datasets, using unique transcriptome data from separate patient cohorts, that DSCAML1 was expressed at lower levels in ovarian tumors than in healthy ovarian tissue. This was statistically significant in both cases (Figure 1; GSE11254, $p<0.0001$; GSE54388; $p=0.0002$).

Thus, we identified DSCAML1 as differentially expressed in the tumors of women diagnosed with ovarian cancer and found that the expression of DSCAML1 was significantly lower in ovarian tumors than in normal ovary.

Discussion

Ovarian cancer is a significant medical problem as the fifth leading cause of cancer death for women in the United States. Ninety percent of women diagnosed with ovarian cancer possess an epithelial type of cancer, and the most common epithelial ovarian cancer is the serous type. The five-year survival rate for women with stage III or stage IV serous ovarian cancer is 42% and 26%, respectively. There are zero targeted treatments available for women diagnosed with ovarian cancer. Towards this goal, we performed systems-level transcriptome analysis of ovarian tumors as compared to the normal ovary using two independent datasets. We found that across datasets, DSCAML1 was among the genes whose expression was most significantly different between ovarian tumor and the normal ovary, and that it was expressed at significantly lower levels in ovarian tumors than in the normal ovary.

DSCAML1 is a cell adhesion molecule involved in synaptic adhesion, axon guidance, and self-avoidance in neurodevelopment (7). This is the first report of differential expression of DSCAML1 in ovarian cancers. Whether DSCAML1 can support tumor initiation, tumor growth or tumor maintenance in vivo should be assessed using xenograft mouse models of ovarian cancer (6). Manipulation of DSCAML1 expression or activity may represent a novel therapeutic strategy in ovarian cancer.

References

1. Chornokur, G., Amankwah, E.K., Schildkraut, J.M. and Phelan, C.M., 2013. Global ovarian cancer health disparities. *Gynecologic oncology*, 129(1), pp.258-264.
2. Torre, L.A., Trabert, B., DeSantis, C.E., Miller, K.D., Samimi, G., Runowicz, C.D., Gaudet, M.M., Jemal, A. and Siegel, R.L., 2018. Ovarian cancer statistics, 2018. *CA: a cancer journal for clinicians*, 68(4), pp.284-296.
3. Dong, S., Wang, R., Wang, H., Ding, Q., Zhou, X., Wang, J., Zhang, K., Long, Y., Lu, S., Hong, T. and Ren, H., 2019. HOXD-AS1 promotes the epithelial to mesenchymal transition of ovarian cancer cells by regulating miR-186-5p and PIK3R3. *Journal of Experimental & Clinical Cancer Research*, 38(1), p.110.
4. Yeung, T.L., Leung, C.S., Wong, K.K., Gutierrez-Hartmann, A., Kwong, J., Gershenson, D.M. and Mok, S.C., 2017. ELF3 is a negative regulator of epithelial-mesenchymal transition in ovarian cancer cells. *Oncotarget*, 8(10), p.16951.
5. Agarwala, K.L., Ganesh, S., Tsutsumi, Y., Suzuki, T., Amano, K. and Yamakawa, K., 2001. Cloning and functional characterization of DSCAML1, a novel DSCAM-like cell adhesion molecule that mediates homophilic intercellular adhesion. *Biochemical and biophysical research communications*, 285(3), pp.760-772.
6. Pizer, E.S., Wood, F.D., Heine, H.S., Romantsev, F.E., Pasternack, G.R. and Kuhajda, F.P., 1996. Inhibition of fatty acid synthesis delays disease progression in a xenograft model of ovarian cancer. *Cancer research*, 56(6), pp.1189-1193.
7. Fuerst, P.G., Bruce, F., Tian, M., Wei, W., Elstrott, J., Feller, M.B., Erskine, L., Singer, J.H. and Burgess, R.W., 2009. DSCAM and DSCAML1 function in self-avoidance in multiple cell types in the developing mouse retina. *Neuron*, 64(4), pp.484-497.

Table 1

Rank (dataset)	ID	p-value	t	B	Gene / Transcript
GSE119056					
143	A_23_P47340	2.75E-06	-9.2062356	5.17053	NM_020693
GSE54388					
141	232059_at	1.02E-09	-10.0232456	12.439732	DSCAML1

Table 1: DSCAML1 is differentially expressed in ovarian tumors compared to healthy, non-affected ovarian tissue.

The rank of differential expression globally out of 65534 transcripts (GSE1190560) or 54675 transcripts (GSE54388), probe ID, the p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between tumor and control groups, the log fold-change in expression between tumor and control groups, and gene/transcript symbol or Genbank accession information are shown in this chart (extracted from GEO2R).

Figure 1: DSCAML1 is expressed at significantly lower levels in ovarian tumors than in healthy ovarian tissue.

The expression of DSCAML1 in benign ovarian tissues (green) and ovarian cancers (tumor tissues, red) is shown by bar graph with the error bars denoting the standard error of the mean (left) and by a violin plot representing minimum to maximum expression with the dotted horizontal line denoting the median value and the finer lines denoting the 25th and 75th percent quartiles of expression (right). Data from GSE119056 is shown in the top row, while data from GSE54388 is shown in the bottom row. A statistical test was performed to evaluate the significance of expression between tumor (ovarian cancer) and control (ovary) groups, and the results of those tests, a p-value, is shown on each graph.