Title: A Guilt-Free Strategy Increase Self-Reported Non-Compliance with COVID-19 Preventive Measures: Experimental Evidence from 12 Countries

Authors

Daoust, Jean-François^{1*}, Éric Bélanger², Ruth Dassonneville³, Erick Lachapelle³, Richard Nadeau³, Michael Becher⁴, Sylvain Brouard⁵, Martial Foucault⁵, Christoph Hönnige⁶, Daniel Stegmüller⁷

¹ *University of Edinburgh;*

² *McGill University;*

³ *Université de Montréal;*

⁴ *IE University;*

⁵ *Sciences Po – CEVIPOF;*

⁶ *University of Hanover*

⁷ *Duke University*

* *Corresponding author:* jf.daoust@ed.ac.uk

Chrystal Macmillan Building

15a George Square

University of Edinburgh

Edinburgh

EH8 9LD

Word count: 4,892 words

Acknowledgement

The data were collected for the project "Citizens' Attitudes Under COVID-19 Pandemic" by the following research team: Sylvain Brouard (Sciences Po – CEVIPOF & LIEPP), Michael Becher (Institute for Advanced Studies in Toulouse & Université Toulouse Capitole 1), Martial Foucault (Sciences Po – CEVIPOF), Pavlos Vasilopoulos (University of York), Vincenzo Galasso (Bocconi University), Christoph Hönnige (University of Hanover), Eric Kerrouche (Sciences Po – CEVIPOF), Sandra Leon Alfonso (University Carlos III), Vincent Pons (Harvard Business School), Hanspeter Kriesi (EUI), Dominique Reynié (Sciences Po – CEVIPOF), Richard Nadeau (Université de Montréal), Éric Bélanger (McGill), Jean-François Daoust (University of Edinburgh), Ruth Dassonneville (Université de Montréal) and Erick Lachapelle (Université de Montréal).

M. Foucault and S. Brouard acknowledge the financial support from ANR – REPEAT grant (Special COVID-19), CNRS, Fondation de l'innovation politique, as well as regions Nouvelle-Aquitaine and Occitanie. Richard Nadeau and Éric Bélanger acknowledge the financial support from the SSHRC/CRSH.

The authors declare no conflict of interest.

A Guilt-Free Strategy to Increase Self-Reported Non-Compliance with COVID-19 Preventive Measures: Experimental Evidence from 12 Countries

Abstract

Studies of citizens' compliance with COVID-19 preventive measures routinely rely on survey data. While essential, public health restrictions provide clear signals of what is socially desirable in this context, creating a potential source of response bias in self-reported measures of compliance. In this research, we examine whether the results of a guilt-free strategy that was used by Daoust et al. (2020) to loosen this constraint are generalizable across twelve countries, and whether the treatment effect varies across subgroups. Our findings show that the guilt-free strategy is a very useful tool in every country included, increasing respondents' proclivity to report non-compliance by 9 to 16 percentage points. This effect holds for different subgroups based on gender, age and education. We conclude that the inclusion of this strategy should be the new standard for survey research that aims to provide crucial data on the current pandemic.

Keywords: COVID-19; Public health; Pandemic; Compliance; Measurement; Social desirability bias; Experimental method

In the fight against the spread of the COVID-19 virus, research that aims to explain compliance with public health preventive measures is of utmost importance. The severity of virus activity is, in no small part, a function of citizens' behaviours (Chan et al. 2020; Haushofer & Metcalf 2020). Therefore, much research has focused on understanding who complies, and what are the socio-demographic and attitudinal correlates of (non)compliance. Answering these questions is critical, pushing governments and public health agencies to gather reliable data on compliance with preventive measures (e.g. social distancing, the use of face masks, etc.). A great deal of this data comes from survey research.

The need for high quality data on compliance with public health measures has led some researchers to investigate the reliability of survey data more closely. In particular, some studies have examined the possibility that public health restrictions in the pandemic context have created common social norms for behaviours that are valued (e.g. social distancing). In turn, these norms create an incentive for respondents to under-report behaviours that are proscribed (e.g. social gatherings). The resulting social desirability bias can considerably affect the quality of data used by policy-makers and public health officials in their decision-making, which is problematic. Both Larsen et al. (2020) and Munzert and Selb (2020) considered the possibility that citizens' reported compliance with public health measures was affected by social desirability bias. These two single-country studies relied on a list experiment approach (also called unmatched item count technique). Reassuringly, they failed to detect social desirability in the self-reported behaviour of Danish (Larsen et al. 2020) and German (Munzert and Selb 2020) citizens. Findings from these list experiments, however, stand in contrast to the results of a recent study by Daoust et al. (2020), which tested different "face-saving" (or guilt-free) strategies designed to loosen the social norm

of compliance with public health measures in the context of three surveys conducted in Canada. The goal was to reduce social desirability in respondents' answers.

When exposed to a short preamble (referring to the fact that some people have altered their behaviours since the beginning of the pandemic, while others have continued to pursue various activities) combined with guilt-free answer choices (including “only when necessary”), respondents in the Daoust et al. (2020) study were substantially more likely than respondents from the control group to report non-compliant behaviour in the context of the COVID-19 pandemic. This suggests that there is a social desirability bias in citizens' self-reported behaviour when no guilt-free option is provided. While promising, these results are based on a single country, i.e. Canada. We do not know how effective the strategy is beyond the Canadian case. The question of generalizability is particularly important given that the list experimental approaches used in Denmark and Germany showed no signs of social desirability (Larsen et al. 2020; Munzert and Selb 2020).

In this research, we extend the most effective face-saving strategy identified by Daoust et al. (2020) to twelve countries. Doing so, we examine whether results are generalizable beyond Canada and across different contexts. In addition, we test whether the impact of the face-saving (guilt-free) answer option is homogenous across different subsets of the population to assess potential differentiated effects conditional on individual characteristics (Barari et al. 2020; Brouard et al. 2020). To preview our results, we show that the guilt-free strategy is a very useful tool in every country examined, increasing respondents' proclivity to report non-compliance by 9 to 16 percentage points. This effect holds for different subgroups based on gender, age and education.

We conclude that this method should become the new standard for survey research of citizens' compliance with COVID-19 preventive measures, ultimately providing higher-quality data to governments and public health agencies.

Measuring citizens' compliance with COVID-19 preventive measures

Social desirability bias has been a problem in survey research well before the COVID-19 pandemic and researchers have addressed the issue in several ways. Some work suggests that survey mode is an important factor, as the presence of a live interviewer can create greater incentives for respondents to provide more socially desirable answers. A meta-analysis on the topic suggests that an online mode is the best way to allow people to report undesirable behaviour (Gnambs and Kaspar 2014). But even with online surveys (e.g. without a live interviewer), there can still be social desirability bias. Survey researchers are therefore interested in developing additional ways and methodological tools to reduce social desirability and obtain more accurate estimates of undesirable behaviour (see for example Becher et al. 2020).

Focusing on the current pandemic context, some of the first experimental studies of measurement strategies to reduce social desirability bias found limited evidence of social desirability bias when employing the list experimental technique (Larsen et al. 2020; Munzert and Selb 2020). Munzert and Selb (2020) report a difference in the prevalence of not following social distancing between the treated and untreated group of 6 percentage points that is "barely significant at conventional levels due to the vast measurement error of the list estimate." (p. 206) They also find that the prevalence of the undesirable behavior is higher when using a multivalued response option compared to either a simple yes-no question or the list experimental estimate. This shows that

dichotomous direct questions should not be taken at face value. Discussing these findings, Munzert and Selb (2020) suggest that face-saving strategies might be a "valuable alternative" (p. 207). In this research, we test this face-saving strategy using a comparative experimental design and our results strengthen this conclusion.

The objective of the face-saving strategy is to reduce social desirability in respondents' answers by adding a (short) preamble and one or more guilt-free answering options. These steps should loosen the norm around a desirable response and make it more acceptable for respondents to admit non-compliance with the desirable outcome. Such an approach has been applied to a wide range of topics. For example, political scientists used it to study voter turnout (Belli et al. 2006; Morin-Chassé et al. 2017) where there is a clear norm that voting is the right thing to do. It has also proven to be useful to study illicit behavior like shoplifting (Holtgraves et al. 1997), sexual/health behaviour such as using a condom during sexual intercourse (Catania et al. 1996; Raghurir and Menon 1996), or consumer choice in market research (Nancarrow et al. 2001).

In the current context of the COVID-19 pandemic where it is crucial to understand citizens' level of compliance with preventive measures, Daoust et al. (2020) applied the face-saving strategy to analyze a potential social desirability bias. This latter work showed that face-saving strategies can increase the proportion of citizens who self-report non-compliance with a range of public health preventive measures in Canada. They argue that this increase in self-reported non-compliance is a consequence of reduced social desirability. While there is no objective benchmark to compare these estimates, they substantiate this claim by showing that a similar increase in self-reported

non-compliance is not observed when the same face-saving strategy is applied to a series of placebo behaviours that are not prohibited (e.g., grocery shopping).

While promising, the results from Daoust et al. (2020) suffer from a key limitation: Their focus on a single context, that is, the Canadian case in April 2020. In this paper, we address this issue by implementing the authors' approach in twelve different countries. By doing so, we can ascertain whether results are specific to a single context and time period, or whether the effectiveness of a face-saving answer option to reduce social desirability in self-reported compliance with public health measures applies more generally. In the next section, we detail our data and how we implemented the face-saving strategy.

Data and Indicators

We ran a face-saving (or guilt-free) experiment in twelve countries: Australia, Austria, Brazil, France, Germany, Italy, New Zealand, Poland, Spain, Sweden, the United Kingdom and the United States. We have obtained the written consent from participants who answered the survey. The data were analyzed anonymously. The project "Citizens' Attitudes Under COVID-19 Pandemic" has received the ethical certificate from Science Po Paris (France). The online surveys were conducted by three different data collection firms: IPSOS (for all countries except Australia, United States and Spain), CSA (for Australia and the United States) and Netquest (Spain). While it does not entail major differences, having different firms involved in data collection reduces the risk of bias due to potential "house effects." Data collection mainly occurred in mid- June 2020 within a period of a few days (maximum five), producing a nationally representative sample of about 1,000

respondents in most of the countries.¹ Countries included in the study experienced different levels of infection and death rates (Bosancianu et al. 2020) ranging from a low in Australia and New Zealand (with less than .5 deaths per 100 000 inhabitants) to a high experienced in the United Kingdom (with more than 59 deaths per 100 000 inhabitants). Although the countries in our sample were similarly influenced by public health guidelines established by global health authorities, the timing, stringency and components of public health measures adopted to combat the pandemic also strongly varied from Sweden (the least stringent) to New Zealand (the most stringent). Moreover, the countries in our sample also reflect different levels of politicization, with relatively more politicization of the pandemic response in Brazil and the USA than in other countries. Table A.1 of the Supplementary Material lists the exact dates of data collection in each country, as well as the number of observations. Moreover, Table A.2 shows the population per country and the death rates (per 100 000 inhabitants) as of June 15th and Table A.3. provides an overview of the preventive measures that were recommended and compulsory in each country.

Here, we make use of the most effective face-saving strategy identified by Daoust et al. (2020), which is also the strategy they recommend for future work (their Study 3). Extending this work to other countries, half of each national sample was randomly allocated a direct question while the other half received the face-saving treatment. The direct question was: “Have you done any of the following activities in the last week?” followed by a set of four items and yes/no answer choices.² The face-saving question preamble was:

“Some people have altered their behaviour since the beginning of the pandemic, while others have continued to pursue various activities. Some may also want to change their

¹ In France, Germany and the USA, the sample size is about 2000 respondents. In these cases, half of the sample answered the questions used in this research.

² Respondents could skip the question. Less than 0.5% did so in every country.

behaviour but cannot do so for different reasons. Have you done any of the following activities in the last week?”

Respondents in the treatment group received the answer options yes/occasionally/only when necessary/no. The first three answer choices indicated (and were coded as) non-compliance with the items. Of these three options, ‘occasionally’ and ‘only when necessary’ were the guilt-free answer choices. The four items, displayed in a random order, were:

- Go shopping or take public transportation without a face mask or taking it off during it
- Meet friends, family or colleagues greeting them by shaking hands, hugging or kissing
- Have a group of friends or family over at your place
- Participate in social activities (work, sport, religious ceremony...) without respecting physical distancing

These items refer to behaviours that are known to be crucial to minimize the spread of the disease among the population, that is, wearing a face mask and different forms of social distancing (Chu et al. 2020; Courtemanche et al. 2020). Moreover, greeting people by shaking hands, hugging or kissing was clearly not recommended, while hosting a gathering at one’s place was allowed though not without some level of risk (e.g., Center for Disease Control and Prevention 2020, see the “Hosting gatherings or cook-outs” section).

In a second step, we explore whether the effects of the treatments are heterogeneous. We consider respondents' gender (female or male), age (treated as linear, from 18 to 91), and their level of education. Education was measured using different categories in each country given their different educational systems. We use a 0 to 1 “university graduate” scale in order to consistently model the effect of having obtained a university education in each country. We also provide individual analyses for each country. For descriptive statistics on how much these preventive measures were

respected in the twelve countries, see Figure B.1 of the Supplementary Material.³ Descriptive statistics for gender, age and education, are reported in Appendix B of the Supplementary Material.

Results

Our main goal is to ascertain whether providing a short preamble and guilt-free answer options increases citizens' likelihood of reporting non-compliance with important public health measures like mask-wearing and social distancing. To shed light on this question, we analyze the data discretely across the four items as well as across the twelve countries. We thus provide a complete picture of the experimental effects and avoid pooling to ensure that the results are not driven by certain items or countries. Figure 1 displays the proportion of non-compliers with the preventive measures for the control (direct question) and the treatment (face-saving) groups. The treatment group is depicted in grey, while the black bars indicate levels of reported compliance in the control condition. With four items in twelve countries, Figure 1 plots a total of 48 effects of interest.

³ Figure B.2 in the appendix distinguishes between the two guilt-free answer choices (“Occasionally” and “Only when necessary”), showing that there are no substantial differences between them with the only two exceptions being Brazil and Spain, where non-compliers tend to prefer the “only when necessary” answer choice.

Figure 1. Non-compliance in the control and treatment (face-saving) groups

Note: Means of non-compliance are shown with 95% confidence intervals included. ATE=Average treatment effect.

As clear from eyeballing the graphs, the means of non-compliance are systematically higher in the face-saving group (the grey bars). More precisely, the impact of the treatment is positive in 47 out of 48 cases (the single exception being Austria for the face mask item). Substantially speaking, the average treatment effect (averaged across all countries) of receiving the treatment ranges from 9 (the face mask) to 16 percentage points (hosting at home). Greeting people with non-recommended behaviours and not respecting physical distancing show an ATE of 12 and 14 percentage points, respectively. While we prefer to focus on the substantive effect, we note that in most cases (45 out of 47 positive effects), the differences are significant at $p < .05$ (based on two-sided t -test). Moreover, we implemented the Romano-Wolf correction given the number of tests that we presented (Clarke et al. 2020; Romano & Wolf 2007). Tables D.2-5 show that the differences

between the treatment and the control group are systematically different (that is, in 45 cases) even using Romano-Wolf p -values.

Does the impact of the face-saving treatment differ across various subgroups of the population? In an exploratory fashion, we look at the potential moderation effect from gender, age and education, which are known to be linked to compliance with preventive measures (Barari et al. 2020; Brouard et al. 2020; Galasso et al. 2020). For this analysis, we rely on a pooled dataset that includes the information from all countries. Figure 2 shows the coefficients from logistic regressions for different groups. The full regression outputs can be found in Table C.1-3 of Appendix C. The results for potential heterogeneous effects across subgroups based on gender, age and education are quite clear: Based on the evidence, we cannot reject the null hypothesis of no moderation effect. The interaction coefficients never reach statistical significance at $p < .05$. The direction of the effects is perfectly split for gender, three out of four coefficients are negative for age and three are positive for education.

Figure 2. Interaction coefficients for the guilt-free treatment, by gender, age and education

Note: The values of the interaction coefficients estimated in Tables C.1-3 of the Supplementary Material are shown, with 95% confidence intervals included.

All in all, we find that the face-saving strategy is effective. In total, 47 out of 48 effects are positive, and, their effects are substantive, ranging from 9 to 16 percentage points. Moreover, the impact of the treatment effect is fairly homogenous. As shown in Figure 2, it does not seem to be conditioned by peoples' gender, age or level of education.

We made sure that our conclusions were robust in several ways. First, although we should be cautious about randomization checks in an experimental context (Mutz et al. 2019), we verified that both control and treatment groups were similar in terms of age, gender and education. The average age was identical at 47 (control v/s treatment); 52% of people in the control group were women compared to 51% in the treatment group, and means for education were the same at .60.

Second, using weights for age, gender, education and region does not alter our conclusions. More specifically, we replicated Figure 1 and 2 for the weighted dataset (see Figure D.1 and D.2 in the Supplementary Material). One interaction out of twelve reaches $p < .05$, that is, the interaction between age and treatment for the ‘no mask’ behaviour. Focusing on the coefficients rather than p -values, our findings are very similar.

Third, we made sure that our results of the tests for heterogeneous treatment effects were not driven by particular countries by estimating the effects with a model that includes country fixed effects. Figure D.3 and D.4 replicate and Table D.1-4. shows the results of this test and leads to essentially the same conclusion.

Fourth, we tackle the possibility that our results in fact are a “false positive” (Hoglinger and Jann 2018). In a nutshell, our experimental design is based on the assumption that the differences in the proportion of self-reported non-compliance between the direct and the face-saving questions are related to a reduction of incentives to report socially desirable behaviours. While we think this is a very plausible assumption, we also examine the possibility that another mechanism might explain our results (for example the mere change in the number of response options) using a Canadian survey that was in field during the same period as the twelve surveys examined here. In this Canadian survey, we used the same face-saving strategy but with a broader battery of items, 8 instead of 4. This larger battery included behaviours that were not officially prohibited, i.e. where there should be much less social desirability. In panel A of Figure D.4, we show that there is a strong effect of the treatment (about 10 percentage points) for behaviours that the government prescribed, such as wearing a face-mask in public, and that this effect is much less important in

panel B (average of about 3 percentage points) for behaviours that perhaps entail a risk, but that the government did not proscribe – such as taking public transportation or shopping for non-essential products. While we do not have these ‘placebo’ items for twelve countries included in the analyses, this result for Canada is reassuring and increases our confidence that the greater proportion of self-reported non-compliance in the face-saving group is not a ‘false positive.’

Fifth, we also check whether or not a potential experimenter demand effect (de Quidt et al. 2018) might drive the results using data from a survey experiment in France. The design of the experiment is presented in the Supplementary Material. Beyond both question formats analyzed earlier, we rely on a list experiment (and a question used by Munzert & Selb (2020)) to estimate the levels of compliance with preventive behaviors from alternative measurement strategies. Results are shown in Figure D.5 of the Supplementary Material. The list experiment approach was designed to reduce social desirability bias and is usually employed for this purpose (Blair and Imai 2012). Most importantly, the risk that there would be an experimenter demand effect is very limited. Overall, even if this experiment only includes one country, using estimates from the list experiment as a baseline brings no evidence of an experimenter demand effect associated to the guilt-free question format that would systematically bias our estimates.

Discussion and implications

West et al. (2020: 451) argued that “there is an urgent need to develop and evaluate interventions to promote effective enactment of these behaviours and provide a preliminary analysis to help guide this.” We agree with the authors and address the issue of social desirability and its impact on reported compliance with public health measures. While the work of Daoust et al. (2020) on

this topic showed a face-saving strategy is a promising approach to attenuate social desirability, evidence was lacking on whether this face-saving strategy was effective beyond Canada.

In this research, we tested the face-saving strategy using a survey experiment in twelve countries to examine the benefits of this approach. We replicated the findings of Daoust et al. (2020) and most importantly, did so in a diverse set of contexts, with different countries and political systems, in different stages of the pandemic (deconfinement in most countries), with different levels of infections, etc. Based on four public health preventive measures related to the wearing of face masks, greetings, hosting gatherings at one's home, and social distancing, we found that the face-saving strategy increased the proportion of citizens who readily answer that they did not comply for 47 out of 48 cases. Most importantly, the effects were substantial. They ranged from 9 percentage points (the mask item) to 16 percentage points (hosting at home) and are robust to several additional tests. Given that there is no objective benchmark for the four preventive measures analyzed in our study that are readily accessible for researchers, we must acknowledge that there is no way to compare our estimates with external measures. Altogether, our experimental design and extensive robustness checks bolster the conclusion that the differences we find can be at least partly attributed to a social desirability bias.

There have already been major advances in the development of observational measures of citizens' behaviour (i.e. not from survey data). Among others, France has recently used cameras in its subway stations to quantify the proportion of people who wear a mask when in public transport, and several countries are developing applications to track the inter-regional movement of their residents (Olivier et al. 2020). While very useful, we cannot rely solely on behavioral data in the fight against the pandemic. First, observational measures are not available for several important

preventive public health measures, as many measures cannot be examined in public, such as the respect of social distancing if one hosts a gathering at their private home. Second, even behavioral data like that obtained from cameras or tracing applications have some major drawbacks. Most importantly, due to technical and privacy constraints this approach does not easily provide researchers important auxiliary information about who complies and what makes people comply or not. For these reasons, we believe that survey research is a crucial complement to other data sources.

In summary, policy-makers and public health experts require survey data, and we should aim for data of the best possible quality. Our research confirms that using a guilt-free strategy is an effective approach and is relevant to anyone who aims to provide data on citizens' compliance with COVID-19 preventive measures. This type of data is crucial for governments and public health agencies to make enlightened decisions. Moreover, as the strategy simply implies the addition of a very short preamble and guilt-free answer choices, there are very limited additional costs involved to implement this method compared to a direct question. While replications would be welcome to strengthen the validity of the approach, we believe that our comparative research provides a firm ground for what should become the standard when measuring citizens' compliance with public health preventive measures.

References

- Barari, S., Caria, S., Davola, A., Falco, P., Fetzer, T., Fiorin, S., ... and Kraft-Todd, G. (2020). Evaluating COVID-19 Public Health Messaging in Italy: Self-Reported Compliance and Growing Mental Health Concerns. *medRxiv*. doi: <https://doi.org/10.1101/2020.03.27.20042820>
- Belli, R.F., Moore, S. E., VanHoewyk, J. (2006). An experimental comparison of question forms used to reduce vote overreporting. *Electoral Studies*, 25(4): 751-759.

Bosancianu, C. M., Dionne, K. Y., Hilbig, H., Humphreys, M., KC, S., Lieber, N., & Scacco, A. (2020, Version 1). Political and Social Correlates of Covid-19 Mortality. SocArxiv. <https://doi.org/10.31235/osf.io/ub3zd>

Becher, M., Stegmüller, D., Brouard, S. and Kerrouche, E. (2020). Comparative Experimental Evidence on Compliance with Social Distancing During the Covid-19 Pandemic. SSRN: <http://dx.doi.org/10.2139/ssrn.3652543>

Blair, G., & Imai, K. (2012). Statistical analysis of list experiments. *Political Analysis*, 20(1), 47-77.

Brouard, S., Vasilopoulos, P. and Becher, M. (2020). Sociodemographic and psychological correlates of compliance with the Covid-19 public health measures in France. *Canadian Journal of Political Science*. DOI: <https://doi.org/10.1017/S0008423920000335>

Catania, J. A., Binson, D., Canchola, J., Pollack, L. M., Hauck, W., & Coates, T. J. (1996). Effects of interviewer gender, interviewer choice, and item wording on responses to questions concerning sexual behavior. *Public Opinion Quarterly*, 60(3), 345-375.

Center for Disease Control and Prevention. (2020). Personal and Social Activities. Link: <https://www.cdc.gov/coronavirus/2019-ncov/daily-life-coping/personal-social-activities.html> (consulted on August 2nd)

Chan, D. K., Zhang, C. Q., & Josefsson, K. W. (2020). Why people failed to adhere to COVID-19 preventive behaviors? Perspectives from an integrated behavior change model. *Infection Control & Hospital Epidemiology*, 1-6.

Chu, D. K., Akl, E. A., Duda, S., Solo, K., Yaacoub, S., Schünemann, H. J., ... & Hajizadeh, A. (2020). Physical distancing, face masks, and eye protection to prevent person-to-person transmission of SARS-CoV-2 and COVID-19: a systematic review and meta-analysis. *The Lancet*.

Clarke, D., Romano, J. P., & Wolf, M. (2020). The Romano-Wolf Multiple Hypothesis Correction in Stata. IZA Working Paper. <https://www.iza.org/publications/dp/12845/the-romano-wolf-multiple-hypothesis-correction-in-stata>

Courtemanche, C., Garuccio, J., Le, A., Pinkston, J., & Yelowitz, A. (2020). Strong Social Distancing Measures In The United States Reduced The COVID-19 Growth Rate: Study evaluates the impact of social distancing measures on the growth rate of confirmed COVID-19 cases across the United States. *Health Affairs*, 10-1377.

Daoust, J. F., Nadeau, R., Dassonneville, R., Lachapelle, E., Bélanger, É., Savoie, J., & van der Linden, C. (2020). How to survey citizens' compliance with COVID-19 public health measures. Evidence from three survey experiments. *Journal of Experimental Political Science*.

- Gnambs, T., & Kaspar, K. (2015). Disclosure of sensitive behaviors across self-administered survey modes: a meta-analysis. *Behavior research methods* 47(4): 1237-1259.
- Galasso, V., Pons, V., Profeta, P., Becher, M., Brouard, S., and Foucault, M. (2020). Gender differences in COVID-19 attitudes and behavior: Panel evidence from eight countries. *PNAS*, 117 (44): 27285-27291.
- Haushofer, J., & Metcalf, C. J. E. (2020). Which interventions work best in a pandemic? *Science* 368 (6495): 1063-1065.
- Höglinger, M., & Jann, B. (2018). More is not always better: An experimental individual-level validation of the randomized response technique and the crosswise model. *PLoS one*, 13(8), e0201770.
- Holtgraves, T., Eck, J. and Lasky, B. (1997). Face Management, Question Wording, and Social Desirability. *Journal of Applied Social Psychology*, 27: 1650-1671
- Larsen, M. V., Petersen, M. B., & Nyrup, J. (2020). Do Survey Estimates of the Public's Compliance with COVID-19 Regulations Suffer from Social Desirability Bias? *Journal of Behavioral Public Administration*, 3(2).
- Morin-Chassé, A., D. Bol, L. Stephenson, and S. Labbé St-Vincent (2017). How to Survey About Electoral Turnout? The Efficacy of the Face-Saving Response Items in 19 Different Contexts. *Political Science Research and Methods* 5(3): 575–584.
- Munzert, S., & Selb, P. (2020). Can we directly survey adherence to non-pharmaceutical interventions? Evidence from a list experiment conducted in Germany during the early Corona pandemic. *Survey Research* 14(2): 205-209.
- Mutz D., R Pemantle & Philip Pham. (2019). “The Perils of Balance Testing in Experimental Design: Messy Analyses of Clean Data.” *The American Statistician*, 73:1, 32-42
- Nancarrow, C., Brace, I. and Wright, L. T. (2001). “Tell me Lies, Tell me Sweet Little Lies”: Dealing with Socially Desirable Responses in Market Research. *The Marketing Review*, 2(1): 55-69.
- Oliver, N., Lepri, B., Sterly, H., Lambiotte, R., Deletaille, S., De Nadai, M., ... & Colizza, V. (2020). Mobile phone data for informing public health actions across the COVID-19 pandemic life cycle. *Science Advances*.
- de Quidt, J., Haushofer, J., & Roth, C. P. (2018). Measuring and Bounding Experimenters Demand. *American Economic Review*, 108(11), 3266–3302. <https://doi.org/10.1257/aer.20171330>
- Raghubir, P., & Menon, G. (1996). Asking sensitive questions: The effects of type of referent and frequency wording in counterbiasing methods. *Psychology & Marketing*, 13(7), 633-652.

Romano, J. P., & Wolf, M. (2007). Control of generalized error rates in multiple testing. *The Annals of Statistics*, 35(4), 1378-1408. <http://dx.doi.org/10.1214/009053606000001622>

West, R., Michie, S., Rubin, G. J., & Amlôt, R. (2020). Applying principles of behaviour change to reduce SARS-CoV-2 transmission. *Nature Human Behaviour* 4 (May): 451-459.

Supplementary Material

A Guilt-Free Strategy to Increase Self-Reported Non-Compliance with COVID-19 Preventive Measures: Experimental Evidence from 12 Countries

PLOS One

Appendix A: Description of the samples and national contexts

Table A.1. Description of the samples

Country	Dates on the field	Number of observations
Australia	June, 15 th – 19 th	1,003
Austria	June, 23 rd – 27 th	1,011
Brazil	June, 23 rd – 28 th	1,000
France	June, 22 nd – 25 th	1,006
Germany	June, 23 rd – 27 th	1,004
Italy	June, 24 th – 27 th	1,003
New Zealand	June, 23 rd – 28 th	1,000
Poland	June, 24 th – 27 th	1,014
Spain	July, 6 th – 10 th	961
Sweden	June, 23 rd – 27 th	1,017
United Kingdom	June, 21 st – 24 th	1,014
United States	June, 16 th – 22 nd	1,001
Canada*	June, 16 th – 24 th	1,002

Note. *: used for robustness checks only.

Table A.2. Population and mortality rate across countries

Country	Population (million)	COVID-19 Mortality rate per 100 000 inhabitants (June 15)
Australia	25.36	0.40
Austria	8.87	7.63
Brazil	211.05	20.82
France	67.06	43.80
Germany	83.13	10.59
Italy	60.29	57.00
New Zealand	4.91	0.44
Poland	37.97	3.30
Spain	47.07	57.64
Sweden	10.28	47.55
US	328.24	36.19
United Kingdom	66.83	58.94
Canada*	37.58	21.88

*Note: Source for the population is the World Bank and the Mortality rates and computed using John Hopkins Coronavirus Resource Center data (accessible on github). *: used for robustness checks only.*

Table A.3. COVID-19 Preventive Measures across countries

	Mandatory use of masks		Mandatory physical distancing		Stay at home requirement	
	in shops and public transportation		Policy	Scope	Policy	Scope
Australia	No	National	Yes	National	No	National
Austria	In public transportation	National	Yes	National	No	National
Brazil	In shops and/or public transportation	In specific areas in state or in whole state	Yes	In specific areas in state or in whole state	Require not leaving house with exceptions for daily exercise, grocery shopping, and 'essential' trips	In specific areas
France	In public transportation	National	Yes	National	No	National
Germany	In public transportation and shops	National (agreement of federal and state governments) / all states except Berlin	Yes	National	No	National
Italy	In shops and public transportation	National	Yes	National	Recommend not leaving house	National
New Zealand	No	National	No	National	No	National
Poland	In shops and public transportation	National	yes	National	No	National
Spain	In shops and restaurants; also in all public spaces when social distancing not possible	Specific regions / national	Yes	National	Recommend not leaving house	In specific regions
Sweden	No	National	No	National	Recommend not leaving house	National
United Kingdom	In public transportation	National	Yes	National	Recommend not leaving house	In specific areas
United States	In shops and/or public transportation	In specific areas in state or in whole state	Yes	In specific areas in state or in whole state	Require not leaving house with exceptions for daily exercise, grocery shopping, and 'essential' trips	In specific areas in state or in whole state
Canada*	For air passengers / in shops and public transportation	National / in some provinces or in specific areas of some provinces	Yes	National	Recommend not leaving house	National

Source: <https://github.com/amel-github/covid19-interventionmeasures>; <https://masks4all.co/what-countries-require-masks-in-public>; <https://airtable.com/shrxEdbaBJIQwMUgA/tbl5o6qUd54BL9wkw>; <https://www.bsg.ox.ac.uk/research/research-projects/coronavirus-government-response-tracker>; <https://kubinec.shinyapps.io/coronanet/>; and further authors' corroborations with official and media sources.

Appendix B: Descriptive statistics, by country

Figure B.1. Non-compliance with preventive measures

Note: Means are shown for each item in each country, with 95% confidence included.

Figure B.2. Non-compliance, by guilt-free answer choices

Note: Means are shown for each item and guilt-free answer choices, with 95% confidence included.

Table B.1. Descriptive statistics for age, gender and education.

	Australia		Austria		Brazil		France		Germany		Italy	
	Mean	Std. Dev.	Mean	Std. Dev.	Mean	Std. Dev.	Mean	Std. Dev.	Mean	Std. Dev.	Mean	Std. Dev.
Age	46.64	14.43	47.30	16.47	41.33	15.01	3.97	1.70	51.02	15.91	47.68	16.48
Gender	0.52	0.50	0.47	0.50	0.52	0.50	0.49	0.50	0.51	0.50	0.53	0.50
Education	0.46	0.27	0.49	0.22	0.63	0.15	0.72	0.31	0.62	0.28	0.57	0.19
	NZ		Poland		Spain		Sweden		UK		USA	
Age	48.37	15.58	43.96	15.37	48.27	17.54	51.45	16.14	46.65	16.03	46.41	17.90
Gender	0.53	0.50	0.53	0.50	0.52	0.50	0.49	0.50	0.51	0.50	0.53	0.50
Education	0.55	0.26	0.71	0.26	0.54	0.19	0.65	0.34	0.61	0.29	0.58	0.21

Note: For gender, female=1

Appendix C. Regression tables

Table C.1. Regressions outputs for panel a of Figure 2

	Face mask	Greeting	Host	Distance
Gender (1=Female)	-0.14 (0.05)	-0.08 (0.06)	-0.03 (0.06)	-0.22 (0.07)
Treatment	0.40 (0.05)	0.48 (0.05)	0.76 (0.06)	0.78 (0.06)
Gender $\times$ Treatment	-0.05 (0.07)	0.09 (0.08)	-0.07 (0.08)	0.02 (0.09)
Constant	-0.50 (0.04)	-0.77 (0.04)	-0.99 (0.04)	-1.44 (0.05)
Observations	12026	12026	12023	12028

Note: Logistic regression coefficients with standard errors in parentheses.

Table C.2. Regressions outputs for panel b of Figure 2

	Face mask	Greeting	Host	Distance
Age	-0.0089 (0.0017)	-0.0284 (0.0018)	-0.0168 (0.0018)	-0.0316 (0.0022)
Treatment	0.5295 (0.1151)	0.6885 (0.1200)	0.9339 (0.1202)	0.8246 (0.1327)
Age $\times$ Treatment	-0.0032 (0.0023)	-0.0028 (0.0025)	-0.0042 (0.0025)	0.0002 (0.0029)
Constant	-0.1536 (0.0822)	0.5044 (0.0859)	-0.2233 (0.0881)	-0.1353 (0.1009)
Observations	12026	12026	12023	12028

Note: Logistic regression coefficients with standard errors in parentheses.

Table C.3. Regressions outputs for panel c of Figure 2

	Face mask	Greeting	Host	Distance
Education	-0.24 (0.10)	0.22 (0.11)	0.24 (0.11)	0.21 (0.13)
Treatment	0.23 (0.09)	0.63 (0.09)	0.69 (0.10)	0.71 (0.11)
Education $\times$ Treatment	0.24 (0.14)	-0.17 (0.14)	0.05 (0.15)	0.12 (0.17)
Constant	-0.44 (0.07)	-0.94 (0.07)	-1.15 (0.07)	-1.68 (0.09)
Observations	11946	11946	11943	11948

Note: Logistic regression coefficients with standard errors in parentheses.

Appendix D. Robustness checks

Figure D.1. Replication of Figure 1 with weights

Note: Means of non-compliance are shown with 95% confidence intervals included. ATE=Average treatment effect.

Figure D.2. Replication of Figure 2 with weights

Note: The values of the interaction coefficients with 95% confidence intervals included.

Figure D.3. Average treatment effects controlling for country fixed effects

Note: Average treatment effects are estimated from Table D.1. and are shown with 95% confidence intervals.

Table D.1. Regressions models with country fixed effects

<i>Dependent variable =</i>	No mask	Greeting	Host	No distance
Guilt-free treatment	0.49 (0.04)	0.58 (0.04)	0.76 (0.04)	0.86 (0.05)
Austria	-1.28 (0.10)	0.07 (0.09)	-0.09 (0.09)	-0.12 (0.10)
Brazil	-2.81 (0.11)	-0.72 (0.10)	-0.86 (0.10)	-0.74 (0.11)
France	-1.88 (0.10)	-0.29 (0.09)	0.49 (0.09)	-0.67 (0.11)
Germany	-2.58 (0.11)	-0.44 (0.10)	-0.37 (0.10)	-0.58 (0.11)
Italy	-2.38 (0.11)	0.12 (0.09)	0.41 (0.09)	-0.26 (0.10)
New Zealand	0.52 (0.11)	1.18 (0.09)	0.35 (0.09)	1.15 (0.10)
Poland	-1.45 (0.10)	0.78 (0.09)	0.46 (0.09)	0.50 (0.10)
Spain	-3.60 (0.14)	0.11 (0.09)	-0.13 (0.10)	-0.36 (0.11)
Sweden	-0.40 (0.10)	-0.51 (0.10)	-0.48 (0.10)	-0.64 (0.11)
United Kingdom	-1.43 (0.10)	-1.59 (0.12)	-1.09 (0.11)	-1.38 (0.13)
United States	-1.69 (0.10)	-0.19 (0.09)	-0.19 (0.10)	0.09 (0.10)
Constant	0.85 (0.08)	-0.76 (0.07)	-0.94 (0.07)	-1.43 (0.08)
Observations	12026	12026	12023	12028

Note. Logistic regression predicting non-compliance. Standard errors in parentheses. The reference category for the countries is Australia.

Figure D.4. Prohibited and non-prohibited (placebo) items

Note: Means of non-compliance are shown with 95% confidence intervals included

Figure D.5. Comparison with other estimation techniques

Note: Means with 95% confidence intervals are shown. D.Q. indicates estimation from the “dichotomous” question format; P.Q. indicates estimation from the “polytomous” question format; F.S. indicates estimation from the “face saving” question format; L.E. indicates estimation from the list-experiment estimation strategy. Masks refers to the following items: Go shopping or take public transportation without a face mask or taking it off during it; Greetings refers to the following items: Meet friends, family or colleagues greeting them by shaking hands, hugging or kissing; Activities refers to the following items: Participate in social activities (work, sport, religious ceremony...) without respecting physical distancing. Friends refers to the following items: Have a group of friends or family over at your place without respecting physical distancing.

Data: We ran a quota-based online survey in France that was in field from July 9-19th. As nearly all surveys in France, sampling was done using a quota method based on age, gender, occupation, regions and type of residential area. The sample is designed to be representative of the electorate’s features in each of the 12 metropolitan French regions. 10,686 respondents answered our questionnaire.

The same four items included in the survey were:

- Go shopping or take public transportation without a face mask or taking it off during it;
- Meet friends, family or colleagues greeting them by shaking hands, hugging or kissing;
- Have a group of friends or family over at your place without respecting physical distancing;
- Participate in social activities (work, sport, religious ceremony...) without respecting physical distancing.

We used four different strategies to address the issue of social desirability. Three approaches focusing on the question wording and answer options, and a list-experimental approach:

- First, a set of 1343 respondents were exposed to the following “dichotomous” question wording discussed by Daoust et al. (2020): “Have you done any of the following activities in the last week?”. The response categories were (1) yes, (2) no.
- Second we administered to another group of 1337 respondents the “polytomous” question used by Munzert & Selb (2020) : “How often have you done the following activities in the last week?”. The response categories were (1) daily, (2) several times, (3) once, (4) never.
- The third type of questions included is the “face-saving” question proposed by Daoust et al. (2020): “Some people have altered their behaviour since the beginning of the pandemic, while others have continued to pursue various activities. Some may also want to change their behaviour but cannot do so for different reasons. Have you done any of the following activities in the last week?”. 1345 respondents were to choose between the following categories: (1) Yes, (2) Occasionally, (3) Only when necessary, (4) No.
- Finally, we also embedded a list experiment (Becher 2020; Larsen 2020; Munzert & Selb 2020) in this survey. Faced with a list of items, a fourth group of respondents were asked how many things they had done last week, but not which specific ones. Respondents were randomly assigned to either the control group (N=1334) or one of the four treatment groups (N=1332; N=1331; N=1335; N=1329). The treated groups received an additional item capturing the violation of the social distancing norm. The control group received a list of four behaviours that are generally permissible under existing health recommendations. The question reads as follow: “How many of these things have you done last week? You do not need to tell me which ones you have done, just how many.” The following four activities were listed: I went to the doctor or to the hospital; I watched a movie or TV show online using a streaming service (e.g., Amazon, Netflix); I exercised outdoors; I ordered food using an online delivery service. In addition to these four items, the treated groups received one of the four more sensitive items we study. The estimated level of non-compliance was obtained from the simple differences-in-means.

Turning to the results from these different approaches, as shown in Figure D.5., for 3 out of 4 items the level of non-compliance estimated by the face-saving question (the estimates denoted by F.S.) is in the 95% confidence interval of the level of non-compliance estimated by the list experiment (the estimates denoted by L.E.). For one item, the face-saving estimate is 4.5 points below and for both other ones it is 2 points higher. For the items related to mask-wearing, greetings and social-distancing in activities, estimates from the dichotomous questions are systematically below the list-experiment estimates and beyond its 95% confidence intervals. The “face-saving” treatment does not seem to induce a systematic experimenter demand effect that would inflate non-compliance, at least compared with an alternative measurement strategy - list experiment – that is well known for decreasing desirability bias (Blair and Imai 2012) and does not appear prone to any experimenter demand effect due to the lack of any associated framing.

The pattern described above differs only for non-compliance in social distancing while hosting friends. In this instance, the list experiment estimate is the lowest. Only the estimate from the dichotomous question does not statistically differ from it. Both the estimates of the face-saving and polytomous questions are significantly higher. The polytomous question is devoid of any framing in the wording of the question. Therefore, even if we have no explanation for this

idiosyncratic pattern, it is unlikely that the higher estimate produced by the face-saving question relative to the list experiment is due to a potential experimenter demand effect induced by the face-saving treatment.

Overall, even if this experiment only includes France, it brings no evidence of an experimenter demand effect that would systematically bias our estimates.

Table D.2. Romano-Wolf test for the “No Mask” item

	Model p-value	Resample p-value	Romano-Wolf p-value
Australia	<i>0.0039</i>	<i>0.0050</i>	<i>0.0200</i>
Austria	<i>0.0156</i>	<i>0.0230</i>	<i>0.0509</i>
Brazil	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>
France	<i>0.0002</i>	<i>0.0020</i>	<i>0.0030</i>
Germany	<i>0.0074</i>	<i>0.0060</i>	<i>0.0320</i>
Italy	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>
New Zealand	<i>0.0600</i>	<i>0.0619</i>	<i>0.1249</i>
Poland	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>
Spain	<i>0.0001</i>	<i>0.0010</i>	<i>0.0030</i>
Sweden	<i>0.0000</i>	<i>0.0010</i>	<i>0.0020</i>
United Kingdom	<i>0.1295</i>	<i>0.1518</i>	<i>0.1518</i>
United States	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>

Note: Default options from the rwolf package was used, with the number of repetitions set at 1000.

Table D.3. Romano-Wolf test for the “Greeting” item

	Model p-value	Resample p-value	Romano-Wolf p-value
Australia	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>
Austria	<i>0.0071</i>	<i>0.0080</i>	<i>0.0210</i>
Brazil	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>
France	<i>0.0004</i>	<i>0.0010</i>	<i>0.0010</i>
Germany	<i>0.0001</i>	<i>0.0010</i>	<i>0.0010</i>
Italy	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>
New Zealand	<i>0.0002</i>	<i>0.0010</i>	<i>0.0010</i>
Poland	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>
Spain	<i>0.0132</i>	<i>0.0150</i>	<i>0.0260</i>
Sweden	<i>0.0004</i>	<i>0.0010</i>	<i>0.0010</i>
United Kingdom	<i>0.0380</i>	<i>0.0360</i>	<i>0.0360</i>
United States	<i>0.0001</i>	<i>0.0010</i>	<i>0.0010</i>

Note: Default options from the rwolf package was used, with the number of repetitions set at 1000.

Table D.4. Romano-Wolf test for the “Host” item

	Model p-value	Resample p-value	Romano-Wolf p-value
Australia	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>
Austria	<i>0.0035</i>	<i>0.0050</i>	<i>0.0050</i>
Brazil	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>
France	<i>0.0001</i>	<i>0.0010</i>	<i>0.0020</i>
Germany	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>
Italy	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>
New Zealand	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>
Poland	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>
Spain	<i>0.0002</i>	<i>0.0010</i>	<i>0.0030</i>
Sweden	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>
United Kingdom	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>
United States	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>

Note: Default options from the rwolf package was used, with the number of repetitions set at 1000.

Table D.5. Romano-Wolf test for the “Social Distance” item

	Model p-value	Resample p-value	Romano-Wolf p-value
Australia	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>
Austria	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>
Brazil	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>
France	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>
Germany	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>
Italy	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>
New Zealand	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>
Poland	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>
Spain	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>
Sweden	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>
United Kingdom	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>
United States	<i>0.0000</i>	<i>0.0010</i>	<i>0.0010</i>

Note: Default options from the rwolf package was used, with the number of repetitions set at 1000.