ACADL is differentially expressed in malignant mesothelioma.

Shahan Mamoor, MS¹

¹Thomas Jefferson School of Law

San Diego, CA 92101

mamoorsk@tjisl.edu

Mesothelioma is an aggressive cancer often caused by exposure to asbestos (1) with a median survival of 7 months upon treatment with first-line chemotherapy (2). There are zero targeted treatment options available for mesothelioma. We mined two public datasets (3, 4) to identify the genes most differentially expressed in mesotheliomas compared to the tissues and organ in which it emerges - the peritoneum and the lung parenchyma. We found that the expression of ACADL changed more than greater than 99% of the genome between the tumors of patients with mesothelioma and the normal lung and peritoneum. This was conserved across two types of mesotheliomas and across independent datasets. This is the first report of ACADL differential expression in mesothelioma. ACADL should be evaluated as a novel therapeutic target in mesothelioma.

Keywords: mesothelioma, thoracic cancers, ACADL, systems oncology, tumor transcriptome analysis. targeted therapeutics in mesothelioma.

Introduction

Mesothelioma is a cancer of the mesothelial cells that compose the pleural mesothelium, the cardiac mesothelium, and the serosal mesothelium, that line the lungs, the heart, and the gut (as peritoneum) (5). Exposure to the industrial substance asbestos is a common cause of mesothelioma but the SV40 virus has also been implicated (1). The most frequently prescribed first line chemotherapy consists of pemetrexed with carboplatin or cisplatin with gemcitabine as an option for addition upon retreatment (2, 6, 7). First-line treatment provides 7 month overall survival and retreatment can add 5 more months (2). A recent study demonstrated that addition of bevacizumab extended median overall survival (8). There are zero targeted treatments available for patients with mesothelioma.

In this study, we performed global differential gene expression profiling in mesothelioma by comparing the transcriptomes of mesothelioma cancer tissue to that of the normal lung or peritoneum using two separate datasets (3, 4). We found that the acyl Co-A dehydrogenase, long chain, or ACADL, was among the genes whose expression changed the most between mesothelioma tumor and normal tissues of the lung and peritoneum. This finding was consistent between both datasets. ACADL should be evaluated for its role to support tumor initiation, maintenance, or progression in xenograft mouse models of mesothelioma (9).

Methods

Datasets GSE51024 (3) and GSE112154 (4) were utilized here. Investigators that created GSE51024 utilized Affymetrix Human Genome U133 Plus 2.0 Array technology. Investigators that created GSE112154 utilized Illumina HumanHT-12 V4.0 expression beadchip. In GSE51024, n=41 normal lung parenchyma (paired) samples and n=55 mesothelioma (malignant pleural). In GSE112154, n=3 normal peritoneum samples and n=45 mesothelioma samples (diffuse malignant peritoneal).

The Benjamini and Hochberg method of p-value adjustment was utilized, and p-value adjustment was set to “Auto-detect”. The NCBI generated category of platform annotation was utilized. To evaluate the statistical significance of differences in RNA expression values in control versus tumor groups, a two-tailed, unpaired t test with Welch’s correction was performed to obtain a p-value. P-values less than 0.05 were considered statistically significant.

Results

We performed whole transcriptome analysis of mesothelioma using two separate datasets (3, 4) by comparing the transcriptomes of mesothelioma tissue to that of normal lung and normal peritoneum. We found that ACADL was among the most differentially expressed genes in mesothelioma tumor tissues.

ACADL is one of the genes whose expression changes most significantly between the transcriptomes of mesothelioma tumor and normal tissues of the lung and peritoneum.

We asked which genes were expressed the most significantly differently between the transcriptomes of 55 malignant pleural mesothelioma samples and 41 normal lung parenchyma samples (3). ACADL was identified as the most differentially expressed gene (rank 1) out of 54675 transcripts detected and measured in this dataset. This was statistically quite significant (Table 1; $p=1.27E-40$).

We probed a separate dataset to determine if this could be validated using independent data (4). We asked what genes were expressed most significantly differently between the transcriptomes of 3 normal peritoneum samples and 45 diffuse malignant peritoneal mesothelioma. Again, ACADL was identified as one of the most differentially expressed genes between the mesothelioma tumor and control, healthy tissue state, ranking 63 out of 47323 total transcripts. This was statistically significant (Table 1; $4.8E-16$).

ACADL mRNA is produced in significantly lower quantities in mesothelioma tissues than in normal lung and peritoneum.

Next, we extracted the exact mRNA expression values for ACADL in each of the datasets, for each patient tumor and each control tissue sample. In both datasets, ACADL was

expressed at lower quantities in mesothelioma tumor tissue. This was statistically significant for both the malignant pleural mesothelioma cohort (Figure 1; $p < 0.0001$) and for the diffuse malignant peritoneal cohort (Figure 1; $p < 0.0120$).

ACADL is differentially expressed in mesothelioma and it is expressed at significantly lower levels in mesothelioma tumor tissues than in normal lung and peritoneum.

Discussion

We performed systems-level whole transcriptome analysis of mesothelioma tumor tissues using two separate datasets (3, 4) from patients diagnosed with malignant pleural mesothelioma and from patients diagnosed with diffuse malignant peritoneal mesothelioma. We found that ACADL was out of over 50,000 transcripts the most differentially expressed gene in the mesothelioma transcriptome, and ranked 63 out of over 47,000 transcripts in another dataset.

There are zero targeted treatment options available for patients diagnosed with mesothelioma. ACADL catalyzes the first step in fatty acid oxidation in the mitochondria (10). Mitochondrial fatty acid oxidation (FAO) is the primary means by which energy is harvested by metabolism of fatty acids (11). The first step in mitochondrial FAO is an α - β dehydrogenation of the acyl-CoA ester by chain-length specific acyl-CoA dehydrogenases, one of which is long-chain acyl Co-A dehydrogenase ACADL (also known as LCAD) (10, 11). We identified ACADL as among the most differentially expressed genes across different mesothelioma types and found that it is expressed at significantly lower levels in mesothelioma tumor tissues than in normal lung and peritoneum. Manipulation of ACADL expression presents itself as a potential novel therapeutic modality and should be evaluated for its ability to halt, delay, or reverse mesothelioma progression *in vivo* (9).

References

1. Robinson, B.W., Musk, A.W. and Lake, R.A., 2005. Malignant mesothelioma. *The Lancet*, 366(9483), pp.397-408.
2. Beebe-Dimmer, J.L., Fryzek, J.P., Yee, C.L., Dalvi, T.B., Garabrant, D.H., Schwartz, A.G. and Gadgeel, S., 2016. Mesothelioma in the United States: a Surveillance, Epidemiology, and End Results (SEER)–Medicare investigation of treatment patterns and overall survival. *Clinical epidemiology*, 8, p.743.
3. Suraokar, M.B., Nunez, M.I., Diao, L., Chow, C.W., Kim, D., Behrens, C., Lin, H., Lee, S., Raso, G., Moran, C. and Rice, D., 2014. Expression profiling stratifies mesothelioma tumors and signifies deregulation of spindle checkpoint pathway and microtubule network with therapeutic implications. *Annals of oncology*, 25(6), pp.1184-1192.
4. Sciarillo, R., Wojtuszkiewicz, A., El Hassouni, B., Funel, N., Gandellini, P., Lagerweij, T., Buonamici, S., Blijlevens, M., van der Laan, E.A.Z., Zaffaroni, N. and Deraco, M., 2019. Splicing modulation as novel therapeutic strategy against diffuse malignant peritoneal mesothelioma. *EBioMedicine*, 39, pp.215-225.
5. Hiriart, E., Deepe, R. and Wessels, A., 2019. Mesothelium and malignant mesothelioma. *Journal of developmental biology*, 7(2), p.7.
6. Ryan, C.W., Herndon, J. and Vogelzang, N.J., 1998. A review of chemotherapy trials for malignant mesothelioma. *Chest*, 113(1), pp.66S-73S.
7. Vogelzang, N.J., Rusthoven, J.J., Symanowski, J., Denham, C., Kaukel, E., Ruffie, P., Gatzemeier, U., Boyer, M., Emri, S., Manegold, C. and Niyikiza, C., 2003. Phase III study of pemetrexed in combination with cisplatin versus cisplatin alone in patients with malignant pleural mesothelioma. *Journal of clinical oncology*, 21(14), pp.2636-2644.
8. Zalcman, G., Mazieres, J., Margery, J., Greillier, L., Audigier-Valette, C., Moro-Sibilot, D., Molinier, O., Corre, R., Monnet, I., Gounant, V. and Rivire, F., 2016. Bevacizumab for newly diagnosed pleural mesothelioma in the Mesothelioma Avastin Cisplatin Pemetrexed Study (MAPS): a randomised, controlled, open-label, phase 3 trial. *The Lancet*, 387(10026), pp. 1405-1414.
9. Bertino, P., Piccardi, F., Porta, C., Favoni, R., Cilli, M., Mutti, L. and Gaudino, G., 2008. Imatinib mesylate enhances therapeutic effects of gemcitabine in human malignant mesothelioma xenografts. *Clinical Cancer Research*, 14(2), pp.541-548.
10. Kurtz, D.M., Rinaldo, P., Rhead, W.J., Tian, L., Millington, D.S., Vockley, J., Hamm, D.A., Brix, A.E., Lindsey, J.R., Pinkert, C.A. and O'Brien, W.E., 1998. Targeted disruption of mouse long-chain acyl-CoA dehydrogenase gene reveals crucial roles for fatty acid oxidation. *Proceedings of the National Academy of Sciences*, 95(26), pp.15592-15597.
11. Mannaerts, G.P. and Debeer, L.J., 1982. Mitochondrial and peroxisomal β -oxidation of fatty acids in rat liver. *Annals of the New York Academy of Sciences*, 386(1), pp.30-39.

Rank	ID	p-value	t	B	Gene	Gene name
GSE51024						
1	206069_s_at	1.27E-40	-22.8046846	81.61521	ACADL	acyl-CoA dehydrogenase, long chain
GSE112154						
63	ILMN_1660890	4.8E-16	-1.2E+01	26.26571	ACADL	acyl-CoA dehydrogenase, long chain

Table 1: ACADL is one of the genes whose expression changes most significantly between the transcriptomes of mesothelioma tumor and normal tissues of the lung and peritoneum.

The rank of differential expression globally out of 54675 total transcripts (GSE51024) or 47323 total transcripts (GSE112154), probe ID, the p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between tumor and control groups, and gene symbol and gene name are shown in this chart (extracted from GEO2R).

Figure 1: ACADL mRNA is produced in significantly lower quantities in mesothelioma tissues than in normal lung and peritoneum.

The expression of ACADL in normal peritoneum (top row) or lung (bottom row) (green) and mesothelioma tumor tissues (red) is shown by bar graph with the error bars denoting the standard error of the mean (left) by a violin plot representing minimum to maximum expression with the dotted horizontal line denoting the median value and the finer lines denoting the 25th and 75th percent quartiles of expression (middle), and with dots representing the expression level in each individual patient and the dotted horizontal line representing the mean value (right). Data from GSE51024 is shown in the top row, while data from GSE112154 is shown in the bottom row. A statistical test was performed to evaluate the significance of expression between tumor (mesothelioma) and control (peritoneum or lung) groups, and the results of those tests, a p-value, is shown on each graph.