DETERMINANTAL FORMS AND RECURSIVE RELATIONS OF THE DELANNOY TWO-FUNCTIONAL SEQUENCE

FENG QI, MUHAMMET CIHAT DAĞLI, AND WEI-SHIH DU

Dedicated to people facing and battling COVID-19

ABSTRACT. In the paper, the authors establish closed forms for the Delannoy two-functional sequence and its difference in terms of the Hessenberg determinants, derive recursive relations for the Delannoy two-functional sequence and its difference, and deduce closed forms in terms of the Hessenberg determinants and recursive relations for the Delannoy one-functional sequence, the Delannoy numbers, and central Delannoy numbers.

CONTENTS

1. Introduction	1
2. Determinantal forms of the Delannoy two-functional sequence and its difference	4
3. Recursive relations of the Delannoy two-functional sequence and its difference	6
4. Determinantal forms and recursive relations for the Delannoy one-functional sequence, the Delannoy numbers, and central Delannoy numbers	7
5. Remarks	8
References	9

1. INTRODUCTION

A tridiagonal determinant is a determinant whose nonzero elements locate only on the diagonal and slots horizontally or vertically adjacent the diagonal. Technically speaking, a determinant $H = |h_{ij}|_{n \times n}$ is called a tridiagonal determinant if $h_{ij} = 0$ for all pairs (i, j) such that $|i - j| > 1$. For more information, please refer to the paper [11]. A determinant $H = |h_{ij}|_{n \times n}$ is called a lower (or an upper, respectively) Hessenberg determinant if $h_{ij} = 0$ for all pairs (i, j) such that $i + 1 < j$ (or $j + 1 < i$, respectively). For more information, please refer to the paper [18].

In combinatorial number theory, the Delannoy number $D(m, n)$ for $n, m \geq 0$ can be regarded as the number of lattice paths from $(0, 0)$ to (m, n) in which only east

2010 *Mathematics Subject Classification.* 05A10, 11B83, 11C20, 11Y55, 26C05.

Key words and phrases. closed form; recursive relation; difference; Hessenberg determinant; Delannoy two-functional sequence; Delannoy one-functional sequence; Delannoy number; central Delannoy number.

This paper was typeset using $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{L}\mathcal{A}\mathcal{T}\mathcal{E}\mathcal{X}$.

$(1, 0)$, north $(0, 1)$, and northeast $(1, 1)$. The Delannoy numbers $D(m, n)$ can be defined by means of the recurrence relation

$$D(m, n) = D(m-1, n) + D(m-1, n-1) + D(m, n-1)$$

with initial values $D(0, n) = D(m, 0) = D(0, 0) = 1$. The historic significance of these numbers $D(m, n)$ was explained in the paper [1]. The Delannoy numbers $D(m, n)$ can be computed by explicit formulas

$$D(m, n) = \sum_{k=0}^n \binom{n}{k} \binom{m}{k} 2^k \quad \text{and} \quad D(m, n) = \sum_{\ell=0}^n \binom{n}{\ell} \binom{m+n-\ell}{n}$$

and can be generated by

$$\frac{1}{1-x-y-xy} = \sum_{n,m \geq 0} D(m, n) x^m y^n.$$

For more information on the Delannoy numbers $D(m, n)$, please refer to [1, 19, 58, 62] and closely related references therein.

The numbers $D(k) = D(k, k)$ for $k \geq 0$ are known as central Delannoy numbers. In [26, Theorem 1.3], central Delannoy numbers $D(k)$ were represented by

$$D(k) = \frac{1}{\pi} \int_{3-2\sqrt{2}}^{3+2\sqrt{2}} \frac{1}{\sqrt{(t-3+2\sqrt{2})(3+2\sqrt{2}-t)}} \frac{1}{t^{k+1}} dt, \quad k \geq 0.$$

In [47, Section 4], the central Delannoy numbers $D(k)$ were generalized as

$$D_{a,b}(k) = \frac{1}{\pi} \int_a^b \frac{1}{\sqrt{(t-a)(b-t)}} \frac{1}{t^{k+1}} dt, \quad k \geq 0, \quad b > a > 0$$

and, from [26, Lemma 2.4], it is derived that the quantities $D_{a,b}(k)$ have the generating function

$$\frac{1}{\sqrt{(x+a)(x+b)}} = \sum_{k=0}^{\infty} D_{a,b}(k) x^k.$$

By virtue of conclusions in [24, Section 2.4] and [37, Remark 4.1], we can find that the quantities $D_{a,b}(k)$ for $k \geq 0$ can be computed by

$$D_{a,b}(k) = \frac{1}{G(a, b)} \frac{(-1)^k}{[2A(a, b)]^k} \sum_{\ell=0}^k (-1)^\ell 2^{2\ell} \frac{(2\ell-1)!!}{(2\ell)!!} \binom{\ell}{k-\ell} \left[\frac{A(a, b)}{H(a, b)} \right]^\ell$$

and

$$D_{a,b}(k) = \frac{1}{\sqrt{ab}} \frac{1}{b^k} \sum_{\ell=0}^k \frac{(2\ell-1)!!}{(2\ell)!!} \frac{[2(k-\ell)-1]!!}{[2(k-\ell)]!!} \left(\frac{b}{a} \right)^\ell,$$

where $A(a, b) = \frac{a+b}{2}$, $G(a, b) = \sqrt{ab}$, and $H(a, b) = \frac{2}{1/a+1/b}$ are respectively known as arithmetic, geometric, and harmonic means of $a, b > 0$. The quantities $D_{a,b}(k)$ can be further generalized as

$$D_{a,b;\lambda}(z) = \frac{\sin(\lambda\pi)}{\pi} \int_a^b \frac{1}{(t-a)^\lambda (b-t)^{1-\lambda}} \frac{1}{t^{z+1}} dt,$$

where $0 < a < b$, $\lambda \in (0, 1)$, and $z \in \mathbb{C}$. It is obvious that $D_{a,b;1/2}(k) = D_{a,b}(k)$. Theorems 2.1 and 2.2 in the paper [37] demonstrate that the quantities $D_{a,b;\lambda}(k)$ can be generated by

$$\frac{1}{(x+a)^\lambda(x+b)^{1-\lambda}} = \sum_{k=0}^{\infty} D_{a,b;\lambda}(k)x^k$$

and can be computed by the closed forms

$$D_{a,b;\lambda}(k) = \frac{1}{n!} \frac{1}{a^\lambda b^{1-\lambda}} \frac{1}{b^n} \sum_{\ell=0}^n \binom{n}{\ell} (\lambda)_\ell (1-\lambda)_{n-\ell} \left(\frac{b}{a}\right)^\ell$$

and

$$D_{a,b;\lambda}(k) = \frac{1}{a^\lambda b^{1-\lambda}} \frac{1}{b^n} \sum_{k=0}^n \left(\frac{b}{a}\right)^k \sum_{\ell=0}^k \langle \lambda \rangle_\ell \binom{k-1}{\ell-1} \frac{1}{\ell!} \left(1 - \frac{a}{b}\right)^\ell,$$

where

$$(x)_n = \begin{cases} \prod_{k=0}^{n-1} (x+k), & n \geq 1 \\ 1, & n = 0 \end{cases} \quad \text{and} \quad \langle x \rangle_n = \begin{cases} \prod_{k=0}^{n-1} (x-k), & n \geq 1 \\ 1, & n = 0 \end{cases}$$

are known as the rising and falling factorials respectively. The ideas, significance, and reasonability of these generalizations $D_{a,b}(k)$ and $D_{a,b;\lambda}(z)$ come from the papers [25, 33, 36, 37, 38, 42, 51, 52, 53, 54] and closely related references therein.

Inspired by the identity

$$\sum_{k=0}^n \binom{n}{k} \binom{x}{k} t^k = \sum_{k=0}^n \binom{n}{k} \binom{x+k}{n} (t-1)^k$$

in [8, eq. 3.17], the Delannoy numbers $D(m, n)$ were extended to the functional sequence

$$D(x; n) = \sum_{k=0}^n \binom{n}{k} \binom{x}{k} 2^k = \sum_{k=0}^n \binom{n}{k} \binom{x+k}{n}$$

in [60]. Various arithmetic properties and congruence relations for the Delannoy one-functional sequence $D(x; n)$ have been studied in [9, 16, 17, 58, 59, 62].

In [57], the Delannoy one-functional sequence $D(x; n)$ was further generalized as

$$D(x, r; n) = \sum_{k=0}^n \binom{x+r+k}{k} \binom{x-r}{n-k},$$

the generating function

$$\frac{(1+t)^{x-r}}{(1-t)^{x+r+1}} = \sum_{n=0}^{\infty} D(x, r; n) t^n, \quad |t| < 1 \quad (1.1)$$

was derived, and a plenty of identities for the Delannoy two-functional sequence $D(x, r; n)$ were acquired.

In [7], the Delannoy two-functional sequence $D(x, r; n)$ was generalized again to the Delannoy two-functional polynomials

$$D(x, r; n; y) = \sum_{k=0}^n \binom{x+r+k}{k} \binom{x-r}{n-k} y^k$$

and, among other things, the generating function and recurrence formula for the Delannoy two-functional polynomials $D(x, r; n; y)$ were derived.

It is noted that $D(x, 0; n) = D(x; n)$ and $D(x, r; n; 1) = D(x, r; n)$.

In combinatorial number theory, it is significant to express concrete sequences or arrays of integer numbers or polynomials in terms of tridiagonal determinants or the Hessenberg determinants. In this respect, the Bernoulli numbers and polynomials [2, 6, 20, 27, 35, 39, 61], the Euler numbers and polynomials [28, 31, 63], (central) Delannoy numbers and polynomials [10, 19, 25, 26, 30, 46, 47, 48], the Horadam polynomials [40], (generalized) Fibonacci numbers and polynomials [13, 14, 15, 21, 25, 32, 45, 49], the Lucas polynomials [40, 45], and the like, have been represented via tridiagonal determinants or the Hessenberg determinants, and consequently many remarkable relations have been obtained. For more information in this area and direction, please refer to [12, 22, 23, 29, 34, 41, 43, 44, 50, 55, 56] and closely related references therein.

In this paper, we will present closed forms [3] for the Delannoy two-functional sequence $D(x, r; n)$ and its difference $D(x, r; n) - D(x, r; n - 1)$ in terms of the Hessenberg determinants, derive recursive relations for the Delannoy two-functional sequence $D(x, r; n)$ and its difference $D(x, r; n) - D(x, r; n - 1)$, and deduce closed forms in terms of the Hessenberg determinants and recursive relations for the Delannoy one-functional sequence $D(x; n)$, the Delannoy numbers $D(m, n)$, and central Delannoy numbers $D(k)$.

2. DETERMINANTAL FORMS OF THE DELANNOY TWO-FUNCTIONAL SEQUENCE AND ITS DIFFERENCE

In this section, we will present closed forms for the Delannoy two-functional sequence $D(x, r; n)$ and for its difference $D(x, r; n) - D(x, r; n - 1)$ in terms of the Hessenberg determinants.

Theorem 2.1. *The Delannoy two-functional sequence $D(x, r; n)$ for $n \geq 0$ can be determinantly expressed by*

$$D(x, r; n) = \frac{(-1)^n}{n!} |P_{(n+1) \times 1}(x, r) \quad B_{(n+1) \times n}(x, r)|_{(n+1) \times (n+1)}, \quad (2.1)$$

where

$$P_{(n+1) \times 1}(x, r) = (\langle x - r \rangle_0 \quad \langle x - r \rangle_1 \quad \langle x - r \rangle_2 \quad \dots \quad \langle x - r \rangle_n)^T,$$

$$B_{(n+1) \times n}(x, r) = \left(\binom{i-1}{j-1} (-1)^{i-j} \langle x + r + 1 \rangle_{i-j} \right)_{(n+1) \times n},$$

and T denotes the transpose of a matrix.

The difference $D(x, r; n) - D(x, r; n - 1)$ for $n \geq 1$ can be determinantly expressed by

$$D(x, r; n) - D(x, r; n - 1) = \frac{(-1)^n}{n!} |P_{(n+1) \times 1} \quad C_{(n+1) \times n}|_{(n+1) \times (n+1)}, \quad (2.2)$$

where

$$C_{(n+1) \times n}(x, r) = \left(\binom{i-1}{j-1} (-1)^{i-j} \langle x + r \rangle_{i-j} \right)_{(n+1) \times n}.$$

Proof. Let $p_{x,r}(t) = (1+t)^{x-r}$ and $q_{x,r}(t) = (1-t)^{x+r+1}$. Then

$$p_{x,r}^{(k)}(t) = \langle x - r \rangle_k (1+t)^{x-r-k} \rightarrow \langle x - r \rangle_k$$

and

$$q_{x,r}^{(k)}(t) = (-1)^k \langle x+r+1 \rangle_k (1-t)^{x-r-k} \rightarrow (-1)^k \langle x+r+1 \rangle_k$$

as $t \rightarrow 0$.

The equation (1.1) implies that

$$D(x, r; n) = \frac{1}{n!} \lim_{t \rightarrow 0} \frac{d^n}{dt^n} \left[\frac{(1+t)^{x-r}}{(1-t)^{x+r+1}} \right]. \quad (2.3)$$

Let $u(t)$ and $v(t) \neq 0$ be two n th differentiable functions for $n \in \mathbb{N}$. Exercise 5) in [4, p. 40] reads that the n th derivative of the ratio $\frac{u(t)}{v(t)}$ can be computed by

$$\frac{d^n}{dx^n} \left[\frac{u(t)}{v(t)} \right] = (-1)^n \frac{|W_{(n+1) \times (n+1)}(t)|}{v^{n+1}(t)}, \quad (2.4)$$

where $|W_{(n+1) \times (n+1)}(t)|$ is the determinant of the $(n+1) \times (n+1)$ matrix

$$W_{(n+1) \times (n+1)}(t) = \begin{pmatrix} U_{(n+1) \times 1}(t) & V_{(n+1) \times n}(t) \end{pmatrix}_{(n+1) \times (n+1)},$$

the matrix $U_{(n+1) \times 1}(t)$ is an $(n+1) \times 1$ matrix whose elements satisfy $u_{k,1}(t) = u^{(k-1)}(t)$ for $1 \leq k \leq n+1$, and $V_{(n+1) \times n}(t)$ is an $(n+1) \times n$ matrix whose elements meet $v_{i,j}(t) = \binom{i-1}{j-1} v^{(i-j)}(t)$ for $1 \leq i \leq n+1$ and $1 \leq j \leq n$. The formula (2.4) is a general and fundamental, but non-extensively circulated, formula for derivatives of a ratio of two differentiable functions.

Combining (2.3) and (2.4) gives

$$\begin{aligned} D(x, r; n) &= \frac{(-1)^n}{n!} \lim_{t \rightarrow 0} \frac{|U_{(n+1) \times 1}(t) \quad V_{(n+1) \times n}(t)|}{q_{x,r}^{n+1}(t)} \\ &= \frac{(-1)^n}{n!} \frac{|\lim_{t \rightarrow 0} U_{(n+1) \times 1}(t) \quad \lim_{t \rightarrow 0} V_{(n+1) \times n}(t)|}{\lim_{t \rightarrow 0} q_{x,r}^{n+1}(t)} \\ &= \frac{(-1)^n}{n!} |\lim_{t \rightarrow 0} U_{(n+1) \times 1}(t) \quad \lim_{t \rightarrow 0} V_{(n+1) \times n}(t)|, \end{aligned}$$

where

$$\begin{aligned} \lim_{t \rightarrow 0} U_{(n+1) \times 1}(t) &= \lim_{t \rightarrow 0} \begin{pmatrix} p_{x,r}^{(0)}(t) & p_{x,r}^{(1)}(t) & p_{x,r}^{(2)}(t) & \dots & p_{x,r}^{(n)}(t) \end{pmatrix}^T \\ &= (\langle x-r \rangle_0 \quad \langle x-r \rangle_1 \quad \langle x-r \rangle_2 \quad \dots \quad \langle x-r \rangle_n)^T \end{aligned}$$

and

$$\begin{aligned} \lim_{t \rightarrow 0} V_{(n+1) \times n}(t) &= \left(\lim_{t \rightarrow 0} \binom{i-1}{j-1} q_{x,r}^{(i-j)}(t) \right)_{(n+1) \times n} \\ &= \left(\binom{i-1}{j-1} (-1)^{i-j} \langle x+r+1 \rangle_{i-j} \right)_{(n+1) \times n}. \end{aligned}$$

The determinantal form (2.1) is thus proved.

The generating function in (1.1) can be rewritten as

$$\begin{aligned} \frac{(1+t)^{x-r}}{(1-t)^{x+r}} &= (1-t) \sum_{n=0}^{\infty} D(x, r; n) t^n \\ &= D(x, r; 0) + \sum_{n=1}^{\infty} [D(x, r; n) - D(x, r; n-1)] t^n. \end{aligned}$$

By the same arguments as in the derivation of the determinantal form (2.1), we can obtain the determinantal forms (2.2) immediately. The proof of Theorem 2.1 is complete. $\square$

3. RECURSIVE RELATIONS OF THE DELANNOY TWO-FUNCTIONAL SEQUENCE AND ITS DIFFERENCE

In this section, with the aid of Theorem 2.1, we will derive recursive relations for the Delannoy two-functional sequence $D(x, r; n)$ and for its differences $D(x, r; n) - D(x, r; n-1)$.

Theorem 3.1. *The Delannoy two-functional sequence $D(x, r; n)$ for $n \geq 1$ satisfies*

$$D(x, r; n) = \frac{\langle x-r \rangle_n}{n!} + (-1)^n \sum_{s=1}^n (-1)^s \frac{\langle x+r+1 \rangle_{n-s+1}}{(n-s+1)!} D(x, r; s-1) \quad (3.1)$$

and the differences $D(x, r; n) - D(x, r; n-1)$ for $n \geq 2$ meet

$$\begin{aligned} D(x, r; n) - D(x, r; n-1) &= \frac{\langle x-r \rangle_n}{n!} \\ &+ (-1)^n \sum_{s=2}^n (-1)^s \frac{\langle x+r \rangle_{n-s+1}}{(n-s+1)!} [D(x, r; s-1) - D(x, r; s-2)]. \end{aligned} \quad (3.2)$$

Proof. Let $Q_0 = 1$ and

$$Q_n = \begin{vmatrix} e_{1,1} & e_{1,2} & 0 & \dots & 0 & 0 \\ e_{2,1} & e_{2,2} & e_{2,3} & \dots & 0 & 0 \\ e_{3,1} & e_{3,2} & e_{3,3} & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots & \vdots \\ e_{n-2,1} & e_{n-2,2} & e_{n-2,3} & \dots & e_{n-2,n-1} & 0 \\ e_{n-1,1} & e_{n-1,2} & e_{n-1,3} & \dots & e_{n-1,n-1} & e_{n-1,n} \\ e_{n,1} & e_{n,2} & e_{n,3} & \dots & e_{n,n-1} & e_{n,n} \end{vmatrix}$$

for $n \in \mathbb{N}$. In [5, p. 222, Theorem], it was proved that the sequence Q_n for $n \geq 0$ satisfies $Q_1 = e_{1,1}$ and

$$Q_n = \sum_{s=1}^n (-1)^{n-s} e_{n,s} \left(\prod_{j=s}^{n-1} e_{j,j+1} \right) Q_{s-1} \quad (3.3)$$

for $n \geq 2$, where any empty product is understood to be 1 while any empty sum is understood to be 0.

Substituting $(-1)^{n-1}(n-1)!D(x, r; n-1)$ for Q_n , replacing $e_{k,1}$ for $1 \leq k \leq n$ by $\langle x-r \rangle_{k-1}$, switching $e_{i,j}$ for $1 \leq i \leq n$ and $2 \leq j \leq n$ into $\binom{i-1}{j-2}(-1)^{i-j+1}\langle x+r+1 \rangle_{i-j+1}$ in (3.3), and simplifying arrive at

$$\begin{aligned} (-1)^{n-1}(n-1)!D(x, r; n-1) &= (-1)^{n-1}\langle x-r \rangle_{n-1} \\ &+ \sum_{s=2}^n (-1)^{n-s} \binom{n-1}{s-2} (-1)^{n-s+1} \langle x+r+1 \rangle_{n-s+1} (-1)^{s-2}(s-2)!D(x, r; s-2) \end{aligned}$$

which can be reformulated as

$$(-1)^{n-1}(n-1)!D(x, r; n-1) = (-1)^{n-1}\langle x-r \rangle_{n-1}$$

$$- \sum_{s=2}^n (-1)^s (s-2)! \binom{n-1}{s-2} \langle x+r+1 \rangle_{n-s+1} D(x, r; s-2).$$

The recursive relation (3.1) is thus proved.

Similarly, substituting $(-1)^{n-1}(n-1)![D(x, r; n-1) - D(x, r; n-2)]$ for Q_n , replacing $e_{k,1}$ for $1 \leq k \leq n$ by $\langle x-r \rangle_{k-1}$, switching $e_{i,j}$ for $1 \leq i \leq n$ and $2 \leq j \leq n$ into $\binom{i-1}{j-2}(-1)^{i-j+1}\langle x+r \rangle_{i-j+1}$ in (3.3), and straightforwardly simplifying lead to the recursive relation (3.2). The proof of Theorem 3.1 is complete. $\square$

4. DETERMINANTAL FORMS AND RECURSIVE RELATIONS FOR THE DELANNOY ONE-FUNCTIONAL SEQUENCE, THE DELANNOY NUMBERS, AND CENTRAL DELANNOY NUMBERS

In this section, with the help of Theorems 2.1 and 3.1, we deduce closed forms in terms of the Hessenberg determinants and recursive relations for the Delannoy one-functional sequence $D(x; n)$, the Delannoy numbers $D(m, n)$, and central Delannoy numbers $D(k)$.

Since $D(x, 0; n) = D(x; n)$, when taking $r = 0$ in Theorems 2.1 and 3.1, we derive the following conclusions.

Theorem 4.1. *The Delannoy one-functional sequence $D(x; n)$ for $n \geq 0$ can be determinantly expressed by*

$$D(x; n) = \frac{(-1)^n}{n!} |\mathfrak{P}_{(n+1) \times 1}(x) \quad \mathfrak{B}_{(n+1) \times n}(x)|_{(n+1) \times (n+1)}, \quad (4.1)$$

where

$$\mathfrak{P}_{(n+1) \times 1}(x) = (\langle x \rangle_0 \quad \langle x \rangle_1 \quad \langle x \rangle_2 \quad \dots \quad \langle x \rangle_n)^T$$

and

$$\mathfrak{B}_{(n+1) \times n}(x) = \left(\binom{i-1}{j-1} (-1)^{i-j} \langle x+1 \rangle_{i-j} \right)_{(n+1) \times n}.$$

The difference $D(x; n) - D(x; n-1)$ for $n \geq 1$ can be determinantly expressed by

$$D(x; n) - D(x; n-1) = \frac{(-1)^n}{n!} |\mathfrak{P}_{(n+1) \times 1}(x) \quad \mathfrak{C}_{(n+1) \times n}(x)|_{(n+1) \times (n+1)},$$

where

$$\mathfrak{C}_{(n+1) \times n}(x) = \left(\binom{i-1}{j-1} (-1)^{i-j} \langle x \rangle_{i-j} \right)_{(n+1) \times n}.$$

Theorem 4.2. *The Delannoy one-functional sequence $D(x; n)$ for $n \geq 1$ satisfies*

$$D(x; n) = \frac{\langle x \rangle_n}{n!} + (-1)^n \sum_{s=1}^n (-1)^s \frac{\langle x+1 \rangle_{n-s+1}}{(n-s+1)!} D(x; s-1)$$

and the differences $D(x; n) - D(x; n-1)$ for $n \geq 2$ meet

$$\begin{aligned} D(x; n) - D(x; n-1) &= \frac{\langle x \rangle_n}{n!} \\ &\quad + (-1)^n \sum_{s=2}^n (-1)^s \frac{\langle x \rangle_{n-s+1}}{(n-s+1)!} [D(x; s-1) - D(x; s-2)]. \end{aligned}$$

When taking $x = m$ in Theorems 4.3 and 4.4, we derive the following conclusions.

Theorem 4.3. *The Delannoy numbers $D(m; n)$ for $m, n \geq 0$ can be determinantal expressed by*

$$D(m; n) = \frac{(-1)^n}{n!} \left| \mathcal{P}_{(n+1) \times 1}(m) \quad \mathcal{B}_{(n+1) \times n}(m) \right|_{(n+1) \times (n+1)}, \quad (4.2)$$

where

$$\mathcal{P}_{(n+1) \times 1}(m) = (\langle m \rangle_0 \quad \langle m \rangle_1 \quad \langle m \rangle_2 \quad \dots \quad \langle m \rangle_n)^T$$

and

$$\mathcal{B}_{(n+1) \times n}(m) = \left((-1)^{i-j} \binom{i-1}{j-1} \langle m+1 \rangle_{i-j} \right)_{(n+1) \times n}.$$

The differences $D(m; n) - D(m; n-1)$ for $m \geq 0$ and $n \geq 1$ can be determinantal expressed by

$$D(m; n) - D(m; n-1) = \frac{(-1)^n}{n!} \left| \mathcal{P}_{(n+1) \times 1}(m) \quad \mathcal{C}_{(n+1) \times n}(m) \right|_{(n+1) \times (n+1)},$$

where

$$\mathcal{C}_{(n+1) \times n}(m) = \left(\binom{i-1}{j-1} (-1)^{i-j} \langle m \rangle_{i-j} \right)_{(n+1) \times n}.$$

Theorem 4.4. *The Delannoy numbers $D(m; n)$ for $m \geq 0$ and $n \geq 1$ satisfy*

$$D(m; n) = \binom{m}{n} + (-1)^{n-1} \sum_{s=0}^{n-1} (-1)^s \binom{m+1}{n-s} D(m; s) \quad (4.3)$$

and the differences $D(m; n) - D(m; n-1)$ for $m \geq 0$ and $n \geq 2$ meet

$$\begin{aligned} D(m; n) - D(m; n-1) &= \binom{m}{n} \\ &+ (-1)^n \sum_{s=2}^n (-1)^s \binom{m}{n-s+1} [D(m; s-1) - D(m; s-2)]. \end{aligned}$$

When taking $m = n = k$ in Theorem 4.3, we derive the following conclusions.

Theorem 4.5. *Central Delannoy numbers $D(k)$ for $k \geq 0$ can be determinantal expressed by*

$$D(k) = \frac{(-1)^k}{k!} \left| \mathcal{P}_{(k+1) \times 1} \quad \mathcal{B}_{(k+1) \times k} \right|_{(k+1) \times (k+1)}, \quad (4.4)$$

where

$$\mathcal{P}_{(k+1) \times 1} = (\langle k \rangle_0 \quad \langle k \rangle_1 \quad \langle k \rangle_2 \quad \dots \quad \langle k \rangle_k)^T$$

and

$$\mathcal{B}_{(k+1) \times k} = \left(\binom{i-1}{j-1} (-1)^{i-j} \langle k+1 \rangle_{i-j} \right)_{(k+1) \times k}.$$

5. REMARKS

The determinantal forms (4.2) and (4.4) in Theorem 4.3 and Theorem 4.5 recover [19, Theorem 2.1].

The formula (4.3) coincides with the first result in [19, Theorem 3.1]. Letting $m = n$ in (4.3) recovers the second result in [19, Theorem 3.1].

The determinantal form (4.4) in Theorem 4.5 is different from

$$D(k) = (-1)^k \begin{vmatrix} a_1 & 1 & 0 & \cdots & 0 & 0 & 0 \\ a_2 & a_1 & 1 & \cdots & 0 & 0 & 0 \\ a_3 & a_2 & a_1 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ a_{k-2} & a_{k-3} & a_{k-4} & \cdots & a_1 & 1 & 0 \\ a_{k-1} & a_{k-2} & a_{k-3} & \cdots & a_2 & a_1 & 1 \\ a_k & a_{k-1} & a_{k-2} & \cdots & a_3 & a_2 & a_1 \end{vmatrix}, \quad (5.1)$$

where

$$a_k = \frac{(-1)^{k+1}}{6^k} \sum_{\ell=1}^k (-1)^\ell 6^{2\ell} \frac{(2\ell-3)!!}{(2\ell)!!} \binom{\ell}{k-\ell}.$$

The determinantal form (5.1) was established in [26, Theorem 1.1]. See also related texts in the papers [19].

REFERENCES

- [1] C. Banderier and S. Schwer, *Why Delannoy numbers?*, J. Statist. Plann. Inference **135** (2005), no. 1, 40–54; available online at <https://doi.org/10.1016/j.jspi.2005.02.004>.
- [2] R. Booth and H. D. Nguyen, *Bernoulli polynomials and Pascal's square*, Fibonacci Quart. **46/47** (2008/2009), no. 1, 38–47.
- [3] J. M. Borwein and R. E. Crandall, *Closed forms: what they are and why we care*, Notices Amer. Math. Soc. **60** (2013), no. 1, 50–65; available online at <https://doi.org/10.1090/noti936>.
- [4] N. Bourbaki, *Functions of a Real Variable, Elementary Theory*, Translated from the 1976 French original by Philip Spain. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 2004; available online at <https://doi.org/10.1007/978-3-642-59315-4>.
- [5] N. D. Cahill, J. R. D'Errico, D. A. Narayan, and J. Y. Narayan, *Fibonacci determinants*, College Math. J. **33** (2002), no. 3, 221–225; available online at <https://doi.org/10.2307/1559033>.
- [6] F. Costabile, F. Dell'Accio, M. I. Gualtieri, *A new approach to Bernoulli polynomials*, Rend. Mat. Appl. (7) **26** (2006), no. 1, 1–12.
- [7] M. C. Dağlı, *A new generalization of Delannoy numbers*, accepted for publication in Indian Journal of Pure and Applied Mathematics.
- [8] H. W. Gould, *Combinatorial Identities: A standardized set of tables listing 500 binomial coefficient summations*, Henry W. Gould, Morgantown, W.Va., 1972.
- [9] V. J. W. Guo, *Proof of Sun's conjectures on integer-valued polynomials*, J. Math. Anal. Appl. **444** (2016), no. 1, 182–191; available online at <https://doi.org/10.1016/j.jmaa.2016.06.028>.
- [10] B.-N. Guo and F. Qi, *Some explicit and recursive formulas of the large and little Schröder numbers*, Arab J. Math. Sci. **23** (2017), no. 2, 141–147; available online at <https://doi.org/10.1016/j.ajmsc.2016.06.002>.
- [11] V. Higgins and C. Johnson, *Inverse spectral problems for collections of leading principal submatrices of tridiagonal matrices*, Linear Algebra Appl. **489** (2016), 104–122; available online at <https://doi.org/10.1016/j.laa.2015.10.004>.
- [12] S. Hu and M.-S. Kim, *Two closed forms for the Apostol–Bernoulli polynomials*, Ramanujan J. **46** (2018), no. 1, 103–117; available online at <https://doi.org/10.1007/s11139-017-9907-4>.
- [13] A. İpek and K. Ari, *On Hessenberg and pentadiagonal determinants related with Fibonacci and Fibonacci-like numbers*, Appl. Math. Comput. **229** (2014), 433–439; available online at <https://doi.org/10.1016/j.amc.2013.12.071>.
- [14] M. Janjić, *Determinants and recurrence sequences*, J. Integer Seq. **15** (2012), no. 3, Article 12.3.5, 21 pages.
- [15] R. K. Kittappa, *A representation of the solution of the n th order linear difference equation with variable coefficients*, Linear Algebra Appl. **193** (1993), 211–222; available online at [https://doi.org/10.1016/0024-3795\(93\)90278-V](https://doi.org/10.1016/0024-3795(93)90278-V).

- [16] J.-C. Liu, *A supercongruence involving Delannoy numbers and Schröder numbers*, J. Number Theory **168** (2016), 117–127; available online at <https://doi.org/10.1016/j.jnt.2016.04.019>.
- [17] J.-C. Liu, L. Li, and S.-D. Wang, *Some congruences on Delannoy numbers and Schröder numbers*, Int. J. Number Theory **14** (2018), no. 7, 2035–2041; available online at <https://doi.org/10.1142/S1793042118501221>.
- [18] R. S. Martin and J. H. Wilkinson, *Handbook Series Linear Algebra: Similarity reduction of a general matrix to Hessenberg form*, Numer. Math. **12** (1968), no. 5, 349–368; available online at <https://doi.org/10.1007/BF02161358>.
- [19] F. Qi, *A determinantal expression and a recursive relation of the Delannoy numbers*, arXiv preprint (2020), available online at <https://arxiv.org/abs/2003.12572v1>; Acta Univ. Sapientiae Math. **12** (2020), no. 2, in press.
- [20] F. Qi, *A double inequality for the ratio of two non-zero neighbouring Bernoulli numbers*, J. Comput. Appl. Math. **351** (2019), 1–5; available online at <https://doi.org/10.1016/j.cam.2018.10.049>.
- [21] F. Qi, *Denying a short proof of a determinantal formula for generalized Fibonacci polynomials*, J. Math. Anal. **11** (2020), no. 1, 52–57.
- [22] F. Qi, *Determinantal expressions and recurrence relations for Fubini and Eulerian polynomials*, J. Interdiscip. Math. **22** (2019), no. 3, 317–335; available online at <https://doi.org/10.1080/09720502.2019.1624063>.
- [23] F. Qi, *Derivatives of tangent function and tangent numbers*, Appl. Math. Comput. **268** (2015), 844–858; available online at <https://doi.org/10.1016/j.amc.2015.06.123>.
- [24] F. Qi and V. Čerňanová, *Some discussions on a kind of improper integrals*, Int. J. Anal. Appl. **11** (2016), no. 2, 101–109.
- [25] F. Qi, V. Čerňanová, and Y. S. Semenov, *Some tridiagonal determinants related to central Delannoy numbers, the Chebyshev polynomials, and the Fibonacci polynomials*, Politehn. Univ. Bucharest Sci. Bull. Ser. A Appl. Math. Phys. **81** (2019), no. 1, 123–136.
- [26] F. Qi, V. Čerňanová, X.-T. Shi, and B.-N. Guo, *Some properties of central Delannoy numbers*, J. Comput. Appl. Math. **328** (2018), 101–115; available online at <https://doi.org/10.1016/j.cam.2017.07.013>.
- [27] F. Qi and R. J. Chapman, *Two closed forms for the Bernoulli polynomials*, J. Number Theory **159** (2016), 89–100; available online at <https://doi.org/10.1016/j.jnt.2015.07.021>.
- [28] F. Qi and B.-N. Guo, *A determinantal expression and a recurrence relation for the Euler polynomials*, Adv. Appl. Math. Sci. **16** (2017), no. 9, 297–309.
- [29] F. Qi and B.-N. Guo, *A diagonal recurrence relation for the Stirling numbers of the first kind*, Appl. Anal. Discrete Math. **12** (2018), no. 1, 153–165; available online at <https://doi.org/10.2298/AADM170405004Q>.
- [30] F. Qi and B.-N. Guo, *Explicit and recursive formulas, integral representations, and properties of the large Schröder numbers*, Kragujevac J. Math. **41** (2017), no. 1, 121–141; available online at <https://doi.org/10.5937/KgJMath1701121F>.
- [31] F. Qi and B.-N. Guo, *Explicit formulas for special values of the Bell polynomials of the second kind and for the Euler numbers and polynomials*, Mediterr. J. Math. **14** (2017), no. 3, Article 140, 14 pages; available online at <https://doi.org/10.1007/s00009-017-0939-1>.
- [32] F. Qi and B.-N. Guo, *Expressing the generalized Fibonacci polynomials in terms of a tridiagonal determinant*, Matematiche (Catania) **72** (2017), no. 1, 167–175; available online at <https://doi.org/10.4418/2017.72.1.13>.
- [33] F. Qi and B.-N. Guo, *Lévy-Khintchine representation of Toader–Qi mean*, Math. Inequal. Appl. **21** (2018), no. 2, 421–431; available online at <https://doi.org/10.7153/mia-2018-21-29>.
- [34] F. Qi and B.-N. Guo, *Several explicit and recursive formulas for generalized Motzkin numbers*, AIMS Math. **5** (2020), no. 2, 1333–1345; available online at <https://doi.org/10.3934/math.2020091>.
- [35] F. Qi and B.-N. Guo, *Some determinantal expressions and recurrence relations of the Bernoulli polynomials*, Mathematics **4** (2016), no. 4, Article 65, 11 pages; available online at <https://doi.org/10.3390/math4040065>.
- [36] F. Qi and B.-N. Guo, *The reciprocal of the geometric mean of many positive numbers is a Stieltjes transform*, J. Comput. Appl. Math. **311** (2017), 165–170; available online at <https://doi.org/10.1016/j.cam.2016.07.006>.

- [37] F. Qi and B.-N. Guo, *The reciprocal of the weighted geometric mean is a Stieltjes function*, Bol. Soc. Mat. Mex. (3) **24** (2018), no. 1, 181–202; available online at <https://doi.org/10.1007/s40590-016-0151-5>.
- [38] F. Qi and B.-N. Guo, *The reciprocal of the weighted geometric mean of many positive numbers is a Stieltjes function*, Quaest. Math. **41** (2018), no. 5, 653–664; available online at <https://doi.org/10.2989/16073606.2017.1396508>.
- [39] F. Qi and B.-N. Guo, *Two nice determinantal expressions and a recurrence relation for the Apostol–Bernoulli polynomials*, J. Indones. Math. Soc. (MIHMI) **23** (2017), no. 1, 81–87; available online at <https://doi.org/10.22342/jims.23.1.274.81-87>.
- [40] F. Qi, C. Kizilates, and W.-S. Du, *A closed formula for the Horadam polynomials in terms of a tridiagonal determinant*, Symmetry **11** (2019), no. 6, 8 pages; available online at <https://doi.org/10.3390/sym11060782>.
- [41] F. Qi, O. Kouba, and I. Kaddoura, *Computation of several Hessenberg determinants*, Math. Slovaca **70** (2020), in press.
- [42] F. Qi and D. Lim, *Integral representations of bivariate complex geometric mean and their applications*, J. Comput. Appl. Math. **330** (2018), 41–58; available online at <https://doi.org/10.1016/j.cam.2017.08.005>.
- [43] F. Qi, M. Mahmoud, X.-T. Shi, and F.-F. Liu, *Some properties of the Catalan–Qi function related to the Catalan numbers*, SpringerPlus (2016), **5**:1126, 20 pages; available online at <https://doi.org/10.1186/s40064-016-2793-1>.
- [44] F. Qi, D.-W. Niu, and B.-N. Guo, *Some identities for a sequence of unnamed polynomials connected with the Bell polynomials*, Rev. R. Acad. Cienc. Exactas Fís. Nat. Ser. A Math. RACSAM **113** (2019), no. 2, 557–567; available online at <https://doi.org/10.1007/s13398-018-0494-z>.
- [45] F. Qi, E. Polatli, and B.-N. Guo, *Determinantal formulas and recurrent relations for bi-periodic Fibonacci and Lucas polynomials*, OSF Preprints (2020), available online at <https://doi.org/10.31219/osf.io/qvxd8>.
- [46] F. Qi, X.-T. Shi, and B.-N. Guo, *Integral representations of the large and little Schröder numbers*, Indian J. Pure Appl. Math. **49** (2018), no. 1, 23–38; available online at <https://doi.org/10.1007/s13226-018-0258-7>.
- [47] F. Qi, X.-T. Shi, and B.-N. Guo, *Some properties of the Schröder numbers*, Indian J. Pure Appl. Math. **47** (2016), no. 4, 717–732; available online at <https://doi.org/10.1007/s13226-016-0211-6>.
- [48] F. Qi, X.-T. Shi, and B.-N. Guo, *Two explicit formulas of the Schröder numbers*, Integers **16** (2016), Paper No. A23, 15 pages.
- [49] F. Qi, J.-L. Wang, and B.-N. Guo, *A determinantal expression for the Fibonacci polynomials in terms of a tridiagonal determinant*, Bull. Iranian Math. Soc. **45** (2019), no. 6, 1821–1829; available online at <https://doi.org/10.1007/s41980-019-00232-4>.
- [50] F. Qi, J.-L. Wang, and B.-N. Guo, *A representation for derangement numbers in terms of a tridiagonal determinant*, Kragujevac J. Math. **42** (2018), no. 1, 7–14; available online at <https://doi.org/10.5937/KgJMath1801007F>.
- [51] F. Qi, X.-J. Zhang, and W.-H. Li, *An elementary proof of the weighted geometric mean being a Bernstein function*, Politehn. Univ. Bucharest Sci. Bull. Ser. A Appl. Math. Phys. **77** (2015), no. 1, 35–38.
- [52] F. Qi, X.-J. Zhang, and W.-H. Li, *An integral representation for the weighted geometric mean and its applications*, Acta Math. Sin. (Engl. Ser.) **30** (2014), no. 1, 61–68; available online at <https://doi.org/10.1007/s10114-013-2547-8>.
- [53] F. Qi, X.-J. Zhang, and W.-H. Li, *Lévy–Khintchine representation of the geometric mean of many positive numbers and applications*, Math. Inequal. Appl. **17** (2014), no. 2, 719–729; available online at <https://doi.org/10.7153/mia-17-53>.
- [54] F. Qi, X.-J. Zhang, and W.-H. Li, *Lévy–Khintchine representations of the weighted geometric mean and the logarithmic mean*, Mediterr. J. Math. **11** (2014), no. 2, 315–327; available online at <https://doi.org/10.1007/s00009-013-0311-z>.
- [55] F. Qi and J.-L. Zhao, *Some properties of the Bernoulli numbers of the second kind and their generating function*, Bull. Korean Math. Soc. **55** (2018), no. 6, 1909–1920; available online at <https://doi.org/10.4134/BKMS.b180039>.

- [56] F. Qi, J.-L. Zhao, and B.-N. Guo, *Closed forms for derangement numbers in terms of the Hessenberg determinants*, Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Mat. RACSAM **112** (2018), no. 4, 933–944; available online at <https://doi.org/10.1007/s13398-017-0401-z>.
- [57] Z.-H. Sun, *A kind of orthogonal polynomials and related identities*, J. Math. Anal. Appl. **456** (2017), no. 2, 912–926; available online at <https://doi.org/10.1016/j.jmaa.2017.07.049>.
- [58] Z.-W. Sun, *Arithmetic properties of Delannoy numbers and Schröder numbers*, J. Number Theory **183** (2018), 146–171; available online at <https://doi.org/10.1016/j.jnt.2017.07.011>.
- [59] Z.-W. Sun, *On Delannoy numbers and Schröder numbers*, J. Number Theory **131** (2011), no. 12, 2387–2397; available online at <https://doi.org/10.1016/j.jnt.2011.06.005>.
- [60] Z.-W. Sun, *Supercongruences involving dual sequences*, Finite Fields Appl. **46** (2017), 179–216; available online at <https://doi.org/10.1016/j.ffa.2017.03.007>.
- [61] H. W. Turnbull, *The Theory of Determinants, Matrices, and Invariants*, 3rd ed., Dover Publications, New York, 1960.
- [62] Y. Wang, S.-N. Zheng, and X. Chen, *Analytic aspects of Delannoy numbers*, Discrete Math. **342** (2019), no. 8, 2270–2277; available online at <https://doi.org/10.1016/j.disc.2019.04.003>.
- [63] C.-F. Wei and F. Qi, *Several closed expressions for the Euler numbers*, J. Inequal. Appl. **2015**, 2015:219, 8 pages; available online at <https://doi.org/10.1186/s13660-015-0738-9>.

INSTITUTE OF MATHEMATICS, HENAN POLYTECHNIC UNIVERSITY, JIAOZUO 454010, HENAN, CHINA; COLLEGE OF MATHEMATICS AND PHYSICS, INNER MONGOLIA UNIVERSITY FOR NATIONALITIES, TONGLIAO 028043, INNER MONGOLIA, CHINA; SCHOOL OF MATHEMATICAL SCIENCES, TIANJIN POLYTECHNIC UNIVERSITY, TIANJIN 300387, CHINA

Email address: qifeng618@gmail.com, qifeng618@hotmail.com, qifeng618@qq.com
URL: <https://qifeng618.wordpress.com>, <https://orcid.org/0000-0001-6239-2968>

DEPARTMENT OF MATHEMATICS, AKDENİZ UNIVERSITY, 07058-ANTALYA, TURKEY

Email address: mcihatdagli@akdeniz.edu.tr
URL: <https://orcid.org/0000-0003-2859-902X>

DEPARTMENT OF MATHEMATICS, NATIONAL KAOHSIUNG NORMAL UNIVERSITY, KAOHSIUNG 82444, TAIWAN

Email address: wsdu@mail.nknu.edu.tw
URL: <https://orcid.org/0000-0001-8996-2270>