Differential expression of topoisomerase II α in cancers of the breast.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY 11730

Breast cancer affects women at relatively high frequency¹. We mined published microarray datasets^{2,3} to determine in an unbiased fashion and at the systems level genes most differentially expressed in the primary tumors of patients with breast cancer. We identified significant differential expression of the gene encoding the DNA topoisomerase II α , TOP2A, when comparing primary tumors of the breast to the tissue of origin, the normal breast. Expression of TOP2A in primary tumors of the breast was significantly higher than in normal breast tissue, and patients with low expression of TOP2A in primary tumors had markedly superior overall and relapse-free survival. TOP2A may be of relevance to initiation, maintenance or progression of cancers of the female breast.

Keywords: breast cancer, TOP2A, DNA topoisomerase II α , systems biology of breast cancer, targeted therapeutics in breast cancer.

Invasive breast cancer is diagnosed in over a quarter of a million women in the United States each year¹ and in 2018, breast cancer was the leading cause of cancer death in women worldwide⁴. While patients with localized breast cancer are provided a 99% 5-year survival rate, patients with regional breast cancer, cancer that has spread to lymph nodes or nearby structures, are provided an 86% 5-year survival rate^{5,6}. Patients with metastasis to distant sites are provided a 27% 5-year survival rate^{5,6}. Understanding how primary tumors are most transcriptionally different from the tissue from which they originate, the breast, can facilitate development of novel diagnostic and therapeutics to promote early detection and enhanced treatment, and contribute to efforts to prevent progression to metastatic stages. We mined published microarray data^{2,3} to understand at the transcriptome level and in an unbiased fashion genes most differentially expressed in primary tumors of the breast as compared to normal breast tissue. The DNA topoisomerase II α , encoded by the TOP2A gene, emerged as among the most differentially expressed genes in cancer of the female breast.

Methods

We utilized datasets GSE109169² and GSE124648³ for this differential gene expression analysis of female breast cancer. GSE109169 was generated using Affymetrix Human Exon 1.0 ST Array technology with $n=25$ normal breast tissue and $n=25$ tumors of the breast and; analysis was performed using platform GPL5175. The tissues from this first dataset are paired tissues (25 tumors matching 25 breast tissues from 25 patients). GSE124648 was generated using Affymetrix Human Genome U133A Array technology with $n=10$ normal breast tissue and $n=126$ tumors of the breast and; analysis was performed using platform GPL96. The Benjamini and Hochberg method of p -value adjustment was used for ranking of differential

expression but raw p -values were used to assess statistical significance of global differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. A statistical test was performed to evaluate whether TOP2A expression was significantly between primary breast tumors and breast tissue using a two-tailed, unpaired t-test with Welch's correction. We used PRISM for all statistical analyses of differential gene expression in human breast cancer (Version 8.4.0)(455). For Kaplan-Meier survival analysis, we used the Kaplan-Meier plotter online tool⁷ for correlation of TOP2A mRNA expression levels with relapse-free survival (disease-free survival) in $n=3951$ breast cancer patients, and overall survival in $n=1402$ breast cancer patients.

Results

We performed discovery of genes associated with breast cancer in females by mining two independently published microarray datasets^{2,3}.

TOP2A is differentially expressed in primary tumors of the breast.

Comparison of 25 normal breast tissues to 25 tumors of the breast² revealed that the DNA topoisomerase II α , TOP2A, was among the genes most differentially expressed in tumors of the breast in female breast cancer (Table 1). When sorting each of the genes expressed in tumors of the breast based on significance of difference as compared to normal breast tissue, TOP2A ranked 220 out of 19076 total transcripts, equating to 98.8% differential expression (Table 1). Differential expression of TOP2A in female breast cancer was statistically significant (Table 1; $p=6.64E-13$).

Analysis of a second microarray dataset³, here in 126 tumors of the breast as compared

to 10 normal breast tissues, again revealed significant differential expression of TOP2A in tumors of the breast in female breast cancer (Table 2). When sorting each of the genes expressed in tumors of the breast based on significance of difference as compared to normal breast tissue, TOP2A ranked 127 out of 22283 total transcripts, equating to 99.4% differential expression (Table 2). Differential expression of TOP2A in female breast cancer was statistically significant (Table 2; $p=4.69E-16$).

TOP2A is expressed at significantly higher levels in breast tumors as compared to the breast.

We obtained exact mRNA expression levels for TOP2A from the breast and from breast tumors to understand the magnitude and direction of TOP2A expression change. In both datasets, TOP2A was expressed at higher levels in tumors of the breast as compared to normal breast tissue (Figure 1 and Figure 2). Increased expression of TOP2A in primary breast tumors was statistically significant (Figure 1: $p<0.0001$ and Figure 2: $p<0.0001$). We calculated a mean fold change of 1.5 ± 0.3 and 1.728 ± 0.242 when comparing primary tumors of the breast to normal breast tissues (Table 1 and Table 2; respectively).

TOP2A expression associates with survival outcomes in female breast cancer.

We performed Kaplan-Meier survival analysis in an attempt to correlate TOP2A mRNA expression levels with patient survival, in 3951 breast cancer patients for relapse-free survival correlation analysis or 1402 breast cancer patients for overall survival analysis. We found a statistically significant correlation between TOP2A expression and both overall survival as well as relapse-free survival (also known as disease-free survival) in patients with breast cancer

(Figure 3 and 4; log rank p -value=3.1e-08 for overall survival, hazard ratio: 1.84 (1.48-2.29) (Fig. 3) and log rank p -value < 1e-16, and hazard ratio: 2 (1.78 - 2.23) for relapse-free survival (Fig 4)). Higher TOP2A mRNA levels were a negative prognostic indicator. Median OS for patients whose tumors expressed low levels of TOP2A was 169.2 months, while median OS for patients whose tumors expressed high levels of TOP2A was 67.4 months (Table 3). Median RFS was 216.66 months for patients whose tumors expressed low levels of TOP2A while median RFS was 163.46 months for patients whose tumors expressed high levels of TOP2A (Table 4).

Thus, through comparative transcriptome analysis of primary tumors of the breast and normal breast tissue, we found that differential expression and up-regulation of TOP2A was among the most significant transcriptional features in primary tumors from patients with breast cancer. TOP2A expression in primary tumors of the breast demonstrated a strong and significant correlation with overall survival as well as relapse-free survival in patients with breast cancer, with lower expression of TOP2A correlating with improved survival.

Discussion

Invasive breast cancer is medical problem with a 27% 5-year survival rate for women whose disease has spread to distant sites. To facilitate understanding of the basic transcriptional differences between primary tumors of the breast and the tissues from which these tumors originate, normal breast tissues, we performed comparative transcriptome analysis using two independently published microarray datasets^{2,3} and identified significant differential expression of the DNA topoisomerase IIa, TOP2A, in breast cancer.

1 DNA topoisomerase function to resolve supercoiling in the double-stranded DNA helix
2 during DNA replication and transcription of mRNAs. Human DNA TopoII α contains an amino-
3 terminal ATPase domain that utilizes energy derived from ATP during unwinding of supercoiling,
4 a centrally located DNA-binding and cleavage domain, as well as a carboxy-terminal tail⁸.
5 Relief of supercoiling involves binding of a dimeric enzyme to the helix, cleavage of the
6 opposing strand creating a gated or G-segment, and a conformational change to allow, in a
7 relaxation reaction, passage of a transported or T-segment through the gate created by
8 cleavage⁸.
9

10
11 Co-amplification of the TOP2A gene, along with HER2, has been reported in both urinary
12 bladder cancer⁹ and in colorectal cancer¹⁰. Multiple studies have reported aberrations at the
13 TOP2A locus in breast cancer, either copy number gain (amplification) or loss (deletion). One
14 study from the Danish Breast Cancer Cooperative Group reported copy number variation at the
15 TOP2A locus in 23% of 773 tumors surveyed, with 12% in total harboring TOP2A amplification
16 and 11% harboring TOP2A deletion¹¹. Patients with TOP2A amplification possessed
17 significantly enhanced recurrence-free and overall survival if treated with cyclophosphamide,
18 epirubicin and fluorouracil (CEF) as compared to patients treated with cyclophosphamide,
19 methotrexate and fluorouracil (CMF)¹¹. While this might be interpreted as an indication of
20 superior outcome to CEF chemotherapy based on amplification of TOP2A, we argue this
21 conclusion would be misleading as we find that patients with low expression of TOP2A
22 possessed significantly greater overall and relapse-free survival as compared to patients with
23 high expression of TOP2A, suggesting that the statistical difference observed was not an
24 indication of superior chemotherapeutic response, attributable to unique functions of CEF on a
25 TOP2A amplified genetic background, but an improvement over inferior survival associated with
26
27
28

TOP2A genotype: high expression of TOP2A resulting from amplification, and attributable to any number of other factors or pathways differentially affected by CEF versus CMF. Of note, the effect of chemotherapeutic regimen on overall and recurrence-free survival was not observed in patients with normal TOP2A genotype. A follow-up (“update”) to the Danish Breast Cancer Cooperative Group that when clustered together as a group, TOP2A aberrations, both deletions and amplifications, were associated with significantly shorter recurrence-free and overall survival¹². Moreover, this follow-up described worse prognosis for patients with deletions as compared to patients with amplifications¹². There are two issues here: first, it is misleading to cluster together patients with TOP2A amplifications and deletions as one group and attempt to derive a correlation. Second, and more importantly, we find that patients with low expression of TOP2A (like patients with TOP2A deletions) possess significantly greater, and not worse, overall and recurrence-free survival than patients with high expression of TOP2A (like patients with TOP2A amplification).

Co-amplification of TOP2A in conjunction with HER2 or in association with HER2 status has been described in multiple studies¹³⁻¹⁵; in one study of 670 breast cancer cases, it was concluded that while changes in TOP2A were associated with HER2 status, TOP2A was of limited value as a prognostic marker. Finally, in part based on unsubstantiated rationale that TOP2A is a target or the target of anthracyclines in patients with breast cancer (i.e., that poisoning of TOP2A by anthracyclines is what confers efficacy to anthracyclines in patients with breast cancer¹⁶), multiple studies/analyses have investigated relationships between anthracycline chemotherapy and TOP2A genomic status¹⁷⁻¹⁹. One concluded that TOP2A amplification status was not significantly associated with response to anthracyclines¹⁶. Another study, using pooled data from five adjuvant trials, concluded that patients with TOP2A

1 aberrations benefited from adjuvant anthracycline chemotherapy¹⁷. A third meta-analysis
2 analyzing data from 3102 patients concluded that while there was some value of TOP2A
3 genomic status to predicting response to anthracyclines, the data did not support the use of
4 anthracyclines based on amplification or deletion of TOP2A¹⁸. Thus, changes in copy number at
5 the TOP2A locus - amplifications and deletions - have been reported in human breast cancer,
6 alone and in conjunction with HER2 copy number changes; it has been argued that TOP2A
7 genomic status could predict benefit to adjuvant epirubicin (i.e., CEF vs. CMF), that patients
8 with TOP2A aberrations had significantly worse survival outcomes than those without, but that
9 patients with TOP2A deletions had significantly worse survival than those with amplifications,
10 and multiple studies have arrived at contradictory conclusions as to the utility of TOP2A genomic
11 status in predicting response to adjuvant anthracycline.

12
13
14 We found significant differential expression and up-regulation of the DNA topoisomerase
15 II α , TOP2A in human breast cancer across independent microarray datasets^{2,3}. TOP2A
16 expression was significantly correlated with both overall and relapse-free survival in patients
17 with breast cancer; we found that the median overall survival for breast cancer patients with low
18 tumor expression of TOP2A was more than 8 years greater than that for patients with high tumor
19 expression of TOP2A. Future studies can determine by which mechanisms DNA topoisomerase
20 II α might function to contribute to breast cancer initiation or progression, and which of these are
21 the most significant with respect to driving aggressive behavior.
22
23
24
25
26
27
28

References

1. DeSantis, C.E., Ma, J., Goding Sauer, A., Newman, L.A. and Jemal, A., 2017. Breast cancer statistics, 2017, racial disparity in mortality by state. *CA: a cancer journal for clinicians*, 67(6), pp.439-448.
2. Chang, J.W., Kuo, W.H., Lin, C.M., Chen, W.L., Chan, S.H., Chiu, M.F., Chang, I.S., Jiang, S.S., Tsai, F.Y., Chen, C.H. and Huang, P.H., 2018. Wild-type p53 upregulates an early onset breast cancer-associated gene GAS7 to suppress metastasis via GAS7–CYFIP1-mediated signaling pathway. *Oncogene*, 37(30), pp.4137-4150.
3. Sinn, B.V., Fu, C., Lau, R., Litton, J., Tsai, T.H., Murthy, R., Tam, A., Andreopoulou, E., Gong, Y., Murthy, R. and Gould, R., 2019. SET ER/PR: a robust 18-gene predictor for sensitivity to endocrine therapy for metastatic breast cancer. *NPJ breast cancer*, 5(1), pp. 1-8.
4. Bray, F., Ferlay, J., Soerjomataram, I., Siegel, R.L., Torre, L.A. and Jemal, A., 2018. Global cancer statistics 2018: GLOBOCAN estimates of incidence and mortality worldwide for 36 cancers in 185 countries. *CA: a cancer journal for clinicians*, 68(6), pp.394-424.
5. ACS Cancer Facts & Figures 2019. <https://www.cancer.net/cancer-types/breast-cancer-metastatic/statistics>.
6. Survival Rates for Breast Cancer. <https://www.cancer.org/cancer/breast-cancer/understanding-a-breast-cancer-diagnosis/breast-cancer-survival-rates.html>
7. Györfy, B., Lanczky, A., Eklund, A.C., Denkert, C., Budczies, J., Li, Q. and Szallasi, Z., 2010. An online survival analysis tool to rapidly assess the effect of 22,277 genes on breast cancer prognosis using microarray data of 1,809 patients. *Breast cancer research and treatment*, 123(3), pp.725-731.
8. Champoux, J.J., 2001. DNA topoisomerases: structure, function, and mechanism. *Annual review of biochemistry*, 70(1), pp.369-413.
9. Simon, R., Atefy, R., Wagner, U., Forster, T., Fijan, A., Bruderer, J., Wilber, K., Mihatsch, M.J., Gasser, T. and Sauter, G., 2003. HER-2 and TOP2A coamplification in urinary bladder cancer. *International journal of cancer*, 107(5), pp.764-772.

10. Al-Kuraya, K., Novotny, H., Bavi, P., Siraj, A.K., Uddin, S., Ezzat, A., Al Sanea, N., Al-Dayel, F., Al-Mana, H., Sheikh, S.S. and Mirlacher, M., 2007. HER2, TOP2A, CCND1, EGFR and C-MYC oncogene amplification in colorectal cancer. *Journal of clinical pathology*, 60(7), pp. 768-772.
11. Knoop, Ann S., Helle Knudsen, Eva Balslev, Birgitte B. Rasmussen, Jens Overgaard, Kirsten V. Nielsen, Andreas Schonau et al. "Retrospective analysis of topoisomerase IIa amplifications and deletions as predictive markers in primary breast cancer patients randomly assigned to cyclophosphamide, methotrexate, and fluorouracil or cyclophosphamide, epirubicin, and fluorouracil: Danish Breast Cancer Cooperative Group." *Journal of Clinical Oncology* 23, no. 30 (2005): 7483-7490.
12. Vang Nielsen, K., Ejlersen, B., Møller, S., Trøst Jørgensen, J., Knoop, A., Knudsen, H. and Mouridsen, H.T., 2008. The value of TOP2A gene copy number variation as a biomarker in breast cancer: Update of DBCG trial 89D. *Acta oncologica*, 47(4), pp.725-734.
13. Arriola, E., Marchio, C., Tan, D.S., Drury, S.C., Lambros, M.B., Natrajan, R., Rodriguez-Pinilla, S.M., Mackay, A., Tamber, N., Fenwick, K. and Jones, C., 2008. Genomic analysis of the HER2/TOP2A amplicon in breast cancer and breast cancer cell lines. *Laboratory investigation*, 88(5), pp.491-503.
14. Engstrøm, M.J., Ytterhus, B., Vatten, L.J., Opdahl, S. and Bofin, A.M., 2014. TOP2A gene copy number change in breast cancer. *Journal of clinical pathology*, 67(5), pp.420-425.
15. Olsen, K.E., Knudsen, H., Rasmussen, B.B., Balslev, E., Knoop, A., Ejlersen, B., Nielsen, K.V., Schønau, A., Overgaard, J. and Danish Breast Cancer Co-operative Group, 2004. Amplification of HER2 and TOP2A and deletion of TOP2A genes in breast cancer investigated by new FISH probes. *Acta Oncologica*, 43(1), pp.35-42.
16. Munro, A.F., Cameron, D.A. and Bartlett, J.M.S., 2010. Targeting anthracyclines in early breast cancer: new candidate predictive biomarkers emerge. *Oncogene*, 29(38), pp. 5231-5240.
17. Villman, K., Sjöström, J., Heikkilä, R., Hultborn, R., Malmström, P., Bengtsson, N.O., Söderberg, M., Saksela, E. and Blomqvist, C., 2006. TOP2A and HER2 gene amplification as predictors of response to anthracycline treatment in breast cancer. *Acta oncologica*, 45(5), pp.590-596.

18. Bartlett, J.M., McConkey, C.C., Munro, A.F., Desmedt, C., Dunn, J.A., Larsimont, D.P., O'Malley, F.P., Cameron, D.A., Earl, H.M., Poole, C.J. and Shepherd, L.E., 2015. Predicting anthracycline benefit: TOP2A and CEP17-not only but also. *J Clin Oncol*, 33(15), pp. 1680-1687.
19. Di Leo, A., Desmedt, C., Bartlett, J.M., Piette, F., Ejlertsen, B., Pritchard, K.I., Larsimont, D., Poole, C., Isola, J., Earl, H. and Mouridsen, H., 2011. HER2 and TOP2A as predictive markers for anthracycline-containing chemotherapy regimens as adjuvant treatment of breast cancer: a meta-analysis of individual patient data. *The lancet oncology*, 12(12), pp. 1134-1142.

Rank	ID	p-value	t	B	FC	Gene
220	3756193	6.64E-13	9.52	19.196192	1.5 ± 0.3	topoisomerase (DNA) II alpha

Table 1: TOP2A is differentially expressed in human breast cancer.

Rank of differential expression, probe ID, p-value of global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the groups compared, fold change of TOP2A in primary breast tumors as compared to normal breast, gene and gene name are listed in this chart.

Rank	ID	p-value	t	B	FC	Gene	Gene name
127	201291_s_at	4.69E-16	9.222495	25.99946	1.728 ± 0.242	TOP2A	topoisomerase (DNA) II alpha

Table 2: TOP2A is differentially expressed in human breast cancer.

Rank of differential expression, probe ID, p-value of global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the groups compared, fold change of TOP2A in primary breast tumors as compared to normal breast, gene and gene name are listed in this chart.

Figure 1: TOP2A is expressed at significantly higher levels in primary breast tumors as compared to normal breast tissue.

The mRNA expression level of TOP2A in normal breast tissue (left) and in primary tumors of the breast (right) is graphically depicted with the result of a statistical test evaluating significance of difference in TOP2A expression between normal breast tissue and primary tumors of the breast, a p -value, listed above.

Figure 2: TOP2A is expressed at significantly higher levels in primary breast tumors as compared to normal breast tissue.

The mRNA expression level of TOP2A in normal breast tissue (left) and in primary tumors of the breast (right) is graphically depicted with the result of a statistical test evaluating significance of difference in TOP2A expression between normal breast tissue and primary tumors of the breast, a *p*-value, listed above.

Figure 3: TOP2A expression correlates with overall survival in patients with breast cancer.

Depicted in this Kaplan-Meier plot is the probability of overall survival for $n=1402$ total patients stratified into two groups, based on low or high expression of TOP2A in patient primary tumors. The log rank p -value denoting statistical significance of difference in overall survival when comparing the two groups, as well as hazard ratio for this comparison is listed above. Listed below is the number of patients at risk (number of patients alive) per interval, after stratification based on TOP2A expression; in the first interval, number at risk is number of patients alive; in each subsequent interval, number at risk is the number at risk less those who have expired or are censored.

Figure 4: TOP2A expression correlates with relapse-free survival in patients with breast cancer.

Depicted in this Kaplan-Meier plot is the probability of relapse-free survival (RFS) for $n=3951$ total patients stratified into two groups, based on low or high expression of TOP2A in patient primary tumors. The log rank p -value denoting statistical significance of difference in relapse-free survival when comparing the two groups, as well as hazard ratio for this comparison is listed above. Listed below is the number of patients at risk (number of patients alive) per interval, after stratification based on TOP2A expression; in the first interval, number at risk is number of patients alive; in each subsequent interval, number at risk is the number at risk less those who have expired or are censored.

Low expression cohort (months)	High expression cohort (months)
169.2	67.4

Table 3: Median overall survival is superior in breast cancer patients with low primary tumor expression of TOP2A.

The overall survival of $n=1402$ breast cancer patients based on stratification into low or high tumor expression of TOP2A is listed in this chart.

Low expression cohort (months)	High expression cohort (months)
216.66	163.46

Table 4: Median relapse-free survival is superior in breast cancer patients with low primary tumor expression of TOP2A.

The relapse-free survival of $n=3951$ breast cancer patients based on stratification into low or high tumor expression of TOP2A is listed in this chart.