**2009 Southwestern University Research Congress
Podium Presentation**

Nurses' Performance and Clients' Delight

Ma. Fe Ivy H. Ortiz
Mauro Allan P. Amparado

Abstract

Objectives:

This study determined the nurses' performance and the clients' delight of Ward 8 of the Vicente Sotto Memorial Medical Center, Cebu City, Cebu, Philippines. The findings of the study were the basis for a proposed Client Service Seminar of the Nursing Service.

Specifically, the study answered the following queries:

1. What is the profile of the patients in terms of:
 - 1.1. age;
 - 1.2. gender;
 - 1.3. civil status;
 - 1.4. highest educational attainment;
 - 1.5. religion; and
 - 1.6. monthly income?

2. What is the profile of the staff nurses in terms of:
 - 2.1. age;
 - 2.2. gender;
 - 2.3. civil status;
 - 2.4. highest educational attainment;
 - 2.5. religion;
 - 2.6. client service seminars attended; and

2.7. years of experience?

3. What is the performance of the staff nurses in terms of:

3.1. assessment;

3.2. planning;

3.3. implementation;

3.4. evaluation;

3.5. documentation; and

3.6. work behavior/professional values?

4. What is the level of clients' delight in terms of:

4.1. assurance;

4.2. responsiveness;

4.3. empathy;

4.4. reliability;

4.5. tangibility?

5. Is there a significant relationship between the profile and performance of staff nurses?

6. Is there a significant relationship between the profile and clients' delight?

7. What client service seminar can be proposed based on the findings of the study?

Methods:

The descriptive-correlational design was utilized as research design. The study included 90 patients who had at least 2 weeks stay in the hospital, with ages between 18-50 years old, literate and conscious. The study also includes 9 staff nurses. The study was conducted at Vicente Sotto Memorial Medical Center Ward 8, Cebu

City, Cebu, Philippines. The utilized a modified tool of SERVQUAL which measures service quality.

Findings and Conclusion:

Majority of the patients are male, married, high school graduates, 31-40 years old, and has a monthly income below P5,000.00. They were highly delighted in terms of assurance, reliability, and tangibility. However, they were moderately delighted in terms of responsiveness and empathy.

Majority of the staff nurses are 31-40 years old, married, with 1-4 years of work service, with units in a Master's degree program, and has attended in at least 2 clients' delight seminar in a year. The staff nurses were very satisfactory in terms of performance.

Only two pair of variables have significant relationships: civil status and empathy; and income and responsiveness.

Recommended citation:

Ortiz, M. F. I. H. & Amparado, M. A. P. (2009, March).

Nurses' Performance and Clients' Delight.

Paper presented in the 2009 Southwestern University Research Congress, Southwestern University, Cebu City, Cebu, Philippines, 1(1), 20-22.

**2009 Southwestern University Research Congress
Podium Presentation**

School Health Management Program of a University

Neil S. Lacanlale
Mauro Allan P. Amparado

Abstract

Objectives:

The study evaluated the implementation of the school health management program of a university in Cebu City, Cebu, Philippines. The findings were the bases for an improved school health management manual.

Specifically, the study answered the following queries:

1. What is the level of implementation of the medical health program in the aspects of:
 - 1.1. health and nutrition education;
 - 1.2. health school living;
 - 1.3. health and nutrition services;
 - 1.4. school-community coordination for health and nutrition?
2. What is the level of implementation of the dental health program?
3. What is the level of implementation of the nursing health program?
4. What is the level of implementation of the nutrition program?

5. Is there a significant difference in the extent of implementation of the school health program as perceived by students in the aspects of:
 - 5.1. medical health program;
 - 5.2. dental health program;
 - 5.3. nutrition program;
 - 5.4. nursing health program;
6. What improved school health management manual can be proposed based on the findings of the study?

Methods:

This study utilized the descriptive-evaluative design. The study utilized 295 respondents with three groups: basic education students, high school students, and college students. It was conducted at the University of Southern Philippines Foundation, Cebu City, Cebu, Philippines.

Findings and Conclusion:

The basic education students perceived that there was fair implementation of the medical health, dental health, nutrition, and nursing health programs. The high school students viewed the four programs as moderately implemented. The college students viewed the four programs as fairly implemented. There were significant differences in the four programs as perceived by the three groups.

These health programs must be intensified and strengthened. Services across all programs should be delivered extensively.

Recommended citation:

Lacanalale, N. S. & Amparado, M. A. P. (2009, March).

School Health Management Program of a University.
Paper presented in the 2009 Southwestern University
Research Congress, Southwestern University, Cebu City,
Cebu, Philippines, 1(1), 23-25.

**2009 Southwestern University Research Congress
Podium Presentation**

**Awareness of the Multi-para Mothers on
Family Planning Methods**

Jessica Athena S. Villaflor
Mauro Allan P. Amparado

Abstract

Objectives:

The study determined the awareness of the multi-para mothers on natural and artificial methods of family planning at Village Basak San Nicolas, Cebu City, Cebu, Philippines. The findings of the study served as basis for a proposed enhanced family planning education manual.

Specifically, the study answered the following queries:

1. What is the profile of the multi-para mothers as to:
 - 1.1. age;
 - 1.2. educational attainment;
 - 1.3. number of children; and
 - 1.4. monthly income?
2. What is the level of awareness of the multi-para mothers on natural family planning methods in the aspects of:
 - 2.1. Lactational Amenorrhea method;
 - 2.2. cervical mucus;
 - 2.3. coitus interruptus;
 - 2.4. abstinence; and
 - 2.5. rhythm method?

3. What is the level of awareness of the multi-para mothers on artificial family planning methods in the aspects of:
 - 3.1. intrauterine device;
 - 3.2. condom;
 - 3.3. bilateral tubal ligation;
 - 3.4. pills; and
 - 3.5. Depo-medroxyprogesterone Acetate?
4. Is there a significant relationship between the level of awareness on natural family planning methods and their:
 - 4.1. age;
 - 4.2. educational attainment;
 - 4.3. number of children; and
 - 4.4. monthly income?
5. Is there a significant relationship between the level of awareness on artificial family planning methods and their:
 - 5.1. age;
 - 5.2. educational attainment;
 - 5.3. number of children; and
 - 5.4. monthly income?
6. What enhanced family planning education manual can be proposed based on the findings of the study?
7. What client service seminar can be proposed based on the findings of the study?

Methods:

The descriptive-correlational design was utilized as research design. The study was conducted in Village Basak San Nicolas, Cebu City, Cebu, Philippines. The respondents of the study were 371 multi-para mothers with two or more children. The study used

a researcher-made instrument. Treatment of data used were simple percentage, weighted mean, and chi-square Test of Independence.

Findings and Conclusion:

Majority of the multi-para mothers belong to the 30-49 years old, high school graduates, with 2-4 children, and monthly income below 10,000 Philippine pesos. They were less aware on Lactational Amenorrhea method, Cervical Mucus, Coitus Interruptus, abstinence, and Rhythm method. They were moderately aware on IUD, condom, bilateral tubal ligation, pills, and Depo-medroxyprogesterone acetate. In general, they were less aware on natural family planning methods and moderately aware on artificial family planning methods.

There were significant relationships between the level of awareness on the natural family planning methods and: age, educational attainment, number of children, and monthly income. In addition, there were significant relationships between the level of awareness on artificial planning methods and: educational attainment, number of children, and monthly income.

Recommended citation:

Villaflor, J. A. S. & Amparado, M. A. P. (2009, March).

Awareness of the Multi-para Mothers on Family Planning Methods. Paper presented in the 2009 Southwestern University Research Congress, Southwestern University, Cebu City, Cebu, Philippines, 1(1), 26-28.

**2009 Southwestern University Research Congress
Podium Presentation**

Work Orientation and Motivation of Staff Nurses

Julie Ann Y. Muaña
Mauro Allan P. Amparado

Abstract

Objectives:

This study aimed to determine and evaluate the work orientation and motivation of staff nurses of Sacred Heart Hospital, Cebu City, Cebu, Philippines. The findings of the study served as basis for a proposed staff nurse manual.

Specifically, the study answered the following queries:

1. What is the profile of the staff nurses in terms of:
 - 1.1. age;
 - 1.2. gender;
 - 1.3. area of assignment;
 - 1.4. civil status;
 - 1.5. highest educational attainment;
 - 1.6. length of service?
2. What is the level of work motivation of staff nurses?
3. What is the level of motivation of the staff nurses in terms of:
 - 3.1. existence;
 - 3.2. relatedness; and
 - 3.3. growth?
4. Is there a significant relationship between:

- 4.1. profile and work orientation;
- 4.2. profile and motivation; and
- 4.3. work orientation and motivation?

5. What can be the proposed staff nurse manual based on the findings of the study?

Methods:

The study utilized the descriptive-correlational design. The research locale was Sacred Heart Hospital, Cebu City, Cebu, Philippines. There were 51 staff nurses who served as respondents of the study. The researchers created a questionnaire with three parts: profile, work orientation, and motivation. Statistical treatment used were simple percentage, weighted mean, chi-square Test of Independence, and Pearson-r.

Findings and Conclusion:

Majority of the respondents were 21-23 years old, female, single, has served the hospital for less than two years, BSN graduates, and assigned in the Operating and Delivery Rooms. The level of work motivation of the respondents was low. The level of motivation was also low. The existence needs were moderate, whereas the relatedness needs and growth needs were low.

Moreover, there was no significant relationship between profile and work orientation. It also revealed that there was no significant relationship between profile and motivation. Finally, there was no significant relationship between work orientation and motivation.

Recommended citation:

Muaña, J. A. Y. & Amparado, M. A. P. (2009, March).

Work Orientation and Motivation of Staff Nurses. Paper presented in the 2009 Southwestern University Research Congress, Southwestern University, Cebu City, Cebu, Philippines, 1(1), 29-31.

**2009 Southwestern University Research Congress
Podium Presentation**

**Instructional Innovation and Communication
Skills of Clinical Instructors**

Antonette C. Ramos
Mauro Allan P. Amparado

Abstract

Objectives:

This study determined the instructional innovation and communication skills of Clinical Instructors of the University of Cebu Lapu-Lapu and Mandaue, Mandaue City, Cebu, Philippines. The findings served as basis for a proposed instructional innovation and communication skills seminar.

Specifically, it sought to answer the following queries:

1. What is the level of instructional innovation of the respondents in terms of:
 - 1.1. direct instruction;
 - 1.2. concept teaching;
 - 1.3. cooperative learning;
 - 1.4. problem-based instruction.

2. What is the level of communication skills of the respondents in terms of:
 - 2.1. verbal communication;
 - 2.2. non-verbal communication?

3. Is there a significant relationship between the instructional innovation and verbal communication skills of the respondents?
4. Is there a significant relationship between the instructional innovation and non-verbal communication skills of the respondents?
5. What instructional innovation and communication skills seminars can be proposed based on the findings of the study?

Methods:

The study utilized the descriptive-correlational design. It was conducted in the University of Cebu Lapu-Lapu and Mandaue, Mandaue City, Cebu, Philippines. There were 197 nursing students who evaluated the Clinical Instructors of the school. Stratified sampling was used. The researchers utilized a modified structured instrument on innovation by Arends and communication skills instrument by Hamm. Statistical treatment used were simple percentage, weighted mean, and chi-square Test of Independence.

Findings and Conclusion:

The nursing students perceived the Clinical Instructors to be very good in direct instruction, concept, teaching, cooperation learning, problem-based instruction, verbal communication, and non-verbal communication. There was a significant relationship between instructional innovation and verbal communication. Furthermore, there was a significant relationship between instructional innovation and non-verbal communication.

Recommended citation:

Ramos, A. C. & Amparado, M. A. P. (2009, March).
Instructional Innovation and Communication Skills of Clinical
Instructors. Paper presented in the 2009 Southwestern
University Research Congress, Southwestern University,
Cebu City, Cebu, Philippines, 1(1), 32-34.

**2009 Southwestern University Research Congress
Podium Presentation**

**Exodus of Nurses in a Tertiary Government Hospital:
Nursing Service Workforce Perspectives**

Ma. Gay M. Gonzales
Mauro Allan P. Amparado

Abstract

Objectives:

The purpose of the study was to determine the perspectives of the Nursing service workforce on the exodus of nurses occurring at a government tertiary hospital and their interventions. The outcome of the study served as basis for recommendations to the Human Resource Management of the hospital.

Specifically, the study answered the following questions:

1. What are the perspectives of the nurse administrators on the exodus of nurses in terms of:
 - 1.1. staff replacement;
 - 1.2. scheduling of duties;
 - 1.3. work assignment;
 - 1.4. scheduling of leaves;
 - 1.5. supervision;
 - 1.6. staff promotion; and
 - 1.7. budget needs.
2. What are the perspectives of the staff nurses on the exodus of nurses in terms of:
 - 2.1. work load;
 - 2.2. competence;

- 2.3. promotion and career path; and
 - 2.4. nursing administrators' functions.
3. What are the perspectives of the nursing attendants on the exodus of nurses in terms of:
- 3.1. own workload;
 - 3.2. nurses' work load;
 - 3.3. nurses' competence;
 - 3.4. nurses' promotion and career path;
 - 3.5. its affectations on nursing administrators' functions?
4. What interventions have been put in place to cope with the affectations of the exodus of nurses as perceived by:
- 4.1. nurse administrators;
 - 4.2. staff nurses; and
 - 4.3. nursing attendants?

Methods:

The qualitative study utilized the nursing service workforce of Zamboanga City Medical Clinic, Zamboanga City, Philippines which consists of 20 nurse supervisors, 95 staff nurses, and 41 nursing attendants. A structured interview guide was used.

Findings and Conclusion:

The nurse administrators perceived that the exodus of nurses affected their work as it took time to replace staff. Due to the lack of personnel, scheduling of duties was difficult. They acted as relievers or went "on call" or as "trouble shooters." Granting of leaves was difficult and more supervision was needed over the new hires. However, there was a greater chance of staff nurse promotion. More budget was needed for hiring of contractual health workers to augment the nursing workforce.

The staff nurses also perceived an increase in their work load. Although the senior nurses were competent, the increase in patient and area loads made them less effective, having less time for patient care. However, the staff nurses had more chances of getting promoted.

Some nursing attendants, likewise, perceived an increase in their work load because they worked with the staff nurses as a team. Others, however, claimed that they were not affected of the exodus since they have different functions and responsibilities.

There is an increase in the workload for the nursing service workload due to nurse migration while replacement process is underway. Volunteers and contractual health workers were utilized to augment the workforce. Staff promotion is a welcome event caused by the vacancies due to the resignation of employees.

The exodus of nurses adversely affected the financial status, system and operations of the hospital. However, the nursing attendants were not affected because they maximized the volunteer and contractual workers to augment the workforce.

Recommended citation:

Gonzales, M. G. M. & Amparado, M. A. P. (2009, March).

Exodus of Nurses in a Tertiary Government Hospital: Nursing Service Workforce Perspectives. Paper presented in the 2009 Southwestern University Research Congress, Southwestern University, Cebu City, Cebu, Philippines, 1(1), 35-37.

**2009 Southwestern University Research Congress
Podium Presentation**

**Classical Music as a Burnout Reduction Therapy to
Clinical Instructors**

Joanna Rae C. Semblante
Mauro Allan P. Amparado

Abstract

Objectives:

This study determined the effectiveness of music as a burnout reduction strategy to clinical instructors in Southwestern University, Cebu City, Cebu, Philippines. The findings were the basis for a proposed faculty burnout management strategy.

Specifically, the study answered the following inquiries:

1. What is the profile of the clinical instructors in terms of:
 - 1.1. age;
 - 1.2. gender;
 - 1.3. civil status;
 - 1.4. highest educational attainment;
 - 1.5. monthly income;
 - 1.6. number of years in service?
2. What is the pre-test and post-test levels of the burnout among the clinical instructors in the control and experimental groups in terms of:
 - 2.1. emotional exhaustion;
 - 2.2. depersonalization; and
 - 2.3. lack of personal accomplishment?

3. Is there a significant difference between the pre-test and post-test levels of burnout among the clinical instructors.
4. What are the perceived work-related stressors of the clinical instructors?
5. What will be the proposed faculty burnout strategy based on the findings of the study?

Methods:

This study utilized the experimental design. A stratified random sample of 20 clinical instructors in a control group and 20 clinical instructors in an experimental group completed the Maclach's Burnout Inventory (MBI) which measured emotional exhaustion, depersonalization, and lack of personal accomplishment. The MBI was administered to the respondents before, after the third week, and after the fifth week of exposing the experimental group to classical music and the control group to no music. The perceived work-related stressors of the clinical instructors were also investigated through a focus group discussion.

Findings and Conclusion:

Majority of the respondents in the experimental group and half of those in the control group are 24-29 years old. In both groups, most are females, married, and have BSN degrees with MAN units. In both groups, more than half of them have an average income of 15,000 to 20,000 Philippine pesos and has served the college for 0-5 years.

The emotional exhaustion of the experimental group was moderate and the control group was low. On depersonalization and lack of personal accomplishment, both groups were low.

The experimental group's emotional exhaustion level after exposure to classical music for three weeks was moderate, while the control group was low. On depersonalization and lack of personal accomplishment, both groups were low.

After the fifth week, the experimental group's emotional exhaustion was moderate, while the control group was low. On depersonalization and lack of personal accomplishment, both groups were low.

After exposing the experimental group to classical music for five weeks, it was statistically revealed that there was no significant difference in the pre-test and post-test burnout levels for the two groups.

The perceived work-related stressors of the clinical instructors were: conflicts with colleagues; six work days and graduate study classes; assignment in new clinical areas; proper channeling of concerns/conflicts with immediate heads; unfair distribution of loads; lecture loads given with shorter time to prepare; and being forced to stay in the facility/office for the whole eight hours.

In conclusion, the use of classical music is not an effective strategy in reducing the burnout level of clinical instructors.

Recommended citation:

Semblante, J. R. C. & Amparado, M. A. P. (2009, March).

Classical Music as a Burnout Reduction Therapy to Clinical Instructors. Paper presented in the 2009 Southwestern University Research Congress, Southwestern University, Cebu City, Cebu, Philippines, 1(1), 38-40.

**2009 Southwestern University Research Congress
Podium Presentation**

Safe and Quality Nursing Care Practices

Ma. Golda H. Castro
Mauro Allan P. Amparado

Abstract

Objectives:

This study assessed the safe and quality nursing care practices of senior nursing students of the University of San Jose-Recoletos as perceived by the nursing faculty and senior nursing students. An enhanced educational program was proposed based on the findings of the study.

Specifically, the study answered the following inquiries:

1. What is the profile of the senior nursing students in terms of:
 - 1.1. age;
 - 1.2. gender; and
 - 1.3. average RLE grade?

2. What is the profile of the clinical instructors in terms of:
 - 2.1. age;
 - 2.2. gender; and
 - 2.3. educational attainment?

3. What are the practices of senior nursing students on safe and quality nursing care as perceived by:
 - 3.1. faculty; and
 - 3.2. senior nursing students?

4. Is there a significant relationship between the profile and the level of safe and quality nursing care practices as perceived by the faculty and senior nursing students?
5. What enhanced instructional program can be proposed based on the findings of the study?

Methods:

This study utilized the descriptive-correlational design. The respondents of the study were 90 senior nursing students and 8 faculty members assigned at the medical-surgical and orthopedic wards of Cebu City Medical Center, Cebu City, Cebu, Philippines. A researcher-made questionnaire was used. Statistical treatment used were simple percentage, weighted mean, chi-square test of independence, and t-test of independent samples.

Findings and Conclusion:

Majority of the senior nursing students were 18-20 years old, female, and with very good RLE grades. They perceived that they always practice safe and quality nursing care. On the other hand, the faculty members were 24-26 years old, female, and graduates of the BSN degree with MAN/MN units. They perceived that they always practice safe and quality nursing care.

There was no significant relationship between the profile of the students and the safe and quality nursing care practices as perceived by the two groups.

Recommended citation:

Castro, M. G. H. & Amparado, M. A. P. (2009, March).

Safe and Quality Nursing Care Practices. Paper presented in the 2009 Southwestern University Research Congress, Southwestern University, Cebu City, Cebu, Philippines, 1(1), 41-43.

**2009 Southwestern University Research Congress
Podium Presentation**

**Knowledge, Attitudes and Practices of Hospital
Personnel on Newborn Screening Program**

Patrick Ray M. Gonzales
Mauro Allan P. Amparado

Abstract

Objectives:

This study determined the knowledge, attitudes and practices of hospital personnel of Zamboanga City Medical Center, Zamboanga City, Philippines on the newborn screening program (NBS) of the Department of Health. The findings of which served as the basis for the formulation of a seminar-workshop program.

Specifically, the study answered the following questions:

1. What are the respondents' level of knowledge on the NBS program of the Department of Health per department in terms of:
 - 1.1. mechanics;
 - 1.2. benefits; and
 - 1.3. implementing rules and guidelines?
2. What are the respondents' attitudes on the NBS program per department?
3. What are the respondents' practices in the implementation of the NBS program per department?

4. Is there a significant difference in the knowledge, attitudes and practices of the respondents on the NBS program per department:

4.1. between groups;

4.2. within groups?

5. What deterrents are identified by the respondents and their corresponding suggestions and recommendations for the full implementation of the NBS program?

Methods:

This study utilized the descriptive-evaluative design. The study was conducted at Zamboanga City Medical Center with 60 respondents composed of doctors, nurses and nursing attendants. The departments were Obstetrics, Pediatric, and Emergency Room departments. A 30-item researcher-modified instrument was used based on the NBS program flyer. Statistical treatment used were weighted mean and ANOVA.

Findings and Conclusion:

The Obstetrics and Pediatrics departments have higher knowledge than the Emergency Room department. The Obstetrics and Pediatrics departments find the program desirable while the Emergency Room department find the program less desirable. In terms of practices, the Pediatrics department was rated good, the Obstetrics department was fair, and the Emergency Room department was poor.

There was a significant difference in the knowledge, attitudes and practices between and within groups. The deterrents identified by the respondents were: test is expensive; there is lack of awareness on the importance of NBS among parents; low acceptability of NBS to parents; parents are less interested with

NBS program; parents live too far to avail of the tests and follow-up results; poor information campaign; NBS is not well implemented; lack of knowledge of other hospital personnel; doctor in-charge is not available; results of tests are delayed.

Recommended citation:

Gonzales, P. R. M. & Amparado, M. A. P. (2009, March).
Knowledge, Attitudes and Practices of Hospital Personnel on
Newborn Screening Program. Paper presented in the 2009
Southwestern University Research Congress, Southwestern
University, Cebu City, Cebu, Philippines, 1(1), 44-46.

**2009 Southwestern University Research Congress
Podium Presentation**

Hygiene and Motivational Factors of Clinical Instructors

Corazon B. Dumadag
Mauro Allan P. Amparado

Abstract

Objectives:

This study determined the hygiene and motivational factors of clinical instructors of Southwestern University, Cebu City, Cebu, Philippines. The findings served as basis for a proposed human resource retention plan.

Specifically, the study answered the following queries:

1. What is the profile of the respondents in terms of:
 - 1.1. age;
 - 1.2. gender;
 - 1.3. civil status;
 - 1.4. educational attainment;
 - 1.5. length of service?
2. What is the turnover rate among clinical instructors?
3. What is the level of hygiene and motivational factors of clinical instructors that retains them in the workplace?
4. Is there a significant relationship between the hygiene and motivational factors of the clinical instructors?

5. Is there a significant difference on the hygiene and motivational factors between male and female clinical instructors?

6. Based on the findings of the study, what human resource retention plan can be proposed?

Methods:

This study utilized the descriptive-correlational design. The study was conducted at the College of Nursing, Southwestern University, Cebu City, Cebu, Philippines. The respondents were 25 clinical instructors. A researcher-made questionnaire was used to collect data. Statistical treatment utilized were simple percentage, weighted mean, Pearson r , and t-test for independent samples.

Findings and Conclusion:

Majority of the respondents were 25-35 years old, female, married, finished BSN with units in a master's program, and with 3-5 years of work experience in the college. There were no resignations from clinical instructors hired from 2003 to 2007. The level of hygiene and motivational factors were high.

There was a significant relationship between the hygiene and motivational factors of the clinical instructors. There were no significant differences on the hygiene and motivational factors between male and female clinical instructors.

Recommended citation:

Dumadag, C. B. & Amparado, M. A. P. (2009, March).

Hygiene and Motivational Factors of Clinical Instructors.
Paper presented in the 2009 Southwestern University

Research Congress, Southwestern University, Cebu City,
Cebu, Philippines, 1(1), 47-49.

**2009 Southwestern University Research Congress
Podium Presentation**

**Tardiness and Absenteeism of Clinical Instructors:
Proposed Intervention Plan**

Amalia L. Martinez
Mauro Allan P. Amparado

Abstract

Objectives:

This study determined the perceptions of clinical instructors on tardiness and absenteeism in the College of Nursing, Southwestern University, Cebu City, Cebu, Philippines. The findings of the study served as basis for an intervention plan.

Specifically, the study answered the following queries:

1. What is the profile of the respondents in terms of:
 - 1.1. age;
 - 1.2. gender; and
 - 1.3. civil status?
2. What is the frequency of tardiness and absenteeism among clinical instructors?
3. What is the extent of the factors that contributes to tardiness and absenteeism?
4. Is there a significant relationship between the respondents' profile and the extent of the factors that contributes to tardiness and absenteeism?

5. Based on the findings of the study, what intervention plan can be proposed?

Methods:

This study utilized the descriptive-correlational design. The study was conducted at the College of Nursing, Southwestern University, Cebu City, Cebu, Philippines. The respondents were 50 clinical instructors. A researcher-made questionnaire was used to collect data. Statistical treatment utilized were simple percentage, linear regression, and chi-square test of independence.

Findings and Conclusion:

Majority of the respondents were 22-34 years old, female, and married. The trend of absenteeism was highest in July 2007 but the tardiness was declining from October 2006 to November 2006. Extent of factors that contributed to absenteeism were low for personal, family, work, and environmental aspects. It was high for health aspects. Extent of factors that contributed to tardiness were low for personal, family, health, work, and environmental aspects. There were no significant relationships between the respondents' profile and the extent of the factors that contributes to absenteeism and tardiness.

Recommended citation:

Martinez, A. L. & Amparado, M. A. P. (2009, March).

Tardiness and Absenteeism of Clinical Instructors: Proposed Intervention Plan. Paper presented in the 2009 Southwestern University Research Congress, Southwestern University, Cebu City, Cebu, Philippines, 1(1), 50-51.

**2009 Southwestern University Research Congress
Podium Presentation**

**Exchange Leadership Styles and Core
Values of Nurse Leaders**

Constantina C. Chan
Mauro Allan P. Amparado

Abstract

Objectives:

This study determined the exchange leadership styles and core values of the nurse leaders of the College of Nursing of Southwestern University, Cebu City, Cebu, Philippines. The findings of the study served as basis for a proposed action plan.

Specifically, the study answered the following queries:

1. What is the level of transformational and transactional leadership styles of the nurse leaders in the pre-test and post-test with reference to the following management functions:
 - 1.1. planning;
 - 1.2. organizing;
 - 1.3. directing; and
 - 1.4. controlling?
2. What are the core values identified by the respondents in the pre-test and post-test?
3. Is there a significant difference between the pre-test and post-test results in terms of the following:
 - 3.1. transformational and transactional leadership styles;
 - 3.2. core values?

4. Based on the findings of the study, what action plan may be proposed?

Methods:

This study utilized the descriptive-comparative design. The study was conducted at Southwestern University College of Nursing with 12 subjects. The group attended the leadership seminar on Redefining Core Values in Leadership by a prominent motivational speaker. Statistical treatment used were simple percentage, weighted mean, and paired t-test.

Findings and Conclusion:

The subjects were very satisfactory in the post-test of transformational and transactional leadership styles. It was noted that the respondents improved in organizing and controlling aspects of transformational leadership style. They also improved in the planning and directing aspects of the transactional leadership style. Among the core values, they were very satisfactory in the aspects of commitment, compassion and cooperation. They were satisfactory in the aspect of competence.

There were no significant differences between the pre-test and post-test results in terms of transformational leadership style, transactional leadership style, and core values.

Recommended citation:

Chan, C. C. & Amparado, M. A. P. (2009, March).
Exchange Leadership Styles and Core Values of Nurse Leaders. Paper presented in the 2009 Southwestern University Research Congress, Southwestern University, Cebu City, Cebu, Philippines, 1(1), 52-53.