TCF4 is differentially expressed in the metastases of patients with breast cancer.

Shahan Mamoor, MS¹

¹Thomas Jefferson School of Law

San Diego, CA 92101

mamoorsk@tjssl.edu

Metastasis is a major problem in patients with breast cancer (1). We analyzed published microarray datasets (2, 3) to identify transcription factors whose expression changed most significantly between primary tumors in patients with breast cancer and metastatic tissues. We identified TCF4 as differentially expressed when comparing the transcriptomes of primary tumors of the breast and metastases to the liver (2). Analysis of a separate microarray dataset uncovered similar differential expression of TCF4 in metastases to the brain compared to primary breast tumors (3). TCF4 is transcriptionally perturbed in metastases to brain and the liver in patients with breast cancer.

Keywords: breast cancer, metastasis, metastatic breast cancer, TCF4, systems biology of breast cancer, tumor transcriptome analysis.

Patients diagnosed with metastatic breast cancer are faced with a 5-year overall survival rate of 27% (10). An enhanced understanding of the metastatic process in patients with breast cancer is required in order to design novel therapeutics to halt, reverse or delay metastases in patients with breast cancer. Systems-level analysis of the metastatic transcriptome as compared to primary tumors of the breast can facilitate this understanding by revealing how gene expression is most significantly different in metastatic tissues and help identify therapeutic targets for patients with metastatic breast cancer. In this study we performed such an analysis, mining published transcriptome data to compare global transcriptional profiles of liver and brain metastases from patients with breast cancer to primary tumors of the breast. Across both datasets, we identified the transcription factor TCF4 as among the most differentially expressed genes in metastases when compared to primary tumors of the breast. In lung adenocarcinoma, enrichment for a TCF4 gene signature in patient tumors correlates with metastasis-free survival and recurrence of the disease (4); however, we found that TCF4 was expressed at significantly lower levels in the metastatic tissues of patients with breast cancer as compared to primary tumors of the breast.

Methods

We used GSE124648 (2) and GSE125989 (3) for this differential gene expression analysis in conjunction with GEO2R. GSE124648 was generated using Affymetrix Human Genome U133A array technology with $n=235$ primary tumors of the breast and $n=16$ liver metastases. These primary tumors included stage IA, IIA, IIB, IIIA and IIIC or of a stage not indicated. Tissue sample types included surgical tissue, fine needle aspiration, or core biopsy. GSE125989 was generated using Affymetrix Human Genome U133A 2.0 Array technology with $n=16$ primary tumors of the breast and $n=16$ brain metastases.

The Benjamini and Hochberg method of p-value adjustment was used for ranking of differential expression but raw p-values were used for assessment of statistical significance of

global differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used.

To evaluate the statistical significance of difference in TCF4 expression between primary and metastatic tumor tissues, we used a two-tailed, unpaired t-test with Welch's correction, using PRISM (8.4.0)(455).

Results

We sought to further understand the transcriptional behavior of breast cancer metastases using human data at the systems-level in an unbiased fashion. Thus, we mined published transcriptome data (2, 3) to identify transcription factors whose expression was most significantly different between primary tumors of the breast and metastases in patients with breast cancer.

TCF4 is differentially expressed in metastases to the liver when compared to primary tumors of the breast.

We identified the transcription factor TCF4 (transcription factor four) as among the genes most differentially expressed in metastases to the liver from patients with breast cancer when compared to primary tumors of the breast. When sorting all of the mRNAs detected by microarray based on change in expression between liver metastases and primary tumors of the breast, TCF4 ranked 240 out of 22283. Differential expression of TCF4 in liver metastases when compared to primary tumors from patients with breast cancer was statistically significant (Table 1; $p=1.94E-18$).

TCF4 is differentially expressed in metastases to brain when compared to primary tumors of the breast.

Analysis of a separate microarray dataset, this containing transcriptome information from the primary tumors and brain metastases of patients with breast cancer revealed that TCF4 was

also among the genes most differentially expressed in metastases to brain when compared to primary tumors of the breast. When sorting all of the mRNAs detected by microarray based on change in expression between brain metastases and primary tumors of the breast, TCF4 ranked 59 out of 22277. Differential expression of TCF4 in metastases to brain when compared to primary tumors from patients with breast cancer was statistically significant (Table 2; $p=4.89E-05$).

TCF4 is expressed at significantly lower levels in metastases to the liver when compared to primary tumors of the breast.

We extracted exact mRNA expression levels of TCF4 from each liver metastasis sample and from each primary breast tumor sample to compare expression of TCF4 between metastases and primary tumors. We also performed a statistical test to determine if the difference in TCF4 expression between primary and liver metastases was statistically significant. TCF4 expression was significantly lower in liver metastases when compared to primary tumors of the breast in patients with breast cancer (Figure 1; $p<0.0001$). We calculated a fold change of 0.69 in expression of TCF4 between liver metastases and primary tumors (Table 1).

TCF4 is expressed at significantly lower levels in metastases to the brain when compared to primary tumors of the breast.

Again, we extracted exact mRNA expression levels of TCF4 from each metastasis sample and from each primary breast tumor sample to compare expression of TCF4 between brain metastases and primary tumors. We also performed a statistical test to determine if the difference in TCF4 expression between primary and brain metastases was statistically significant. TCF4 expression was significantly lower in metastases to brain in patients with breast cancer when compared to primary tumors of the breast (Figure 2; $p<0.0001$). We calculated a fold change of 0.87 in expression of TCF4 between brain metastases and primary tumors (Table 2).

Thus, we found that TCF4 was differentially expressed in metastases to the liver and brain when compared to primary tumors from patients with breast cancer, and that it was expressed at significantly lower levels in metastases when compared to primary tumors of the breast.

Discussion

In this study we mined published transcriptome data (2, 3) from the metastatic tissues and primary tumors of patients with breast cancer to identify transcription factors differentially expressed between metastases and primary tumors. We identified TCF4 as among the genes whose expression most significantly changed between metastases to the brain and to the liver as compared to primary breast tumors. TCF4 was expressed at significantly lower levels in metastases at both organ sites when compared to primary tumors from patients with breast cancer.

TCF4 expression negatively correlates with overall patient survival in colorectal cancer (6). TCF4 promotes invasiveness in colorectal cancer through induction of MT1-MMP and LAMC2, enabling cleavage of MT1-MMP by LAMC2 (6). TCF4 binds to the promoter of the transcriptional repressor ZEB1 leading to induction of the epithelial to mesenchymal transition (7). In lung adenocarcinoma, enrichment for a TCF4 gene signature in patient tumors correlated with metastasis-free survival and recurrence of the disease (4). Wnt pathway signaling involving the transcription factor TCF4 leads to transcriptional activation of the osteopontin gene (8), the gene product of which is involved in tumor progression and metastasis; this pathway is active in breast cancer (9). TCF4 also activates the expression of chemokines monocyte chemoattractant protein MCP-1/CCL2 in human breast cancer cells (10). Decreased expression of TCF4 in tumors of the breast relative to non-affected tissues has been demonstrated (11). A study of breast cancer patient-derived xenografts with five cycles of four different types of chemotherapies showed that TCF4 mutation was found *de novo* in all cases; loss of TCF4 supported chemoresistance in breast cancer cell lines (12). miR-591 was reported to be down-

regulated in breast cancer cells, and TCF4 was found to be a miR-591 pathway target, demonstrating miR-591 inhibition of TCF4 and by analogy, TCF4 de-repression by miR-591 down-regulation in breast cancer (13). Cell proliferation *in vitro* and *in vivo* was inhibited by expression of a miR-591 mimic, suggesting that TCF4 expression, enabled by miR-591 down-regulation, promoted cell proliferation in breast cancer (13). Together these data paint a portrait of TCF4 as a promoter of cell proliferation, invasive behavior, and metastasis in cancer generally, as well as specifically in breast cancer. However, we found significantly decreased expression of TCF4 in metastases to both the brain and the liver in patients with breast cancer.

We found that the transcription factor TCF4 was among the genes whose expression changed most significantly when comparing the transcriptomes of primary tumors and metastases in humans with breast cancer. TCF4 expression should be assessed in larger and separate cohorts of patients with breast cancer, in primary tumors and in metastatic tissues to determine if the findings here can be validated. We reported that ZEB1 is also among the genes most differentially expressed in metastases to the liver and to multiple organ sites in patients with breast cancer (14). ZEB1 and TCF4 may function together to promote the epithelial to mesenchymal transition in breast cancer metastasis (7). Transcriptional regulation by transcription factors in breast cancer should be studied as a system with numerous activating and repressive inputs at the transcriptional level producing a single organismal phenotype - metastasis.

References

1. Weigelt, B., Peterse, J.L. and Van't Veer, L.J., 2005. Breast cancer metastasis: markers and models. *Nature reviews cancer*, 5(8), pp.591-60
2. Cejalvo, J.M., de Dueñas, E.M., Galván, P., García-Recio, S., Gasión, O.B., Paré, L., Antolin, S., Martinello, R., Blancas, I., Adamo, B. and Guerrero-Zotano, Á., 2017. Intrinsic subtypes and gene expression profiles in primary and metastatic breast cancer. *Cancer research*, 77(9), pp.2213-2221.
3. Sinn, B.V., Fu, C., Lau, R., Litton, J., Tsai, T.H., Murthy, R., Tam, A., Andreopoulou, E., Gong, Y., Murthy, R. and Gould, R., 2019. SET ER/PR: a robust 18-gene predictor for sensitivity to endocrine therapy for metastatic breast cancer. *NPJ breast cancer*, 5(1), pp.1-8.
4. Nguyen, D.X., Chiang, A.C., Zhang, X.H.F., Kim, J.Y., Kris, M.G., Ladanyi, M., Gerald, W.L. and Massagué, J., 2009. WNT/TCF signaling through LEF1 and HOXB9 mediates lung adenocarcinoma metastasis. *Cell*, 138(1), pp.51-62.
5. ACS Cancer Facts & Figures 2019, <https://www.cancer.net/cancer-types/breast-cancer-metastatic/statistics>.
6. Kriegl, L., Horst, D., Reiche, J.A., Engel, J., Kirchner, T. and Jung, A., 2010. LEF-1 and TCF4 expression correlate inversely with survival in colorectal cancer. *Journal of translational medicine*, 8(1), p.123.
7. Sánchez-Tilló, E., de Barrios, O., Siles, L., Cuatrecasas, M., Castells, A. and Postigo, A., 2011. β -catenin/TCF4 complex induces the epithelial-to-mesenchymal transition (EMT)-activator ZEB1 to regulate tumor invasiveness. *Proceedings of the National Academy of Sciences*, 108(48), pp.19204-19209.
8. El-Tanani, M., Barraclough, R., Wilkinson, M.C. and Rudland, P.S., 2001. Metastasis-inducing DNA regulates the expression of the osteopontin gene by binding the transcription factor Tcf-4. *Cancer research*, 61(14), pp.5619-5629.
9. Ravindranath, A., Yuen, H.F., Chan, K.K., Grills, C., Fennell, D.A., Lappin, T.R. and El-Tanani, M., 2011. Wnt- β -catenin-Tcf-4 signalling-modulated invasiveness is dependent on osteopontin expression in breast cancer. *British journal of cancer*, 105(4), pp.542-551.
10. Mestdagt, M., Polette, M., Buttice, G., Noël, A., Ueda, A., Foidart, J.M. and Gilles, C., 2006. Transactivation of MCP-1/CCL2 by β -catenin/TCF-4 in human breast cancer cells. *International journal of cancer*, 118(1), pp.35-42.
11. Shulewitz, M., Soloviev, I., Wu, T., Koeppen, H., Polakis, P. and Sakanaka, C., 2006. Repressor roles for TCF-4 and Sfrp1 in Wnt signaling in breast cancer. *Oncogene*, 25(31), pp.4361-4369.

12. de Garibay Ruiz, G., Mateo, F., Stradella, A., Valdes-Mas, R., Palomero, L., Serra-Musach, J., Puente, D.A., Díaz-Navarro, A., Vargas-Parra, G., Tornero, E. and Morilla, I., 2018. Tumor xenograft modeling identifies an association between TCF4 loss and breast cancer chemoresistance. *Disease models & mechanisms*, 11(5).
13. Huang, X., Tang, F., Weng, Z., Zhou, M. and Zhang, Q., 2019. MiR-591 functions as tumor suppressor in breast cancer by targeting TCF4 and inhibits Hippo-YAP/TAZ signaling pathway. *Cancer cell international*, 19(1), p.108.
14. Mamoor, S. (2020) The zinc finger E-box binding homeobox transcriptional repressor ZEB1 is differentially expressed in the metastases of patients with breast cancer. OSF Preprints. doi: 10.31219/osf.io/96x74.

Rank	ID	p-value	t	B	Fold change	Gene	Gene name
240	212386_at	1.94E-18	-10.668012	31.412943	0.69	TCF4	transcription factor 4

Table 1: TCF4 is differentially expressed in metastases to the liver when compared to primary tumors of the breast.

The rank of differential expression, probe/transcript ID, the p-value of global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups, the fold change in mRNA expression between the metastases and primary tumors, the gene and gene name are listed in this chart.

Rank	ID	p-value	t	B	Fold change	Gene	Gene name
59	212382_at	4.89E-05	-4.5841812	1.9345	0.87	TCF4	transcription factor 4

Table 2: TCF4 is differentially expressed in metastases to brain when compared to primary tumors of the breast.

The rank of differential expression, probe/transcript ID, the p-value of global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups, the fold change in mRNA expression between the metastases and primary tumors, the gene and gene name are listed in this chart.

Figure 2: TCF4 is expressed at significantly lower levels in metastases to brain when compared to primary tumors of the breast.

The mRNA expression level of TCF4 is shown in primary tumors of the breast (left) and liver metastases (right). The mean mRNA expression levels in each group are marked and the result of a statistical test evaluating the significance of difference in expression of TCF4 between groups, a p-value, is listed.