1 **Title**

2 Longitudinal increases in childhood depression during the COVID-19 lockdown in a UK
3 cohort

4 **Authors**

5 Giacomo Bignardi¹, Edwin S. Dalmaijer¹, Alexander L. Anwyl-Irvine¹, Tess A. Smith¹, Roma
6 Siugzdaite¹, Stepheni Uh¹ & Duncan E. Astle¹

7 ¹MRC Cognition and Brain Sciences Unit, University of Cambridge

8 **Correspondence**

9 Duncan Astle, MRC Cognition and Brain Sciences Unit, University of Cambridge, 15
10 Chaucer Road, CB2 7EF, Duncan.Astle@mrc-cbu.cam.ac.uk

11

12 **Abstract**

13 **Background**

14 There has been widespread concern that so-called ‘lockdown’ measures, including social
15 distancing and school closures, could negatively impact children’s mental health. However,
16 there has been little direct evidence of this impact due to the paucity of longitudinal studies
17 reporting mental health before and during lockdown. This present study provides the first
18 longitudinal examination of changes in childhood mental health, a key component of an
19 urgently needed evidence base that can inform policy and practice surrounding the continuing
20 response to the COVID-19 pandemic.

21 **Methods**

22 Mental health assessments on 168 children (aged 7.5-11.6 years) were taken before and
23 during the UK lockdown. Assessments included self-reports, parent-reports, and teacher-
24 reports. Mean mental health scores before and during the UK lockdown were compared using
25 mixed linear models.

26 **Findings**

27 A significant increase in depression symptoms during the UK lockdown was observed, as
28 measured by the RCADS short form. Confidence intervals suggest a medium-to-large effect
29 size. There were no significant changes in RCADS anxiety subscale and SDQ emotional
30 problems subscale.

31 **Interpretation**

During the UK lockdown children's depression symptoms have increased substantially, relative to before lockdown. The scale of this effect has direct relevance for the continuation of different elements of lockdown policy, such as complete or partial school closures. This early evidence for the direct impact of lockdown must now be combined with larger-scale epidemiological studies that establish which children are most at risk, and tracks their future recovery.

Funding

This study was supported by grant TWCF0159 from the Templeton World Charity Foundation to DEA, and by the UK Medical Research Council. SU was supported by the Gates Cambridge Trust.

Research in context

Evidence before this study

Many countries have enacted policies designed to reduce physical interpersonal interaction, including social distancing and school closures, in response to the COVID-19 pandemic. We searched scopus, PubMed and the OSF preprint archive search on July 21, 2020 using the terms ("COVID-19" OR "CORONAVIRUS") AND ("child*" OR "pediatric" OR "paediatric") AND ("mental health" OR "wellbeing" OR "depression" OR "anxiety"). No additional limits were set. The search yielded 700 records. None of these records report longitudinal studies of mental health in children, with pre-lockdown assessments. The majority of studies in children were cross-sectional during lockdown, sometimes asking parents whether their children's symptoms have worsened during the lockdown, or examining changes in mental health reports throughout the lockdown period. One study in Norway compared mean well-being ratings in adolescents from large-scale, national surveys before and during the lockdown. Several longitudinal surveys in adult samples were found, and studies in adults comparing mental health in individuals who have been quarantine to those who have not.

Added value of this study

This is the first prospective longitudinal study tracking children's mental health from before to during the national COVID-19 "lockdown" in the United Kingdom. We looked at changes in three widely used mental health questionnaire scales, assessing general emotional problems, anxiety and depression. Depression ratings increased substantially during the lockdown compared to baseline, even when controlling for age, gender and socio-economic status. There were no significant changes in emotional problem and anxiety scales.

65 Implications of all the available evidence

66 Upon a backdrop of worsening mental health in children, the government response to the
67 Covid-19 pandemic will very likely have exacerbated this. The current longitudinal data
68 highlights a substantial increase in ratings of depression, and a key implication is that a
69 prompt return to school for all children must be prioritised by policymakers. When children
70 return to school, their wellbeing and socialisation should be prioritised over and above
71 scholastic progress. And in preparation for school resumption, teachers should receive
72 additional guidance on supporting young children experiencing low mood, and highlighting
73 referral pathways for specialist support.

Introduction

In response to the COVID-19 pandemic, the United Kingdom Government implemented a national “lockdown” involving school closures and social distancing. There has been widespread concern that these measures will negatively impact child and adolescent mental health.¹⁻³ To date, however, there is relatively little direct evidence of this. This is in part due to the paucity of studies including pre-lockdown baseline data. Longitudinal changes in mental health within the same individuals represent the most direct way of quantifying the impact of lockdown on children.

There is good reason to suspect that the implementation of a lockdown has a negative impact on children’s mental health. Early evidence from adult cohort studies suggest there has been a clinically meaningful increase in symptoms of anxiety, and a reduction in wellbeing, during lockdown.⁴⁻⁶ One recent systematic review found that loneliness in children was associated with subsequent mental health problems, particularly depression.⁷ There is a particular concern for children already struggling with mental health issues, where access to mental health services has been impacted.^{8,9}

Capitalising on existing baseline data from young children to assess lockdown-related changes in mental health is not straightforward, and this is why it has yet to be done. Unlike adults, these baseline data quickly become obsolete as children pass through different developmental stages, and it is not easy to re-contact families during a national lockdown. We report results from the Resilience in Education and Development (RED) study, a deeply phenotyped representative cohort of children living in the East of England.¹¹ This cohort had been assessed via a combination of parent-, teacher- and self-reports of mental health, alongside a variety of other measures. Around 18 months after this initial assessment, all of

these children were subject to the national lockdown. During lockdown, we contacted a subsample of the families and tested for any changes in their children's mental health and wellbeing. The aim of the study is to test whether changes in emotional wellbeing, anxiety and depression occurred during lockdown since the initial assessment.

Methods

Participants

The RED study comprises two groups. A larger school-group assessed in classrooms (N=567, from 22 classes, 6 schools), and a smaller lab-group (N=92) who completed the same and additional assessments at our Laboratory. Baseline assessments occurred between June 2018 and March 2019 in the school-group, and December 2018 and September 2019 in the laboratory-group. 6 schools were recruited to take part in the study, with all children in eligible year 3 & 4 classroom groups eligible to take part. Eligibility criteria in the lab-group included a medical screener for suitability to undergo medical resonance imaging.

The mental health assessments by parents and teachers, both before and during lockdown, were completed using an online survey. Participation was incentivised with a £5 amazon voucher for completion. We directly contacted all parents of children in the lab-group, and 5 schools contacted parents in the school-group, to complete the survey. One school (representing 84 children with baseline data) did not contact parents. 168 parents completed mental health assessments for their children during lockdown, for whom prior mental health data were available. Their children's (88 girls) mean age was 8.7 (SD = 0.64) and 10.1 (SD = 0.90) years, before and during lockdown testing, respectively. 8% were eligible for free school meals, an indicator of socioeconomic status (SES)¹².

119 *Measures*

120 Three mental health measures were used: the Strengths and Difficulties Questionnaire (SDQ)
121 Emotional Problems subscale, and RCADS-short form subscales for depression and anxiety.
122 ^{13,14} All scales were adapted for computerised testing using continuous slider scales.

123 Before lockdown, teachers and caregivers completed the SDQ for the school and lab-groups,
124 respectively. At this timepoint, children in both groups completed the RCADS, along with
125 parents in the lab-group. Children completed the RCADS on a custom-developed tablet
126 application¹¹, which included audio presentation of each question. Follow-up testing during
127 lockdown took place between 4th of April and 6th of June, 2020, with parents completing the
128 assessments.

129 *Statistical Analysis*

130 We analysed the impact of lockdown by combining child-, teacher- and parent-reports using
131 linear mixed models. Coefficients estimated the effect of lockdown (0=before/1=during
132 lockdown), and responder (0=child/1=parent or 0=teacher/1=parent), on children's mental
133 health, including a random intercept for participant. Children were included in the analysis if
134 an assessment before and during lockdown were available. Participant's age, gender and SES
135 were controlled in sensitivity analyses. SES was measured using a mean of: household
136 income, home-ownership, parent education, and neighbourhood deprivation.

137 Analyses were performed using R (v3.6.2) and the nlme (v.3.1-144) package. Mental health
138 variables were scored using the arithmetic mean response, after recoding each item so that a
139 higher score indicated worse mental health. For mixed linear models, all three mental health
140 outcomes were quantile-normalised to match a standard normal distribution. Normalisation

was performed for each outcome after transforming data into a “long” format, with repeated measurements (including before and during lockdown, from all raters) gathered in a single variable. Normalisation was performed by transforming variables into a percentile rank and applying the normal quantile function. Therefore, model coefficients can be interpreted as standardised effect sizes.

Role of the funding source

The funders had no role in the study design, data collection, data analysis, data interpretation, or writing of the manuscript. The corresponding author had full access to all the data and had the final responsibility for the decision to submit for publication.

Results

First, the baseline data enabled us to estimate the size of any recruitment biases in those who responded during lockdown. There was a small bias for more affluent families to participate. Neighbourhood deprivation (measured using the Index of Multiple Deprivation) and free school eligibility weakly predicted non-participation ($r = -0.17$ & -0.18 respectively).¹⁵ Greater child-reported RCADS depression ($r = -0.08$, 95% CI -0.16 , -0.01) and anxiety ($r = -0.09$, 95% CI -0.16 , -0.01) symptoms at baseline also weakly predicted non-participation.

Correlations between all variables are provided in Figure 1, alongside patterns of available data and missingness. All three mental health measures during lockdown were strongly correlated ($0.53 \leq r \leq 0.69$), though correlations between mental health reports before and during lockdown were generally low ($r \leq 0.34$). The internal consistency of each mental health scale was assessed using Cronbach’s alpha, which was above 0.74 for all measures except child-rated depression symptoms ($\alpha = 0.52$).

Table 1 reports all main effects. Without controlling for any additional confounders, there was no significant change in SDQ emotional problems ($B = -0.25$, 95% CI $-0.54, 0.05$) or RCADS anxiety scores ($B = -0.06$, 95% CI $-0.34, 0.23$) during lockdown compared to before. The confidence interval upper limits suggest that at most a small increase in these symptoms occurred during lockdown. This is consistent with the proportion of children with emotional problem scores in the “abnormal” range, which changed very little, decreasing from 13% (19 children) to 8% (12 children) from before to during lockdown.¹³ In contrast, standardised RCADS depression scores were on average 0.74 (95% CI 0.46, 1.01) higher during lockdown than before. The confidence intervals suggest a medium-to-large increase is likely.

Controlling for demographic factors separately (Age, Gender, and SES) did not strongly impact our estimates for these effects. Interaction effects of these three factors were estimated, to assess whether changes in mental health disproportionately occurred in certain groups. No interaction effects were statistically significant, albeit these estimates are highly uncertain.

A sensitivity analysis using only parent-rated mental health before and during lockdown was performed (see bottom table 1). One-sample t-tests were conducted on the standardised change scores, i.e. the raw score during lockdown minus the before-lockdown score, divided by the change score standard deviation. These analyses found similar effects as the mixed models (see Table 1).

One potential limitation of using mean scale scores is that changes during lockdown may be driven by specific items within the scale. Therefore, changes in responses to each individual question in the mental health scales were examined, using the same t-test approach outlined above (see Figure 3). 4 out of 5 of the depression questions showed significant increases

during lockdown. Only one other question (“Many worries, often seems worried”, from the SDQ) significantly changed, decreasing during lockdown.

Interpretation

National lockdowns with mass school closures are unprecedented, and the evidence-base to guide future policymaking has been slow to emerge. Longitudinal data forms a vital component of that evidence-base. This study is the first longitudinal demonstration that children’s depression ratings significantly increased during lockdown, relative to 18 months beforehand, with a medium-to-large effect.

Implications for policymakers and practitioners

The backdrop is that children’s mental health appears to be worsening across successive cohorts, and even before lockdown the resources for Child and Adolescent Mental Health Services were stretched thin.^{16,17} The current findings suggest that lockdown measures will likely exacerbate this, specifically with an increase in childhood depression, something previously relatively uncommon in children of this age.¹⁸ The education sector will bear the initial brunt of this.¹⁹ A key implication of the current findings is that a prompt return to school for all children must be prioritised by policymakers. When children return to school, their wellbeing, socialisation and enjoyment are paramount. Additional resources and training will likely be required to equip school teachers in how to support children with low mood, and to increase their awareness of referral pathways for professional support.

Future Directions

Future work should follow children over longer time periods to assess long-term effects. Firstly, because there is potential for “sleeping effects” (effects that emerge sometime after an initial adversity, often in a different phase of development)²⁰, and secondly because we need

to test whether children's mood rebounds when school resumes. Larger cohorts with greater statistical power are needed to address whether the epidemic has had disproportionate effects on particular children and households. The current study focuses on a broadly representative sample, but of particular concern are children with existing mental health and other needs, where effect sizes could be larger. For example, some initial reports have highlighted greater challenges during school closures facing children with autism and ADHD.^{21,22} Finally, our analysis of individual questionnaire items suggests that particular symptoms may be differentially affected by the lockdown. Epidemiological studies could further explore this potential differential impact of lockdown on mental health. One study in 80 Dutch students reported preliminary evidence that global mental health problems did not increase across 2 weeks of the COVID-19 pandemic, but depressive symptoms specifically increased, and anxiety symptoms decreased.^{23,24}

Study limitations

The sample size of the current study is constrained by needing recent baseline data on the same children pre-lockdown. As such, the lack of an effect on SDQ scores or RCADS anxiety scale should be interpreted with caution. The current study does not have the statistical power to detect small but clinically meaningful changes. Secondly, as with any study of this kind there is a small recruitment bias. This is directly relevant to all studies that are recruiting during lockdown. However, this cannot explain the significant findings we report: the biases are small relative to the size of the effect, and socio-economic status is controlled for within the linear models. Thirdly, the mixture of reporters is a limitation for the study. Children and adults report mental health symptoms differently. This is why reporter is directly incorporated within the model, and the effects are subsequently replicated in a subsample with just longitudinal parental report.

233 *Conclusions*

234 This study provides the first longitudinal evidence for the negative impact of UK lockdown
235 measures on child mental health. Specifically, with a significant increase in ratings of
236 depression, with a medium-to-large effect size. Our findings emphasise the critical need for a
237 prompt return to school, and a focus on young children's enjoyment and re-socialisation
238 rather than scholastic progress.

	SDQ Emotional Problems					RCADS Anxiety Subscale					RCADS Depression Subscale				
	B	95% CI		P	N	B	95% CI		P	N	B	95% CI		P	N
Model 1															
Lockdown	-.246	-.542	.050	.103	298/149	-.055	-.335	.225	.699	371/162	.736	.458	1.014	<.001	377/165
Responder	.276	-.056	.607	.102	298/149	-.796	-1.077	-.515	<.001	371/162	-1.331	-1.610	-1.052	<.001	377/165
Model 2															
Lockdown	-.161	-.537	.215	.398	298/149	.059	-.284	.402	.736	371/162	.580	.239	.920	.001	377/165
Responder	.274	-.059	.607	.106	298/149	-.796	-1.077	-.514	<.001	371/162	-1.302	-1.580	-1.023	<.001	377/165
Gender	-.025	-.351	.301	.881	298/149	-.033	-.289	.224	.802	371/162	-.328	-.573	-.082	.009	377/165
Lckdwn*Gndr	-.153	-.552	.245	.449	298/149	-.223	-.580	.134	.220	371/162	.249	-.106	.604	.169	377/165
Model 3															
Lockdown	-.111	-.505	.284	.580	296/148	.108	-.238	.455	.538	363/158	.722	.376	1.068	<.001	369/161
Responder	.331	-.006	.668	.055	296/148	-.790	-1.069	-.510	<.001	363/158	-1.320	-1.602	-1.037	<.001	369/161
Age	-.013	-.281	.256	.925	296/148	.003	-.212	.218	.979	363/158	.027	-.182	.236	.798	369/161
Lockdown*Age	-.215	-.511	.081	.153	296/148	-.222	-.477	.034	.089	363/158	-.031	-.289	.228	.816	369/161
Model 4															
Lockdown	-.255	-.551	.041	.090	298/149	-.060	-.340	.220	.674	371/162	.733	.455	1.011	<.001	377/165
Responder	.290	-.040	.620	.085	298/149	-.792	-1.073	-.510	<.001	371/162	-1.329	-1.608	-1.050	<.001	377/165
SES	-.140	-.301	.020	.086	298/149	-.060	-.187	.067	.353	371/162	-.051	-.174	.071	.409	377/165
Lockdown*SES	-.046	-.245	.153	.650	298/149	-.054	-.233	.126	.557	371/162	-.063	-.241	.115	.487	377/165
Paired T-test															
Lockdown	-.195	-.480	.089	.173	50	.145	-.136	.426	.305	51	.713	.432	.994	<.001	51
Note: N for mixed linear models gives the (Number of observations)/(Number of individuals). Continuous variables of age and SES were z-scored, and Lockdown and Responder are binary variables. Gender is coded as male (0) and female (1).															

Table 1: Main effects from Mixed Linear Model

SDQ Before (Teacher)	446	0	0	0	413	413	99	101	104
SDQ Before (Parent)	NA	78	78	78	71	71	50	51	51
GAD Before (Parent)	NA	.60	78	78	71	71	50	51	51
MDD Before (Parent)	NA	.51	.54	78	71	71	50	51	51
GAD Before (Child)	.22	.07	.19	.21	624	624	155	158	161
MDD Before (Child)	.12	-.05	.14	.21	.49	624	155	158	161
SDQ During (Parent)	.27	.24	.14	.22	.17	-.07	164	163	164
GAD During (Parent)	.23	.08	.15	.23	.17	.05	.69	167	167
MDD During (Parent)	-.00	.09	.15	.34	.10	.02	.53	.56	171
	SDQ Before (Teacher)	SDQ Before (Parent)	GAD Before (Parent)	MDD Before (Parent)	GAD Before (Child)	MDD Before (Child)	SDQ During (Parent)	GAD During (Parent)	MDD During (Parent)

Figure 1: Correlations between mental health variables and patterns of missing data.

Below Diagonal: Correlations between SDQ Emotional Problems (SDQ), RCADS anxiety subscale (GAD) and RCADS depression subscale (MDD), before and during lockdown. On diagonal: Number of observations for each variable. Above diagonal: Number of observations with data on two given variables.

Figure 2: Change in RCADS-depression score from before to during the lockdown measures.

All children were asked to rate their own depression (orange points) before lockdown. Parents (blue points) completed the RCADS before lockdown for a subsample ($n = 51$) of participants, and for all participants during lockdown. Error bars represent the estimated marginal means, grouped by child-report (orange bars) and parent-report (blue bars).

Figure 3: Mean change and confidence intervals for each mental health question, comparing during to before lockdown, using solely parent reported mental health at both time points.

Positive numbers indicate worsening of symptom during lockdown. Results support the interpretation that RCADS depression items have increased during lockdown, with more minor changes in other symptoms.

Declaration of interests

The authors have no conflicts of interest relevant to this article to disclose.

Data Sharing Statement

96% of the participants gave consent for their data to be shared, on the condition that researchers applying for access have the appropriate ethical permission. Analysis code and R output are available from <https://osf.io/ajy57>.

Contributors

Bignardi conceptualised and designed the study, collected the data, drafted the initial manuscript, carried out the statistical analysis, and reviewed and revised the manuscript.

Astle conceptualised and designed the study, drafted the initial manuscript, supervised the study, obtained funding, and reviewed and revised the manuscript. Dalmaijer conceptualised and designed the study, collected the data, supervised the study, and reviewed and revised the manuscript. Dr Siugzdaite, Anwyl-Irvine, Smith and Uh conceptualised and designed the study, collected the data, and reviewed and revised the manuscript.

References

1. Brooks SK, Webster RK, Smith LE, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet*. 2020;395(10227):912-920. doi:10.1016/S0140-6736(20)30460-8
2. Lee J. Mental health effects of school closures during COVID-19. *Lancet Child Adolesc Heal*. 2020;4(6):421. doi:10.1016/S2352-4642(20)30109-7
3. Orben A, Tomova L, Blakemore S-J. The effects of social deprivation on adolescent development and mental health. *Lancet Child Adolesc Heal*. June 2020. doi:10.1016/S2352-4642(20)30186-3
4. Kwong ASF, Pearson RM, Adams MJ, et al. Mental health during the COVID-19

pandemic in two longitudinal UK population cohorts. *medRxiv*. January 2020:2020.06.16.20133116. doi:10.1101/2020.06.16.20133116

5. Pierce M, Hope H, Ford T, et al. Mental health before and during the COVID-19 pandemic: a longitudinal probability sample survey of the UK population. *The Lancet Psychiatry*. July 2020. doi:10.1016/S2215-0366(20)30308-4
6. Li HY, Cao H, Leung DYP, Mak YW. The Psychological Impacts of a COVID-19 Outbreak on College Students in China: A Longitudinal Study. *Int J Environ Res Public Health*. 2020;17(11):3933. doi:10.3390/ijerph17113933
7. Loades ME, Chatburn E, Higson-Sweeney N, et al. Rapid Systematic Review: The Impact of Social Isolation and Loneliness on the Mental Health of Children and Adolescents in the Context of COVID-19. *J Am Acad Child Adolesc Psychiatry*. June 2020. doi:10.1016/j.jaac.2020.05.009
8. Witt A, Ordóñez A, Martin A, Vitiello B, Fegert JM. Child and adolescent mental health service provision and research during the Covid-19 pandemic: challenges, opportunities, and a call for submissions. *Child Adolesc Psychiatry Ment Health*. 2020;14(1):19. doi:10.1186/s13034-020-00324-8
9. Ford T. Prof Tamsin Ford: Challenges for children's services as Lockdown eases. Reachwell. <https://reachwell.org/2020/07/16/prof-tamsin-ford-challenges-for-childrens-services-as-lockdown-eases/>. Published 2020. Accessed July 21, 2020.
10. Xie X, Xue Q, Zhou Y, et al. Mental Health Status Among Children in Home Confinement During the Coronavirus Disease 2019 Outbreak in Hubei Province, China. *JAMA Pediatr*. April 2020. doi:10.1001/jamapediatrics.2020.1619
11. Bignardi G, Dalmaijer E, Anwyl-Irvine A, Astle D. *Collecting Big Data with Small Screens: Group-Tests of Children's Cognition with Touchscreen Tablets Are Reliable and Valid.*; 2020. doi:10.31219/osf.io/aw6c5
12. Taylor C. The Reliability of Free School Meal Eligibility as a Measure of Socio-Economic Disadvantage: Evidence from the Millennium Cohort Study in Wales. *Br J Educ Stud*. 2018;66(1):29-51. doi:10.1080/00071005.2017.1330464
13. Goodman R. The Strengths and Difficulties Questionnaire: A Research Note. *J Child Psychol Psychiatry*. 1997;38(5):581-586. doi:10.1111/j.1469-7610.1997.tb01545.x
14. Muris P, Meesters C, Schouten E. Questionnaire of Anxiety and Depression Children. *Clin Psychol Psychother*. 2002;9:430-442.
15. Smith T, Noble M, Noble S, et al. The English indices of deprivation 2015. *London Dep Communities Local Gov*. 2015.
16. Sellers R, Warne N, Pickles A, Maughan B, Thapar A, Collishaw S. Cross-cohort change in adolescent outcomes for children with mental health problems. *J Child Psychol Psychiatry*. 2019;60(7):813-821. doi:10.1111/jcpp.13029

17. Crenna-Jennings W, Hutchinson J. *Access to Child and Adolescent Mental Health Services in 2019*.; 2020.
18. Ghandour RM, Sherman LJ, Vladutiu CJ, et al. Prevalence and Treatment of Depression, Anxiety, and Conduct Problems in US Children. *J Pediatr*. 2019;206:256-267.e3. doi:10.1016/j.jpeds.2018.09.021
19. Newlove-Delgado T, Moore D, Ukoumunne OC, Stein K, Ford T. Mental health related contact with education professionals in the British Child and Adolescent Mental Health Survey 2004. *J Ment Heal Training, Educ Pract*. 2015;10(3):159-169. doi:10.1108/JMHTEP-02-2015-0007
20. Wade M, Prime H, Browne DT. Why we need longitudinal mental health research with children and youth during (and after) the COVID-19 pandemic. *Psychiatry Res*. 2020;290:113143. doi:10.1016/j.psychres.2020.113143
21. Bobo E, Lin L, Acquaviva E, et al. Comment les enfants et adolescents avec le trouble déficit d'attention/hyperactivité (TDAH) vivent-ils le confinement durant la pandémie COVID-19 ? *Encephale*. 2020;46(3):S85-S92. doi:10.1016/j.encep.2020.05.011
22. Colizzi M, Sironi E, Antonini F, Ciceri ML, Bovo C, Zoccante L. Psychosocial and Behavioral Impact of COVID-19 in Autism Spectrum Disorder: An Online Parent Survey. *Brain Sci*. 2020;10(6):341. doi:10.3390/brainsci10060341
23. Eiko F. Preprint: COVID-19 pandemic and student mental health. 2020. <https://eiko-fried.com/preprint-covid-19-pandemic-and-student-mental-health/>. Accessed July 22, 2020.
24. Fried E., Papanikolaou F, Epskamp S. *Mental Health and Social Contact During the COVID-19 Pandemic: An Ecological Momentary Assessment Study*.; 2020. doi:10.31234/osf.io/36xkp