NTRK2 is differentially expressed in the brains of patients with psychotic disorders.

Shahan Mamoor¹

¹shahanmamoor@gmail.com

East Islip, NY 11730

We used published microarray data^{1,2} to identify the most significant gene expression changes in the brains of patients with psychotic disorders. We identified NTRK2 as differentially expressed in the dorsolateral prefrontal cortex of patients with schizophrenia as well as in the dorsolateral prefrontal cortex of patients with schizophrenia and psychotic bipolar disorder. NTRK2 expression was significantly higher in the dorsolateral prefrontal cortex of patients with psychotic disorders.

Keywords: schizophrenia, psychotic disorders, systems biology of schizophrenia, psychosis, non-coding RNAs in human disease.

Psychotic disorders, including schizophrenia, are characterized by the presence of psychosis, or an inability to engage in reality testing^{3,4}. Patients with psychotic disorders can experience positive as well as negative symptoms⁵. Positive symptoms include hallucinations, visual and auditory, as well as delusions, or fixed false beliefs⁵. Negative symptoms include anhedonia, or an inability to experience pleasure, as well as avolition, or loss of motivation⁵. To understand how specific regions of the brain differ most significantly in terms of gene expression in patients with psychotic disorders, we used published microarray data to compare global gene expression profiles of brain tissues from patients with psychotic disorders with that of non-affected, control subjects^{1,2}. We found that NTRK2 was among the genes whose expression changed most significantly in patients with schizophrenia in the dorsolateral prefrontal cortex, and in a separate dataset, in the dorsolateral prefrontal cortex in patients with schizophrenia and psychotic bipolar disorder. NTRK2 may be relevant for future study of genes pertinent to the biology of psychotic disorders.

Methods

We used microarray datasets GSE87610¹ and GSE120340² for this differential gene expression analysis in conjunction with GEO2R. GSE87610 was generated using Affymetrix Human Genome U219 with n=72 for control subjects and n=65 for patients with schizophrenia. GSE120340 was generated using Affymetrix Human Genome U133 Plus 2.0 Array technology with post-mortem tissues from Brodmann's area 46 from control subjects (n=10 samples utilized in this analysis) and from patients with psychotic disorders (schizophrenia and psychotic bipolar disorder) (n=16 samples utilized in this analysis). The Benjamini and Hochberg method of p-value adjustment was used for ranking of differential expression but raw p-values were used for assessment of statistical significance of global differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. A statistical test was performed to evaluate whether NTRK2

expression was significantly different when comparing normal DLPFC tissue from control subjects and DLPFC from patients with schizophrenia or schizophrenia and psychotic bipolar disorder using a two-tailed, unpaired t-test with Welch's correction. Only p-values less than 0.05 were considered statistically significant. We used PRISM for all statistical analyses (Version 8.4.0)(455).

Results

We mined published microarray datasets^{1,2} to identify genes whose expression was most different in the brains of patients with psychotic disorders when compared to control subjects.

NTRK2 is differentially expressed in the dorsolateral prefrontal cortex of patients with schizophrenia.

We identified NTRK2 as among the genes whose mRNA expression was most different in the dorsolateral prefrontal cortex of patients with schizophrenia when compared to control subjects¹. When sorting all of the genes expressed in the dorsolateral prefrontal cortex, as measured by microarray, based on change in expression between psychotic patients and control subjects, NTRK2 ranked 2 out of 12188 total transcripts. Differential expression of NTRK2 in the dorsolateral prefrontal cortex of patients with schizophrenia was statistically significant (Table 1; $p=1.75E-11$).

NTRK2 is differentially expressed in the dorsolateral prefrontal cortex of patients with schizophrenia and psychotic bipolar disorder.

In a separate dataset, we identified NTRK2 as differentially expressed in the dorsolateral prefrontal cortex of patients with schizophrenia or psychotic bipolar disorder when compared to control subjects². When sorting all of the genes expressed in the dorsolateral prefrontal cortex, as measured by microarray, based on change in expression between patients with schizophrenia or psychotic bipolar disorder and control subjects, NTRK2 ranked 137 out of

19070 total transcripts. Differential expression of NTRK2 in the DLPFC of patients with schizophrenia or psychotic bipolar disorder was statistically significant (Table 2; $p=0.0015125$).

NTRK2 expression is significantly higher in the dorsolateral prefrontal cortex of patients with schizophrenia.

Next we obtained exact mRNA expression levels of NTRK2 from the dorsolateral prefrontal cortex of patients with schizophrenia and from the dorsolateral prefrontal cortex of control subjects to directly compare NTRK2 expression between these groups. We also performed a statistical test to evaluate whether this difference in expression was statistically significant. NTRK2 was expressed at significantly higher levels in the dorsolateral prefrontal cortex of patients with schizophrenia when compared to control subjects (Figure 1; $p<0.0001$). We calculated a 1.3036 ± 0.2296 fold change in NTRK2 expression in the dorsolateral prefrontal cortex of patients with schizophrenia when compared to control subjects (Table 1).

NTRK2 expression is significantly higher in the dorsolateral prefrontal cortex of patients with schizophrenia and psychotic bipolar disorder.

We also obtained exact mRNA expression levels of NTRK2 the dorsolateral prefrontal cortex of patients with schizophrenia or psychotic bipolar disorder and from the dorsolateral prefrontal cortex of control subjects to directly compare NTRK2 between these two groups. We also performed a statistical test to evaluate whether this difference in expression was statistically significant. NTRK2 was expressed at significantly higher levels in the dorsolateral prefrontal cortex of patients with schizophrenia or psychotic bipolar disorder when compared to control subjects (Figure 2; $p=0.0022$). We calculated a 1.0845 ± 0.0673 fold change in NTRK2 expression in the dorsolateral prefrontal cortex of patients with schizophrenia and psychotic bipolar disorder when compared to control subjects (Table 2).

Thus, we found using published microarray datasets^{1,2} that NTRK2 was among the genes most differentially expressed in the brain tissues, in the dorsolateral prefrontal cortex of patients with psychotic disorders.

Discussion

We compared transcriptome information from published microarray datasets^{1,2} generated using the brain tissues of patients with psychotic disorders, specifically, in the dorsolateral prefrontal cortex, to that of non-affected control subjects, to identify the most significant changes in gene expression in the brains of patients with psychotic disorders. We found that changes in NTRK2 gene expression were more significant than greater than 99% of transcriptome in group of patients.

There is a wide body of literature that suggests that alterations in NTRK2 at the genetic level may be implicated with the risk of development of psychiatric disorders. Variants in NTRK2 have been associated with depressive disorders in temporal lobe epilepsy⁶ and as protective against obsessive compulsive disorder⁷. NTRK2 polymorphism trended toward association with the risk for development of autism⁸. In patients with first-episode psychosis receiving antipsychotic therapy, NTRK2 polymorphism was associated with serum prolactin levels⁹. A study of Han Chinese population suggested that interactions between polymorphisms in brain-derived neurotrophic factor (BDNF) and NTRK2 might be genetic risk factors for developing paranoid schizophrenia¹⁰. Genome-wide analysis of single-nucleotide polymorphisms in two German cohorts demonstrated that polymorphisms in NTRK2 were associated with the risk of suicidal attempt in depressed patients¹¹. In a subgroup of a separate African American cohort, risk allele (C) at polymorphism rs1147198 was significantly more common in patients with lifelong major depressive disorder that had completed suicidal attempts¹¹. One further study using an International Affective Picture System (IAPS) and

diffuse tensor imaging showed an association between a single nucleotide polymorphism in NTRK2 and emotional arousal as well as brain white-matter mean diffusivity¹².

It is of interest to determine experimentally whether changes in NTRK2 expression in the dorsolateral prefrontal cortex of patients with schizophrenia are associated with disease state or antipsychotic regimen. We found here that small but statistically significant changes in NTRK2 gene expression were among the most significant transcriptome-wide in the dorsolateral prefrontal cortex of patients with a spectrum of psychotic disorders.

References

1. Arion, D., Huo, Z., Enwright, J.F., Corradi, J.P., Tseng, G. and Lewis, D.A., 2017. Transcriptome alterations in prefrontal pyramidal cells distinguish schizophrenia from bipolar and major depressive disorders. *Biological psychiatry*, 82(8), pp.594-600.
2. Abdolmaleky, H.M., Gower, A.C., Wong, C.K., Cox, J.W., Zhang, X., Thiagalingam, A., Shafa, R., Sivaraman, V., Zhou, J.R. and Thiagalingam, S., 2019. Aberrant transcriptomes and DNA methylomes define pathways that drive pathogenesis and loss of brain laterality/asymmetry in schizophrenia and bipolar disorder. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*, 180(2), pp.138-149.
3. Cannon, T.D., 2015. How schizophrenia develops: cognitive and brain mechanisms underlying onset of psychosis. *Trends in cognitive sciences*, 19(12), pp.744-756.
4. Arciniegas, D.B., 2015. Psychosis. *Continuum: Lifelong Learning in Neurology*, 21(3 Behavioral Neurology and Neuropsychiatry), p.715.
5. Andreasen, N.C., Nopoulos, P., Schultz, S., Miller, D., Gupta, S., Swayze, V. and Flaum, M., 1994. Positive and negative symptoms of schizophrenia: past, present, and future. *Acta Psychiatrica Scandinavica*, 90, pp.51-59.
6. Torres, C.M., Siebert, M., Bock, H., Mota, S.M., Castan, J.U., Scornavacca, F., de Castro, L.A., Saraiva-Pereira, M.L. and Bianchin, M.M., 2017. Tyrosine receptor kinase B gene variants (NTRK2 variants) are associated with depressive disorders in temporal lobe epilepsy. *Epilepsy & Behavior*, 71, pp.65-72.
7. Alonso, P., Gratacòs, M., Menchón, J.M., Saiz-Ruiz, J., Segalàs, C., Baca-García, E., Labad, J., Fernández-Piqueras, J., Real, E., Vaquero, C. and Pérez, M., 2008. Extensive genotyping of the BDNF and NTRK2 genes define protective haplotypes against obsessive-compulsive disorder. *Biological psychiatry*, 63(6), pp.619-628.
8. Correia, C.T., Coutinho, A.M., Sequeira, A.F., Sousa, I.G., Lourenco Venda, L., Almeida, J.P., Abreu, R.L., Lobo, C., Miguel, T.S., Conroy, J. and Cochrane, L., 2010. Increased BDNF levels and NTRK2 gene association suggest a disruption of BDNF/TrkB signaling in autism. *Genes, Brain and behavior*, 9(7), pp.841-848.
9. Gassó, P., Mas, S., Bioque, M., Cabrera, B., Lobo, A., González-Pinto, A., Díaz-Caneja, C.M., Corripio, I., Vieta, E., Castro-Fornieles, J. and Sarró, S., 2018. Impact of NTRK2, DRD2 and ACE polymorphisms on prolactin levels in antipsychotic-treated patients with first-episode psychosis. *Journal of Psychopharmacology*, 32(6), pp.702-710

10. Lin, Z., Su, Y., Zhang, C., Xing, M., Ding, W., Liao, L., Guan, Y., Li, Z. and Cui, D., 2013. The interaction of BDNF and NTRK2 gene increases the susceptibility of paranoid schizophrenia. *PLoS One*, 8(9), p.e74264.
11. Kohli, M.A., Salyakina, D., Pfennig, A., Lucae, S., Horstmann, S., Menke, A., Kloiber, S., Hennings, J., Bradley, B.B., Ressler, K.J. and Uhr, M., 2010. Association of genetic variants in the neurotrophic receptor–encoding gene NTRK2 and a lifetime history of suicide attempts in depressed patients. *Archives of general psychiatry*, 67(4), pp.348-359.
12. Spalek, K., Coyne, D., Freytag, V., Hartmann, F., Heck, A., Milnik, A., de Quervain, D. and Papassotiropoulos, A., 2016. A common NTRK2 variant is associated with emotional arousal and brain white-matter integrity in healthy young subjects. *Translational psychiatry*, 6(3), pp.e758-e758.

Rank	ID	p-value	t	B	Fold change	Gene
2	11720079_at	1.75E-11	7.336003	15.23611	1.3036 ± 0.2296	NTRK2

Table 1: NTRK2 is differentially expressed in the dorsolateral prefrontal cortex of patients with schizophrenia when compared to control subjects.

The rank of differential expression, the probe/transcript ID, the p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, the fold change of NTRK2 in the patient population when compared to control subjects, and gene symbol are listed in this chart.

Rank	ID	p-value	t	B	Fold change	Gene name
137	4915_at	0.0015125	3.52149751	-1.06261	1.0845 ± 0.0673	neurotrophic tyrosine kinase, receptor, type 2

Table 2: NTRK2 is differentially expressed in the dorsolateral prefrontal cortex of patients with schizophrenia or psychotic bipolar disorder when compared to control subjects.

The rank of differential expression, the probe/transcript ID, the p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, the fold change of NTRK2 in the patient population when compared to control subjects and gene name are listed in this chart.

Figure 1: NTRK2 is expressed at significantly higher levels in the dorsolateral prefrontal cortex of patients with schizophrenia when compared to control subjects.

The mRNA expression of NTRK2 in the dorsolateral prefrontal cortex of control subjects (left) and in the dorsolateral prefrontal cortex of patients with schizophrenia (right) is represented with mean mRNA expression level marked and the result of a statistical test evaluating the significance of difference in NTRK2 expression between the dorsolateral prefrontal cortex of control subjects and patients with schizophrenia, a p-value, listed above.

Figure 2: NTRK2 is expressed at significantly higher levels in the dorsolateral prefrontal cortex of patients with schizophrenia or psychotic bipolar disorder than in the DLPFC of control subjects.

The mRNA expression of NTRK2 in the dorsolateral prefrontal cortex of control subjects (left) and in the dorsolateral prefrontal cortex of patients with schizophrenia or psychotic bipolar disorder (right) is represented with mean mRNA expression level marked and the result of a statistical test evaluating the significance of difference in NTRK2 expression between the dorsolateral prefrontal cortex of control subjects and patients with schizophrenia or psychotic bipolar disorder, a p-value, listed above.