Research into anatomical abnormalities in South Asia

Jelle Stans ¹ *

¹ Institute for Globally Distributed Open Research and Education, Beringen, Belgium

* jelle.stans@igdore.org

Anatomical and congenital abnormalities can vary widely in their clinical severity. While anencephaly is quasi invariably lethal (1), cervical ribs may not pose any clinical problems at all (2). Research is essential to investigate the epidemiology, clinical implications and potential treatments. Access to these studies may allow patients to treatment strategies that are otherwise not yet available. Therefore, it is important to investigate where studies into anatomical abnormalities are conducted and what these studies investigate. This short letter aims to characterise the studies into anatomical abnormalities in South Asia. For this letter, South Asia was defined as consisting of Afghanistan, Bangladesh, Bhutan, India, Nepal, Pakistan and Sri Lanka.

The clinicaltrials.gov database was searched for relevant studies using the keywords “Anatomical Abnormality”. Afterwards, the search results were limited to studies in South Asia using relevant filters.

A total of 12 relevant studies were identified in South Asia. Eight of these studies were being conducted in India, five in Pakistan and two in Sri Lanka. No studies were identified in Afghanistan, Bangladesh, Bhutan and Nepal. The total number of studies in the region was significantly smaller than other regions such as Europe with 530 studies and North America with 639 studies, despite South Asia having a larger population than both regions. Southeast Asia and North Asia had a similar number of studies, 23 and 15 respectively. East Asia had a larger number of studies (147), with the majority of these studies being conducted in China.

Two out of the studies investigated spinal deformities, one congenital haemangioma, one osteoarthritis in the knee, one clubfoot, one leg length discrepancy and knee deformities, one end stage renal disease, one hypomelanotic macules, one arteriovenous malformations, one skin

pigmentation over face deformities and one microcephaly. The interventions investigated ranged widely and included drugs, surgical procedures and diagnostic tests.

Only three of the studies were currently recruiting, while five were already completed. Of three studies, the status was unknown. One study was active but not recruiting. Unfortunately, none of the studies has results available. Half of the identified studies were observational, while the other half was interventional.

To summarize, there was only a relatively small number of studies being conducted into anatomical abnormalities in the region. These studies however covered a broad range of different abnormalities and tested several interventions. Further research should investigate what the reasons are for this lack of studies and what can be done to stimulate this kind of research.

References

1. Jaquier M, Klein A, Boltshauser E. Spontaneous pregnancy outcome after prenatal diagnosis of anencephaly. BJOG. 2006;113(8):951-3.
2. Spadliński Ł, Cecot T, Majos A, Stefańczyk L, Pietruszewska W, Wysocki G, et al. The Epidemiological, Morphological, and Clinical Aspects of the Cervical Ribs in Humans. Biomed Res Int. 2016;2016:8034613.