The criteria to classify body-proportionality of the small for gestational age newborn: a scoping review protocol

Berkay Karadavut¹, Ilke Smits¹, Jeroen van Dillen² and Marije Hogeveen¹

- 1 Department of Pediatrics, Division of Neonatology, Radboud University Medical Center Amalia Children's Hospital, Nijmegen, The Netherlands
- 2 Department of Obstetrics and Gynecology, Radboud University Medical Center, Nijmegen, The Netherlands

Correspondence: Marije Hogeveen, marije.hogeveen@radboudumc.nl

List of abbreviations:

AC – Abdominal circumference

BW – Birth weight

EFW – Estimated fetal weight

FGR – Fetal growth restriction

HC – Head circumference

IUGR – Intrauterine growth restriction

PI – Ponderal Index

SGA – Small for gestational age

Box 1: Definitions**Fetal growth restriction**

Conceptual definition: The fetus does not reach its full growth potential due to some underlying pathological process.

Statistical definition: Abdominal circumference (AC) or estimated fetal weight (EFW) are < 10th percentile on an intra-uterine growth standard; or a decrease of > 20 percentile points of EFW and/or AC over a time period of > 2 weeks.

Intrauterine growth restriction

Another term used for fetal growth restriction.

Small for gestational age

Conceptual definition: The smallest newborns of a population entailing both newborns who have reached their full growth potential but who are genetically small, and growth restricted newborns.

Statistical definition: Birthweight below a pre-defined threshold for gestational age (generally below the 10th percentile for the gestational age on a sex-specific birthweight standard).

References

- Mandy GT. Infants with fetal (intrauterine) growth restriction. In: Weisman LE, Kim MS (Editors). Waltham, MA: UpToDate;2020 [cited on March 27, 2021].
- Nederlandse Vereniging voor Obstetrie en Gynaecologie. Foetale groeirestrictie (FGR). [Utrecht]: NVOG; 2017 [updated 2018 Aug; cited March 27, 2021]. (Clinical guideline). Available from: <https://www.nvog.nl/wp-content/uploads/2018/08/NVOG-richtlijn-FGR->

Introduction

When a newborn has a birthweight below a predefined threshold that newborn is classified as small for gestational age (SGA). The statistical definition most commonly used is the 10th percentile for sex and gestational age on a birthweight standard (1-3). Conceptionally, SGA does not only include newborns who are constitutionally small, meaning the newborn is small in comparison to the majority of the population due to physiological factors such as genetics, but also newborns who are growth restricted (fetal growth restriction(FGR))(box 1) (1-4). It should be noted that in some studies the terms intrauterine growth restriction (IUGR) or fetal growth restriction (FGR) are used to define a birthweight below the 10th percentile instead of SGA (5, 6). Although these terms are often used interchangeably, they do not refer to identical concepts nor definitions (box) (6, 7). For the sake of clarity, in this review we will use the term SGA for newborns with a birthweight below the 10th percentile, irrespective of the term used in the original article.

In the short term, SGA newborns are at increased risk for several adverse outcomes such as perinatal mortality, meconium aspiration, hypoglycemia, polycythemia and hypothermia (8). In the long term, SGA newborns are at increased risk for developmental disorders, neurodegenerative illnesses, growth failure, reduced strength and cardiovascular diseases (8-10).

SGA newborns can be categorized according to their body-proportionality as symmetric SGA and as asymmetric SGA (5, 11). Symmetric growth restriction may occur when it originates early in pregnancy and asymmetric growth restriction may develop when it arises during late pregnancy (5, 11). Risk factors vary from maternal risk factors to smoking to placental insufficiency to congenital infections (12, 13).

Body-proportionality in the SGA newborn may influence the incidence and prevalence of adverse outcomes. For example, newborns who are asymmetric SGA have been shown to be more likely to develop postpartum hypoglycemia and to be more likely to have memory difficulties than symmetric SGA newborns (14, 15).

Several definitions of body-proportionality can be found: a head circumference (HC) to birth weight (BW) ratio (11), or a birth weight to length ratio, using the ponderal index ($PI = \text{kg/m}^3$) to measure the latter (5).

This apparent lack of uniform criteria to define body proportions in SGA newborns limits interpretation of research in this area. A preliminary search of PROSPERO, Cochrane Database of Systematic Reviews and the JBI Database of Systematic Reviews was conducted on January 20th, 2021 and no current reviews on this topic were found.

The objective of this scoping review will be to broadly examine the published criteria to classify body-proportionality in SGA newborns with the aims to identify knowledge gaps and to map available evidence. This overview will pave the way for a future consensus definition of body-proportionality of the SGA newborn, which will lead to a better comparison of study results so that robust conclusions can be drawn.

Research questions

1. Which criteria are used to classify the body-proportionality of the small for gestational age newborn?
2. What are the outcomes related to body-proportionality?

Inclusion criteria

Population

This scoping review will include articles about SGA newborns regardless of gestational age in which some type of classification is made into symmetric and asymmetric SGA. This review will exclude studies involving animals and studies which only describe body-proportionality in FGR.

Concept

The criteria used to define the body-proportionality of the SGA newborn will be summarized in this scoping review.

Context

Studies will not be limited to a geographic location. As long as the concept of body-proportionality in the SGA newborn is clearly described, the study can be included.

Types of evidence sources

This scoping review will consider all available study designs, however gray and unpublished literature will not be included due to the broadness of the topic. Conference proceedings will be excluded. Reference lists and citations of included studies will also be considered for inclusion.

Methods

A scoping review will be undertaken to map available evidence on this topic (16). The review will be conducted according to the guidance from the Joanna Briggs Institute (17).

Search strategy

Based on a preliminary limited search of MEDLINE (PubMed), a list of index terms and text words is incorporated in a search strategy developed with the help of librarian and according to Peer Review of Electronic Search Strategies (PRESS) 2015 Evidence-Based Checklist (18) (see Appendix I). Based on the research question the search strategy contains two main topics which are put in search blocks. The first search block contains the population "SGA newborn" and the second search block contains the concept "body-proportionality". These two search blocks will be combined to form the search strategy. The reference lists of included studies will be examined (17). Languages will be restricted to English and Dutch.

Source of evidence selection

This scoping review will include MEDLINE (PubMed), Embase (OVID) and the Cochrane Library as sources for information. Articles published before 19 February 2021 will be screened.

Study selection

Studies will be selected according to a step-by-step process. After executing the search strategy in each of the three databases, all the results will be imported into EndNote X9. The results will be

deduplicated according to Bramer et al (19). After deduplication the results will be imported into Rayyan for phase 1 selection. Studies will undergo a title/abstract screening and will be categorized as “clearly not eligible” and “maybe eligible” by one reviewer using the Rayyan software after pilot testing of 10-25 articles. The phase 1 results in the “maybe eligible” category will be imported into Rayyan as phase 2. Two reviewers will independently screen the title/abstract of the articles and categorize them as “not eligible”, “maybe eligible” or “eligible”. The phase 2 results in the “maybe eligible” and the “eligible” categories will undergo a full-text screening after discrepancies between the two reviewers are solved through discussion in the research team. The phase 2 results in the “maybe eligible” and the “eligible” categories will be imported into Rayyan as phase 3. Two reviewers will independently screen these full-text papers. Inclusion criteria are:

Population: the study provides information on newborns who are classified as SGA;

Concept: the study reports on any classification of body-proportionality of the SGA group.

A reason of exclusion will be given for each excluded paper. References of the included articles will be checked and considered for inclusion. Discrepancies will be discussed with the research team again.

Data extraction

Two reviewers will independently extract the data using a data extraction form based on the template from the Joanna Briggs Institute (17). This form will be pilot tested on 2-5 articles. The extracted data will include basic information like first author, title, publication year, country and also subject specific information such as the study population, number of patients, setting, definition of SGA, definition of body-proportionality and the outcome related to body-proportionality. Discrepancies will be discussed with the entire team.

Data presentation

Extracted data will be presented in a representative manner in tabular form (see Appendix II) accompanied by a narrative summary.

Acknowledgements

The authors acknowledge the support and resources provided by the Radboud University and the Radboud University Medical Center.

References

1. Hoftiezer L, Hof MHP, Dijs-Elsinga J, Hogeveen M, Hukkelhoven C, van Lingen RA. From population reference to national standard: new and improved birthweight charts. *Am J Obstet Gynecol*. 2019;220(4):383.e1-.e17.
2. Beune IM, Bloomfield FH, Ganzevoort W, Embleton ND, Rozance PJ, van Wassenhaer-Leemhuis AG, et al. Consensus Based Definition of Growth Restriction in the Newborn. *J Pediatr*. 2018;196:71-6.e1.
3. Schlaudecker EP, Munoz FM, Bardají A, Boghossian NS, Khalil A, Mousa H, et al. Small for gestational age: Case definition & guidelines for data collection, analysis, and presentation of maternal immunisation safety data. *Vaccine*. 2017;35(48 Pt A):6518-28.

4. Gordijn SJ, Beune IM, Thilaganathan B, Papageorgiou A, Baschat AA, Baker PN, et al. Consensus definition of fetal growth restriction: a Delphi procedure. *Ultrasound Obstet Gynecol.* 2016;48(3):333-9.
5. Landmann E, Reiss I, Misselwitz B, Gortner L. Ponderal index for discrimination between symmetric and asymmetric growth restriction: percentiles for neonates from 30 weeks to 43 weeks of gestation. *J Matern Fetal Neonatal Med.* 2006;19(3):157-60.
6. Mandy GT. Infants with fetal (intrauterine) growth restriction. In: Weisman LE, Kim MS (Editors). Waltham, MA: UpToDate;2020 [cited on March 18, 2021].
7. Physical status: the use and interpretation of anthropometry. Report of a WHO Expert Committee. *World Health Organ Tech Rep Ser.* 1995;854:1-452.
8. Sharma D, Farahbakhsh N, Shastri S, Sharma P. Intrauterine growth restriction - part 2. *J Matern Fetal Neonatal Med.* 2016;29(24):4037-48.
9. Hadar O, Sheiner E, Wainstock T. The Association between Delivery of Small-for-Gestational-Age Neonate and Their Risk for Long-Term Neurological Morbidity. *J Clin Med.* 2020;9(10).
10. de Boo HA, Harding JE. The developmental origins of adult disease (Barker) hypothesis. *Aust N Z J Obstet Gynaecol.* 2006;46(1):4-14.
11. Bocca-Tjeertes I, Bos A, Kerstjens J, de Winter A, Reijneveld S. Symmetrical and asymmetrical growth restriction in preterm-born children. *Pediatrics.* 2014;133(3):e650-6.
12. McCowan L, Horgan RP. Risk factors for small for gestational age infants. *Best Pract Res Clin Obstet Gynaecol.* 2009;23(6):779-93.
13. Albu AR, Anca AF, Horhoianu VV, Horhoianu IA. Predictive factors for intrauterine growth restriction. *J Med Life.* 2014;7(2):165-71.
14. Bhat MA, Kumar P, Bhansali A, Majumdar S, Narang A. Hypoglycemia in small for gestational age babies. *Indian J Pediatr.* 2000;67(6):423-7.
15. Geva R, Eshel R, Leitner Y, Fattal-Valevski A, Harel S. Memory functions of children born with asymmetric intrauterine growth restriction. *Brain Res.* 2006;1117(1):186-94.
16. Munn Z, Peters MDJ, Stern C, Tufanaru C, McArthur A, Aromataris E. Systematic review or scoping review? Guidance for authors when choosing between a systematic or scoping review approach. *BMC Med Res Methodol.* 2018;18(1):143.
17. Peters MDJ GC, McInerney P, Munn Z, Tricco AC, Khalil, H. Chapter 11: Scoping Reviews (2020 version). In: Aromataris E, Munn Z (Editors). *JBIM Manual for Evidence Synthesis*, JBI, 2020. Available from <https://synthesismanual.jbi.global>. <https://doi.org/10.46658/JBIMES-20-12>.
18. McGowan J, Sampson M, Salzwedel DM, Cogo E, Foerster V, Lefebvre C. PRESS Peer Review of Electronic Search Strategies: 2015 Guideline Statement. *J Clin Epidemiol.* 2016;75:40-6.
19. Bramer WM, Giustini D, de Jonge GB, Holland L, Bekhuis T. De-duplication of database search results for systematic reviews in EndNote. *J Med Libr Assoc.* 2016;104(3):240-3.

Appendix I: Search strategy

Block 1: SGA newborn	MeSH (PubMed and Cochrane Library): <ul style="list-style-type: none"> - Infant, Low Birth Weight Emtree (Embase): <ul style="list-style-type: none"> - low birth weight (Explode turned ON) Keywords: <ul style="list-style-type: none"> - Small for gestational age - Small for date - SGA - Low: infant, infants, newborn*, gestational age or neonate* - Small: infant, infants, newborn*, gestational age or neonate*
Block 2: Body-proportionality	MeSH (PubMed and Cochrane Library): <ul style="list-style-type: none"> - Body Height (Explode turned OFF) - Birth Weight (Explode turned OFF) Emtree (Embase): <ul style="list-style-type: none"> - body height - birth weight Keywords: <ul style="list-style-type: none"> - asymmetric SGA - asymmetric small - symmetric SGA - symmetric small - proportionate SGA - proportionate small - disproportionate SGA - disproportionate small - body-proportion* - head circumference* - ponderal - percentile - body length

Table 1: The two search blocks with the list of subject headings and keywords used in the search strategy.

(Infant, Low Birth Weight[MeSH] OR Small for gestational age[TIAB] OR Small for date[TIAB] OR SGA[TIAB] OR ((low[TIAB] OR small[TIAB]) AND (infant[TIAB] OR infants[TIAB] OR newborn*[TIAB] OR Gestational Age[TIAB] OR Neonat*[TIAB])))
AND
(Body Height[MeSH:NoExp] OR Birth Weight[MeSH:NoExp] OR asymmetric SGA[TIAB] OR asymmetric small[TIAB] OR symmetric SGA[TIAB] OR symmetric small[TIAB] OR proportionate SGA[TIAB] OR proportionate small[TIAB] OR disproportionate SGA[TIAB] OR disproportionate small[TIAB] OR body-proportion*[TIAB] OR head circumference*[TIAB] OR ponderal[TIAB] OR percentile[TIAB] OR body length[TIAB])

Table 2: The search strategy for PubMed.

Appendix II: Data presentation

Author	Title	Year	Population	No. of patients	Setting	Definition of SGA	Definition of proportionality	Outcome related to body-proportionality

Table 3: A work in progress example of the overview table in which the results of the data extraction will be presented.