Meaf6 is differentially expressed in the brains of patients with schizophrenia.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY USA

Visual and auditory hallucinations are a cardinal feature of psychotic disorders¹. We mined published and public microarray datasets^{2,3} to discover differentially expressed genes in schizophrenia and schizoaffective disorder. We found significant differential expression of Meaf6 in neurons of the dorsolateral pre-frontal cortex from patients with schizophrenia and schizoaffective disorder.

Keywords: schizophrenia, schizoaffective disorder, psychotic disorders, psychosis, systems biology of schizophrenia, Meaf6.

Psychiatric diseases schizophrenia and schizoaffective disorder have in common the existence of psychotic symptoms^{1,4,5}, like visual and auditory hallucinations, that can plague affected patients. Understanding the transcriptional profiles of cells and tissues from the brain of patients diagnosed with schizophrenia or schizoaffective disorder has the potential to provide insights into molecular mechanisms underlying disease initiation, progression and associated symptoms like psychosis. We mined published and public microarray data^{2,3} from layer 3 and layer 5 pyramidal neurons of the dorsolateral pre-frontal cortex from patients with schizophrenia and schizoaffective disorder and from the layer 3 parvalbumin-positive neurons of the dorsolateral prefrontal cortex of patients with schizoaffective disorder to discover genes whose expression was most significantly different in the brains of patients with psychosis at the systems-level. Meaf6 was among the genes most differentially expressed in two cell types of the brain in patients with psychotic disorders.

Methods

We used microarray datasets GSE93577² and GSE87610³ for this differential gene expression analysis in conjunction with GEO2R. GSE93577 was generated using Affymetrix Human Genome U219 Array technology from layer 3 (L3) parvalbumin (PV) positive neurons of the dorsolateral prefrontal cortex, and we used $n=34$ control subjects each with paired replicates ($n=68$ total samples) and $n=35$ patients with schizophrenia or schizoaffective disorder with paired replicates ($n=70$ total samples) from this dataset. These patient diagnoses in this dataset

1 included schizoaffective disorder, paranoid schizophrenia, undifferentiated schizophrenia,
2 chronic paranoid schizophrenia, chronic disorganized schizophrenia, chronic undifferentiated
3 schizophrenia, and disorganized schizophrenia. GSE87610 was generated using Affymetrix
4 Human Genome U219 with $n=72$ for control subjects and $n=65$ for patients with schizophrenia,
5 using L3 and L5 pyramidal neurons from the dorsolateral prefrontal cortex.
6

7
8 The Benjamini and Hochberg method of p-value adjustment was used for ranking of
9 differential expression but raw p-values were used for assessment of statistical significance of
10 global differential expression. Log-transformation of data was auto-detected, and the NCBI
11 generated category of platform annotation was used. A statistical test was performed to evaluate
12 whether Meaf6 expression was significantly different when comparing neurons from the DLPFC
13 from control subjects and or from patients with schizophrenia or schizophrenia and
14 schizoaffective disorder, respectively, using a two-tailed, unpaired t-test with Welch's correction.
15 We used PRISM for all statistical analyses (Version 8.4.0)(455).
16
17
18

19 20 **Results**

21 We mined published and public microarray data^{2,3} to identify in an unbiased fashion and
22 at the global level genes most differentially expressed in the brains of patients with schizophrenia
23 and schizoaffective disorder.
24

25
26 Meaf6 is differentially expressed in layer 3 parvalbumin-positive neurons of the dorsolateral pre-
27 frontal cortex in patients with schizophrenia and schizoaffective disorder.
28

We observed differential expression of Meaf6 in the layer 3 (L3) PV+ neurons of the dorsolateral prefrontal cortex (DLPFC) in patients with schizophrenia and schizoaffective disorder². When sorting each of the transcripts whose expression was measured by microarray based on significance of change in expression between diseased and control patients, in L3 PV+ DLPFC neurons from patients with schizophrenia, Meaf6 ranked 29 out of 49386 total transcripts. Differential expression of Meaf6 was statistically significant (Table 1; $p=9.59E-06$).

Meaf6 is differentially expressed in L3 and L5 pyramidal neurons of the dorsolateral pre-frontal cortex in patients with schizophrenia.

We also observed differential expression of Meaf6 in the layer 3 and layer 5 (L5) pyramidal neurons of the DLPFC in patients with schizophrenia. When sorting each of the transcripts whose expression was measured by microarray based on significance of change in expression between patients and control subjects, in L3 and L5 pyramidal neurons of the DLPFC from patients with schizophrenia, Meaf6 ranked 3869 out of 49386 total transcripts (Table 3), equating to differential expression greater than 92.2% of all transcripts whose expression was detected and measured. Differential expression of Meaf6 in L3 and L5 pyramidal neurons of the DLPFC in patients with schizophrenia was statistically significant (Table 2; $p=1.89E-02$).

Meaf6 expression is significantly lower in layer 3 parvalbumin-positive neurons of the dorsolateral pre-frontal cortex in patients with schizophrenia and schizoaffective disorder.

We obtained exact mRNA expression levels for the differentially expressed Meaf6

transcript, from control subjects and from patients with schizophrenia and schizoaffective disorder, to directly compare Meaf6 expression levels in L3 PV+ DLPFC neurons. We also performed a statistical test to determine if this difference in Meaf6 expression was statistically significant. Meaf6 was expressed at lower levels in the L3 PV+ DLPFC neurons in patients with schizophrenia and schizoaffective disorder, and this difference was statistically significant (Figure 1; $p=0.0001$). We calculated a mean fold change of 0.957 ± 0.058 in Meaf6 expression when comparing L3 PV+ DLPFC neurons from patients with schizophrenia and schizoaffective disorder to that of control subjects (Table 1).

Thus, we found significantly differential expression of Meaf6 in two types of neuronal cell types in patients with schizophrenia; Meaf6 expression was significantly lower in L3 PV+ neurons of the dorsolateral prefrontal cortex in patients with schizophrenia and schizoaffective disorder.

Discussion

There are five major families of histone acetyltransferase (HATs, or KATs histone lysine acetyltransferases), one of which is the Myst family of KATs including Myst1 (Kat8), Myst2 (Kat7), Myst3 (Kat6a), Myst4 (Kat6b), and Tip60 (Kat5)⁶. The MYST Esa1-associated factor (Meaf) Meaf6 is one subunit of the Myst2/Kat7 histone acetyltransferase complex. The function of Meaf6 is poorly characterized. The yeast homolog of Meaf6, Eaf6 is part of the yeast NuA4 HAT complex, and of 12 subunits, Eaf6 is one of 7 not required for cell viability⁷. In embryonic

1 stem cells, Meaf6 can interact with both Myst2/Kat7 and Tip60/Kat5 to form part of each
2 respective HAT complex (Kat7, Brd1, Jade1, Ing4/Ing5 or Brd8, Ruvbl1, Ruvbl2, Actl6a, Epc2,
3 Vsp72, Morfl1 and Ing3, respectively)^{8,9}.

5 In endometrial stromal sarcoma (ESS), chromosomal rearrangements of 6p21,
6 specifically affecting the Phf1 gene; one fusion transcript found in ESS wherein exon 5 of
7 Meaf6, located on 1p34, is fused to exon 2 of Phf1 resulting from a t(1;6) (p34;p21)
8 chromosomal translocation. The fusion protein produced by this transcript includes the NuA4
9 histone acetyltransferase domain of Meaf6 in conjunction with the tudor, PHD zinc finger and
10 MTF2 domains of PHF1^{10,11}.

13 In the male Sprague-Dawley rat, seizures resulting from electroconvulsive therapy result
14 in significant increase in mRNA expression of Brd1, a component of the Myst2/Kat7 histone
15 acetyltransferase complex, in the hippocampus and frontal cortex¹². There is little information
16 regarding involvement of Meaf6 in human diseases of the brain but mutation of Brpf1 results in
17 syndromic intellectual disability¹³; Brpf1 can interact with Kat7 (Myst2), as well as Kat6a
18 (Myst3) and Kat6b (Myst4). Moreover, a number of diseases including features of intellectual or
19 learning disabilities and/or abnormalities of the facial structure have been found to be the result
20 of *de novo* mutations in Myst family histone acetyltransferases, including in Ohdo Syndrome,
21 Genitopatellar syndrome, and in a patient with Noonan-like syndrome¹⁴⁻¹⁸. Thus, mutation of
22 epigenetic machinery belonging to the MYST family of histone lysine acetyltransferases can
23 result in human diseases of the brain but little information exists describing specific requirement
24
25
26
27
28

1 for or involvement of Meaf6 in human diseases outside of cancer.

2 We found differential expression of Meaf6 in neurons from patients with schizophrenia in
3
4 two published or public microarray datasets. Differential expression of Meaf6 should be
5 validated in separate and larger schizophrenia patient datasets. Meaf6 may be of relevance to the
6 biology of schizophrenia.
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

References

1. Andreasen, N.C., Nopoulos, P., Schultz, S., Miller, D., Gupta, S., Swayze, V. and Flaum, M., 1994. Positive and negative symptoms of schizophrenia: past, present, and future. *Acta Psychiatrica Scandinavica*, 90, pp.51-59.
2. GSE93577. University of Pittsburgh, Department of Psychiatry and Neuroscience. Pittsburgh, PA. Enwright JF, Huo Z, Arion D, Corradi JP, Tseng G, Lewis DA. <https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE93577>.
3. Arion, D., Huo, Z., Enwright, J.F., Corradi, J.P., Tseng, G. and Lewis, D.A., 2017. Transcriptome alterations in prefrontal pyramidal cells distinguish schizophrenia from bipolar and major depressive disorders. *Biological psychiatry*, 82(8), pp.594-600.
4. Cannon, T.D., 2015. How schizophrenia develops: cognitive and brain mechanisms underlying onset of psychosis. *Trends in cognitive sciences*, 19(12), pp.744-756.
5. Arciniegas, D.B., 2015. Psychosis. *Continuum: Lifelong Learning in Neurology*, 21(3 Behavioral Neurology and Neuropsychiatry), p.715.
6. Sapountzi, V. and Côté, J., 2011. MYST-family histone acetyltransferases: beyond chromatin. *Cellular and Molecular Life Sciences*, 68(7), pp.1147-1156.
7. Mitchell, L., Lambert, J.P., Gerdes, M., Al-Madhoun, A.S., Skerjanc, I.S., Figeys, D. and Baetz, K., 2008. Functional dissection of the NuA4 histone acetyltransferase reveals its role as a genetic hub and that Eaf1 is essential for complex integrity. *Molecular and cellular biology*, 28(7), pp.2244-2256.
8. Matsuura, K., Tani, N., Usuki, S., Torikai-Nishikawa, S., Okano, M. and Niwa, H., 2020. MEAF6 is essential for cell proliferation and plays a role in the assembly of KAT7 complexes. *Experimental Cell Research*, p.112279.
9. Pardo, M., Yu, L., Shen, S., Tate, P., Bode, D., Letney, B.L., Quelle, D.E., Skarnes, W. and Choudhary, J.S., 2017. Myst2/Kat7 histone acetyltransferase interaction proteomics reveals tumour-suppressor Niam as a novel binding partner in embryonic stem cells. *Scientific reports*, 7(1), pp.1-14.
10. Panagopoulos, I., Micci, F., Thorsen, J., Gorunova, L., Eibak, A.M., Bjerkehagen, B., Davidson, B. and Heim, S., 2012. Novel fusion of MYST/Es1-associated factor 6 and PHF1 in endometrial stromal sarcoma. *PLoS One*, 7(6), p.e39354.

11. Micci, F., Gorunova, L., Gatus, S., Matias-Guiu, X., Davidson, B., Heim, S. and Panagopoulos, I., 2014. MEAF6/PHF1 is a recurrent gene fusion in endometrial stromal sarcoma. *Cancer Letters*, 347(1), pp.75-78.
12. Fryland, T., Elfving, B., Christensen, J.H., Mors, O., Wegener, G. and Børghlum, A.D., 2012. Electroconvulsive seizures regulates the Brd1 gene in the frontal cortex and hippocampus of the adult rat. *Neuroscience letters*, 516(1), pp.110-113.
13. Yan, K., Rousseau, J., Littlejohn, R.O., Kiss, C., Lehman, A., Rosenfeld, J.A., Stumpel, C.T., Stegmann, A.P., Robak, L., Scaglia, F. and Nguyen, T.T.M., 2017. Mutations in the chromatin regulator gene BRPF1 cause syndromic intellectual disability and deficient histone acetylation. *The American Journal of Human Genetics*, 100(1), pp.91-104.
14. Kraft, M., Cirstea, I.C., Voss, A.K., Thomas, T., Goehring, I., Sheikh, B.N., Gordon, L., Scott, H., Smyth, G.K., Ahmadian, M.R. and Trautmann, U., 2011. Disruption of the histone acetyltransferase MYST4 leads to a Noonan syndrome-like phenotype and hyperactivated MAPK signaling in humans and mice. *The Journal of clinical investigation*, 121(9).
15. Clayton-Smith, J., O'Sullivan, J., Daly, S., Bhaskar, S., Day, R., Anderson, B., Voss, A.K., Thomas, T., Biesecker, L.G., Smith, P. and Fryer, A., 2011. Whole-exome-sequencing identifies mutations in histone acetyltransferase gene KAT6B in individuals with the Say-Barber-Biesecker variant of Ohdo syndrome. *The American Journal of Human Genetics*, 89(5), pp.675-681.
16. Simpson, M.A., Deshpande, C., Dafou, D., Vissers, L.E., Woollard, W.J., Holder, S.E., Gillissen-Kaesbach, G., Derks, R., White, S.M., Cohen-Snuijf, R. and Kant, S.G., 2012. De novo mutations of the gene encoding the histone acetyltransferase KAT6B cause Genitopatellar syndrome. *The American Journal of Human Genetics*, 90(2), pp.290-294.
17. Arboleda, V.A., Lee, H., Dorrani, N., Zadeh, N., Willis, M., Macmurdo, C.F., Manning, M.A., Kwan, A., Hudgins, L., Barthelemy, F. and Miceli, M.C., 2015. De novo nonsense mutations in KAT6A, a lysine acetyl-transferase gene, cause a syndrome including microcephaly and global developmental delay. *The American Journal of Human Genetics*, 96(3), pp.498-506.
18. Tham, E., Lindstrand, A., Santani, A., Malmgren, H., Nesbitt, A., Dubbs, H.A., Zackai, E.H., Parker, M.J., Millan, F., Rosenbaum, K. and Wilson, G.N., 2015. Dominant mutations in KAT6A cause intellectual disability with recognizable syndromic features. *The American Journal of Human Genetics*, 96(3), pp.507-513.

Rank	ID	p-value	t	B	FC	Gene
29	11756418_a_at	9.59E-06	-4.60	3.224436	0.957 ± 0.058	MEAF6

Table 1: Meaf6 is differentially expressed in the layer 3 PV+ neurons of the dorsolateral pre-frontal cortex in patients with schizophrenia and schizoaffective disorder.

The rank of differential expression, the probe ID, the *p*-value with respect to global differential expression, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the two groups compared, the fold change of Meaf6 in L3 PV+ DLPFC neurons of patients with schizophrenia and schizoaffective disorder when compared to deep pyramidal neurons from control subjects, the gene and gene name are listed in this chart.

Rank	ID	p-value	t	B	Gene
3869	11756418_a_at	1.89E-02	2.3748399	-3.38351	MEAF6

Table 2: Meaf6 is differentially expressed in the layer 3 and layer 5 pyramidal neurons of the dorsolateral pre-frontal cortex of patients with schizophrenia.

The rank of differential expression, the probe ID, the *p*-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, the gene and gene name are listed in this chart.

Figure 1: Meaf6 is expressed at significantly lower levels in the layer 3 PV+ neurons of the dorsolateral pre-frontal cortex of patients with schizophrenia and schizoaffective disorder.

The mRNA expression of Meaf6 in L3 PV+ neurons of the DLPFC from control subjects (left) and from patients with schizophrenia and schizoaffective disorder (right) is represented with mean mRNA expression level marked and the result of a statistical test evaluating the significance of difference in Meaf6 expression between control L3 PV+ DLPFC neurons and L3 PV+ DLPFC neurons from patients with schizophrenia and schizoaffective disorder, a *p*-value, listed above.