Platelet-derived growth factors α and β are differentially expressed in the primary tumors of breast cancer patients treated with trastuzumab.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY USA

Trastuzumab, a monoclonal antibody targeted against the human epidermal growth factor receptor 2 (HER2) is utilized for the treatment of human breast cancer^{1,2}, but a complete understanding of how tumor signal transduction is modulated by trastuzumab treatment is lacking. By mining published and public microarray and gene expression data^{3,4} from the primary tumors of patients treated with trastuzumab, we found that platelet-derived growth factors α and β , PDGF- α and PDGF- β , are among the genes most differentially expressed in the primary tumors of patients treated with trastuzumab, and expressed at significantly higher levels in the tumors of patients treated with trastuzumab. Thus, the use of trastuzumab in patients with breast cancer is associated with increased expression growth factors important for development of the neural crest and neural tube^{5,6}.

Keywords: breast cancer, trastuzumab, PDGF- α , PDGF- β , systems biology of breast cancer, targeted therapeutics in breast cancer.

The mechanism of action by which trastuzumab exerts its therapeutic function in human breast cancer is said to be binding of the F_{ab} domain of trastuzumab to the extracellular region of the human epidermal growth factor receptor 2 (HER2)², possibly resulting in decreased cellular proliferation⁷. Transcriptional modulation resulting from trastuzumab binding events in the primary tumors of treated patients is less well understood. We mined published and public microarray and gene expression data^{3,4} generated using primary tissues from patients treated with trastuzumab to compare tumor global gene expression with that of patients not treated with trastuzumab. Using this unbiased, systems level methodology, we identified differential expression and up-regulation of PDGF- α and PDGF- β in primary tumors of the breast in patients treated with trastuzumab.

Methods

We used datasets GSE38057³ and GSE130788⁴ for this differential gene expression analysis of the transcriptional response to trastuzumab treatment in patients with breast cancer. GSE38057 was generated using DASL Human Cancer Panel by Probe technology with $n=11$ primary breast tumors from patients treated with induction chemotherapy and/or hormone therapy without trastuzumab and $n=36$ primary breast tumors from patients treated with induction chemotherapy and/or hormone therapy in addition to trastuzumab; analysis was performed using platform GPL5826. GSE130788 was generated using Agilent-014850 Whole Human Genome Microarray 4x44K G4112F technology with $n=11$ primary breast tumors from patients treated with trastuzumab and $n=36$ primary tumors from patients treated with trastuzumab, analysis performed using platform GPL6480. The Benjamini and Hochberg method of p -value adjustment was used for ranking of differential expression but raw p -values were used to assess statistical significance of global or multiplexed differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. A statistical test was performed to evaluate whether PDGF- α and PDGF- β gene expression was significantly between primary breast tumors of patients receiving trastuzumab using a two-tailed, unpaired t-test with Welch's correction. We used PRISM for all statistical analyses of differential gene expression in human breast cancer (Version 8.4.0)(455).

Results

We performed comparative transcriptome analysis of primary tumors from patients treated or not with trastuzumab (Herceptin), using published multiplexed gene expression data³ and a microarray dataset⁴, in order to identify the most significant transcriptional changes in the primary tumors of breast cancer patients following treatment with trastuzumab.

PDGFA is differentially expressed in the primary tumors of patients treated with trastuzumab.

We found that the PDGF- α , encoded by PDGFA, was among the genes whose expression was most significantly different in the primary tumors of patients with breast cancer treated with trastuzumab³. When sorting each of the genes expressed in the primary tumors of breast cancer patients treated with trastuzumab based on significance of difference in expression as compared to the primary tumors of breast cancer patients not treated with trastuzumab, PDGFA ranked 16 out of 1506 total transcripts, equating to 98.9% statistical significance (Table 1). Differential expression of PDGFA in the primary tumors of breast cancer patients treated with trastuzumab was statistically significant (Table 1; $p=0.02462$).

We queried a second dataset⁶ to validate our findings of PDGFA differential expression upon trastuzumab treatment. In a second microarray dataset analyzing gene expression of patients treated with

trastuzumab, we again found that PDGFA was among the genes whose expression was most significantly different transcriptome-wide when comparing the global gene expression profiles of tumors from patients treated with trastuzumab as compared to those not treated with trastuzumab. When sorting each of the genes expressed in the tumors of patients treated with trastuzumab based on significance of change as compared to the tumors of patients not treated with trastuzumab, PDGFA ranked 21346 out of 41000 total transcripts, equating to 47.9% differential expression. Though differential expression of PDGFA was less significant than greater than half of the transcriptome, differential expression of PDGFA in the primary tumors of patients treated with trastuzumab was statistically significant (Table 2; $p=5.8E-03$).

PDGFA is expressed at higher levels in the tumors of patients treated with trastuzumab.

We obtained exact mRNA expression levels for PDGFA from the tumors of patients treated with trastuzumab and from the tumors of patients not receiving trastuzumab to determine magnitude and direction of PDGFA change in the tumors of patients treated with trastuzumab. PDGFA expression was higher in the tumors of patients treated with trastuzumab as compared to the tumors of patients not treated with trastuzumab (Figure 1). Markedly enhanced expression of PDGFA was not deemed statistically significant (Figure 1: $p=0.2340$). We calculated a mean fold change of 1.46 ± 0.76 when comparing the expression of PDGFA in the tumors of patients treated with trastuzumab as compared to the tumors of patients not treated with trastuzumab.

PDGFB is differentially expressed in the primary tumors of patients treated with trastuzumab.

We also found that the PDGF- β , encoded by PDGFB, was among the genes whose expression was most significantly different in the primary tumors of patients with breast cancer treated with trastuzumab³. When sorting each of the genes expressed in the primary tumors of breast cancer patients treated with trastuzumab based on significance of difference in expression as compared to the primary tumors of breast cancer patients not treated with trastuzumab, PDGFB ranked 124 out of 1506 total transcripts, equating to 91.8% statistical significance (Table 3). Differential expression of PDGFB in the primary tumors of breast cancer patients treated with trastuzumab was not deemed statistically significant (Table 3; $p=0.095874$).

We queried a second dataset⁶ to validate our findings of PDGFB differential expression upon trastuzumab treatment. In a second microarray dataset analyzing gene expression of patients treated with trastuzumab, we again found that PDGFB was among the genes whose expression was most significantly different transcriptome-wide when comparing the global gene expression profiles of tumors from patients treated with trastuzumab as compared to those not treated with trastuzumab. When sorting each of the genes expressed in the tumors of patients treated with trastuzumab based on significance of change as compared to the tumors of patients not treated with trastuzumab, PDGFB ranked 19859 out of 41000 total transcripts, equating to 51.6% differential expression. Differential expression of PDGFB in the primary tumors of patients treated with trastuzumab was statistically significant (Table 4; $p=2.45E-03$).

PDGFB is expressed at higher levels in the tumors of patients treated with trastuzumab.

We obtained exact mRNA expression levels for PDGFB from the tumors of patients treated with trastuzumab and from the tumors of patients not receiving trastuzumab to determine magnitude and direction of PDGFB change in the tumors of patients treated with trastuzumab. PDGFB expression was higher in the tumors of patients treated with trastuzumab as compared to the tumors of patients not treated

1 with trastuzumab (Figure 2). Enhanced expression of PDGFB was statistically significant (Figure 2:
2 $p=0.0212$). We calculated a mean fold change of 1.26 ± 0.48 when comparing the expression of PDGFB
3 in the tumors of patients treated with trastuzumab as compared to the tumors of patients not treated with
4 trastuzumab.

5 Thus, we found that the PDGF- α and PDGF- β , encoded by PDGFA and PDGFB, was among the
6 genes most differentially expressed in the primary tumors of patients with breast cancer and that
7 expression of both PDGF- α and PDGF- β were higher in the primary tumors of patients treated with
8 trastuzumab as compared to those not treated.

9 Discussion

10 We provided evidence here that PDGF- α and PDGF- β , encoded by PDGFA and PDGFB were
11 expressed at higher levels in the tumors of patients treated with trastuzumab and among the genes most
12 differentially expressed in the primary tumors of patients treated with trastuzumab. Thus, administration
13 of trastuzumab in patients with breast cancer results in increased expression of growth factors that are
14 essential for development of the mammalian nervous system^{6,7} - in the neural crest and the neural tube of
15 human embryonic stem cells and whose transgenic expression results in the development of glioma-like
16 lesions in the brain¹².

References

1. Slamon, D., Eiermann, W., Robert, N., Pienkowski, T., Martin, M., Press, M., Mackey, J., Glaspy, J., Chan, A., Pawlicki, M. and Pinter, T., 2011. Adjuvant trastuzumab in HER2-positive breast cancer. *New England Journal of Medicine*, 365(14), pp.1273-1283.
2. Hudis, C.A., 2007. Trastuzumab—mechanism of action and use in clinical practice. *New England Journal of Medicine*, 357(1), pp.39-51.
3. Duchnowska, R., Jassem, J., Goswami, C.P., Dundar, M., Gökmen-Polar, Y., Li, L., Woditschka, S., Biernat, W., Sosińska-Mielcarek, K., Czartoryska-Arłukowicz, B. and Radecka, B., 2015. Predicting early brain metastases based on clinicopathological factors and gene expression analysis in advanced HER2-positive breast cancer patients. *Journal of Neuro-oncology*, 122(1), pp.205-216.
4. GSE130788. A Phase II Randomized Trial of Neoadjuvant Trastuzumab or Lapatinib or the Combination of Trastuzumab and Lapatinib, Followed by Docetaxel and Carboplatin with Trastuzumab and/or Lapatinib in Patients with HER2+ Breast Cancer. [https:// www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE130788](https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE130788). Hsiaowang Chen. UCLA, Department of Medicine. Santa Monica, CA.
5. Soriano, P., 1997. The PDGF alpha receptor is required for neural crest cell development and for normal patterning of the somites. *Development*, 124(14), pp.2691-2700.
6. Pringle, N.P. and Richardson, W.D., 1993. A singularity of PDGF alpha-receptor expression in the dorsoventral axis of the neural tube may define the origin of the oligodendrocyte lineage. *Development*, 117(2), pp.525-533.
7. Hall, A., Giese, N.A. and Richardson, W.D., 1996. Spinal cord oligodendrocytes develop from ventrally derived progenitor cells that express PDGF alpha-receptors. *Development*, 122(12), pp. 4085-4094.
8. Mudhar, H.S., Pollock, R.A., Wang, C., Stiles, C.D. and Richardson, W.D., 1993. PDGF and its receptors in the developing rodent retina and optic nerve. *Development*, 118(2), pp.539-552.
9. Smits, A., Kato, M., Westermarck, B., Nister, M., Heldin, C.H. and Funa, K., 1991. Neurotrophic activity of platelet-derived growth factor (PDGF): Rat neuronal cells possess functional PDGF beta-type receptors and respond to PDGF. *Proceedings of the National Academy of Sciences*, 88(18), pp. 8159-8163.
10. Yeh, H.J., Silos-Santiago, I., Wang, Y.X., George, R.J., Snider, W.D. and Deuel, T.F., 1993. Developmental expression of the platelet-derived growth factor alpha-receptor gene in mammalian central nervous system. *Proceedings of the National Academy of Sciences*, 90(5), pp.1952-1956.
11. Fruttiger, M., Calver, A.R., Krüger, W.H., Mudhar, H.S., Michalovich, D., Takakura, N., Nishikawa, S.I. and Richardson, W.D., 1996. PDGF mediates a neuron–astrocyte interaction in the developing retina. *Neuron*, 17(6), pp.1117-1131.
12. Nazarenko, I., Hedrén, A., Sjödin, H., Orrego, A., Andrae, J., Afink, G.B., Nistér, M. and Lindström, M.S., 2011. Brain abnormalities and glioma-like lesions in mice overexpressing the long isoform of PDGF-A in astrocytic cells. *PloS one*, 6(4), p.e18303.

Rank	ID	p-value	t	B	FC	Gene
16	GI_15208655-A-3	0.02462	2.329201	-4.26	1.46 ± 0.76	PDGFA

Table 1: PDGFA is differentially expressed in the primary tumors of breast cancer patients treated with trastuzumab.

The rank of differential expression, probe ID, p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, fold change of expression in PDGFA between the tumors of patients treated with trastuzumab and the tumors of patients not treated with trastuzumab, and gene are listed in this chart.

Rank	ID	p-value	t	B	Gene	Gene name
21346	A_23_P113701	5.8E-03	2.788779	-3.788313	PDGFA	platelet derived growth factor subunit A

Table 2: PDGFA is differentially expressed in the primary tumors of breast cancer patients treated with trastuzumab.

The rank of differential expression, probe ID, p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, fold change of expression in PDGFA between the tumors of patients treated with trastuzumab and the tumors of patients not treated with trastuzumab, and gene are listed in this chart.

Rank	ID	p-value	t	B	FC	Gene
124	GI_15451785-A-6	0.095874	1.69815	-4.23	1.26 ± 0.48	PDGFB

Table 3: PDGFB is differentially expressed in the primary tumors of breast cancer patients treated with trastuzumab.

The rank of differential expression, probe ID, p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, fold change of expression in PDGFB between the tumors of patients treated with trastuzumab and the tumors of patients not treated with trastuzumab, and gene are listed in this chart.

Rank	ID	p-value	t	B	Gene	Gene name
19859	A_24_P339944	2.45E-03	3.069171	-2.999021	PDGFB	platelet derived growth factor subunit B

Table 4: PDGFB is differentially expressed in the primary tumors of breast cancer patients treated with trastuzumab.

The rank of differential expression, probe ID, p-value with respect to global differential expression, t, a moderated t-statistic, B, the log-odds of differential expression between the two groups compared, fold change of expression in PDGFB between the tumors of patients treated with trastuzumab and the tumors of patients not treated with trastuzumab, and gene are listed in this chart.

Figure 1: PDGFA is expressed at higher levels in the primary tumors of patients treated with trastuzumab as compared to those not treated with trastuzumab.

The mRNA expression level of PDGFA is graphically depicted in this chart in the primary tumors of breast cancer patients not treated with trastuzumab (left) and in the primary tumors of breast cancer patients treated with trastuzumab (right), with mean mRNA expression level marked and the result of a statistical test evaluating significance of difference between PDGFA expression in the primary tumors of patients treated with trastuzumab or not, a *p*-value, listed above.

Figure 2: PDGFB is expressed at higher levels in the primary tumors of patients treated with trastuzumab as compared to those not treated with trastuzumab.

The mRNA expression level of PDGFB is graphically depicted in this chart in the primary tumors of breast cancer patients not treated with trastuzumab (left) and in the primary tumors of breast cancer patients treated with trastuzumab (right), with mean mRNA expression level marked and the result of a statistical test evaluating significance of difference between PDGFB expression in the primary tumors of patients treated with trastuzumab or not, a *p*-value, listed above.