GENERAL SOLUTION OF Nth-ORDER LINEAR ORDINARY DIFFERENTIAL EQUATION

RAJNISH KUMAR JHA

ABSTRACT. In this paper we present a solution expression for the general n^{th} -order linear ordinary differential equation as our main result which involves the use of Integrating Factors where the Integrating Factors are determined using a set of equations such that when this set of equations can be solved, the solution of the concerned differential equation can be determined completely. In this regard we also present result for a special case corresponding to the main result where the solution of the general n^{th} -order linear ordinary differential equation can be determined completely when $n - 1$ out of n complementary solutions are known.

1. INTRODUCTION

The general n^{th} -order linear ordinary differential equation is given as

$$(1) \quad y_{(n)} + P_{n1}(x)y_{(n-1)} + \cdots + P_{nn-1}(x)y_{(1)} + P_{nn}(x)y = R(x),$$

where $y_{(k)}$ represents the k^{th} derivative of the dependent variable - y with respect to x , the independent variable, such that

$$(2) \quad y_{(k)} = \frac{d^k y}{dx^k},$$

$P_{ni}(x)$, $i = 1, 2, \dots, n$ and $R(x)$ are the functions of the independent variable - x .

When $R(x) = 0$ then Eqn. (1) is called homogeneous n^{th} -order linear ordinary differential equation [1] otherwise when $R(x) \neq 0$ then it is called non-homogeneous n^{th} -order linear ordinary differential equation [2].

When the coefficients $P_{ni}(x)$, $i = 1, 2, \dots, n$ are constant such that $P_{ni} = p_{ni}$ then Eqn. (1) is referred to as n^{th} -order linear ordinary differential equation with constant coefficients [3].

The solution of the differential equation given by Eqn. (1) is generally difficult to obtain. However, if the coefficients of this differential equation are constant then the differential equation can be solved completely.

In this paper we present an expression for the solution of the general n^{th} -order linear ordinary differential equation such that the expression involves n successive integrals using n Integrating Factors where these

Integrating Factors are themselves determined using a set of equations (See Theorem-1). Although this process may be tedious, the form in which the solution expression is obtained can be helpful in designing special cases for solving such type of equations. One such case is described in this paper where if $n - 1$ out of n complementary solutions of the concerned differential equation are known then its solution can be determined completely (See Corollary-1).

We present our main result in form of a theorem described in Section-2 and the other result, which gives the complete solution of the concerned differential equation when $n - 1$ out of n complementary solutions are known, is described in Section-3.

2. MAIN RESULT

In this section we state the main result as Theorem-1 which is given along with the proof here as follows

Theorem 1. *The solution of the general n^{th} -order linear ordinary differential equation as given in Eqn. (1) is given by*

$$(3) \quad y = \frac{1}{g_n(x)} \int \frac{g_n(x)}{g_{n-1}(x)} \int \dots \int \frac{g_2(x)}{g_1(x)} \int g_1(x) R(x) \underbrace{dx dx \dots dx}_{n \text{ times}} \\ + \frac{C_1}{g_n(x)} \int \frac{g_n(x)}{g_{n-1}(x)} \int \dots \int \frac{g_2(x)}{g_1(x)} \underbrace{dx dx \dots dx}_{(n-1) \text{ times}} \\ + \dots + \frac{C_{n-1}}{g_n(x)} \int \frac{g_n(x)}{g_{n-1}(x)} dx + \frac{C_n}{g_n(x)},$$

where $g_{n-i+1}(x)$ represents $(n - i + 1)^{th}$ Integrating Factor such that

$$(4) \quad g_{n-i+1}(x) = e^{\int a_{i1}(x) dx}, \quad i = 1, 2, \dots, n,$$

and each $a_{i1}(x)$ can be determined from the set of equations given as

$$(5) \quad P_{ij}(x) = a_{ij}(x) + P_{i-1j}(x) \{1 \leq j \leq i - 1; 1 \leq i \leq n\},$$

$$(6) \quad a_{11}(x) = P_{11}(x),$$

$$(7) \quad P_{i-1j}(x) a_{i1}(x) + P'_{i-1j}(x) = a_{ij+1}(x) \{1 \leq j \leq i - 1; 1 \leq i \leq n\},$$

$$(8) \quad P_{i-1i-1}(x) a_{i1}(x) + P'_{i-1i-1}(x) = P_{ii}(x) \{1 < i \leq n\}$$

and $i, j, n \in \mathbb{N}$.

Proof. We shall prove this theorem by Mathematical Induction and in this regard we consider a statement $S(n)$ corresponding to the statement of Theorem-1 given as follows

S(n): The solution of the general n^{th} -order linear ordinary differential equation,

$$(9) \quad y_{(n)} + P_{n1}(x)y_{(n-1)} + \dots + P_{nn-1}(x)y_{(1)} + P_{nn}(x)y = R(x),$$

is given by

$$\begin{aligned}
(10) \quad y &= \frac{1}{g_n(x)} \int \frac{g_n(x)}{g_{n-1}(x)} \int \dots \int \frac{g_2(x)}{g_1(x)} \int g_1(x) R(x) \underbrace{dx dx \dots dx}_{n \text{ times}} \\
&+ \frac{C_1}{g_n(x)} \int \frac{g_n(x)}{g_{n-1}(x)} \int \dots \int \frac{g_2(x)}{g_1(x)} \underbrace{dx dx \dots dx}_{(n-1) \text{ times}} \\
&+ \dots + \frac{C_{n-1}}{g_n(x)} \int \frac{g_n(x)}{g_{n-1}(x)} dx + \frac{C_n}{g_n(x)},
\end{aligned}$$

where

$$(11) \quad g_{n-i+1}(x) = e^{\int a_{i1}(x) dx}, \quad i = 1, 2, \dots, n$$

and each $a_{i1}(x)$ can be determined from the set of equations given as

$$(12) \quad P_{ij}(x) = a_{ij}(x) + P_{i-1j}(x) \{1 \leq j \leq i-1; 1 \leq i \leq n\},$$

$$(13) \quad a_{11}(x) = P_{11}(x),$$

$$(14) \quad P_{i-1j}(x) a_{i1}(x) + P'_{i-1j}(x) = a_{ij+1}(x) \{1 \leq j \leq i-1; 1 \leq i \leq n\},$$

$$(15) \quad P_{i-1i-1}(x) a_{i1}(x) + P'_{i-1i-1}(x) = P_{ii}(x) \{1 < i \leq n\}$$

and $i, j, n \in \mathbb{N}$.

Consider the statement $S(n)$ for $n = 1$ such that corresponding to $S(1)$ we have the general first order linear ordinary differential equation given as

$$(16) \quad y_{(1)} + P_{11}(x)y = R(x)$$

and the corresponding solution, according to $S(1)$, given as

$$(17) \quad y = \frac{1}{g_1(x)} \int g_1(x) R(x) dx + \frac{C_1}{g_1(x)}$$

where

$$(18) \quad g_1(x) = e^{\int a_{11}(x) dx}$$

$$(19) \quad \text{and } a_{11}(x) = P_{11}(x).$$

But the differential equation given in Eqn. (16) is a first order linear ordinary differential equation whose solution is generally given as (See [4])

$$(20) \quad y = \frac{1}{e^{\int P_{11}(x) dx}} \int e^{\int P_{11}(x) dx} R(x) dx + \frac{C_1}{e^{\int P_{11}(x) dx}},$$

such that the solution given in Eqn. (20) is equivalent to the solution given by Eqn. (17).

Therefore, $S(1)$ is true such that $S(n)$ is true for $n = 1$ and Theorem-1 follows for $n = 1$.

Now, we consider $S(k)$ to be true for $k \in \mathbb{N}$ such that we can state the result corresponding to $S(k)$ as

$$\begin{aligned}
(21) \quad & y_{(k)} + P_{k1}(x)y_{(k-1)} + \cdots + P_{kk-1}(x)y_{(1)} + P_{kk}(x)y = R_0(x) \\
& \implies y = \frac{1}{g_k(x)} \int \frac{g_k(x)}{g_{k-1}(x)} \int \cdots \int \frac{g_2(x)}{g_1(x)} \int g_1(x) R_0(x) \underbrace{dx dx \dots dx}_{k \text{ times}} \\
(22) \quad & + \frac{C_1}{g_k(x)} \int \frac{g_k(x)}{g_{k-1}(x)} \int \cdots \int \frac{g_2(x)}{g_1(x)} \underbrace{dx dx \dots dx}_{(k-1) \text{ times}} \\
& + \cdots + \frac{C_{k-1}}{g_k(x)} \int \frac{g_k(x)}{g_{k-1}(x)} dx + \frac{C_k}{g_k(x)},
\end{aligned}$$

where

$$\begin{aligned}
(23) \quad & g_{k-i+1}(x) = e^{\int a_{i1}(x) dx}, \{i = 1, 2, \dots, k\}, \\
(24) \quad & P_{ij}(x) = a_{ij}(x) + P_{i-1j}(x) \{1 \leq j \leq i-1; 1 \leq i \leq k\}, \\
(25) \quad & a_{11}(x) = P_{11}(x), \\
(26) \quad & P_{i-1j}(x)a_{i1}(x) + P'_{i-1j}(x) = a_{ij+1}(x) \{1 \leq j \leq i-1; 1 \leq i \leq k\}, \\
(27) \quad & P_{i-1i-1}(x)a_{i1}(x) + P'_{i-1i-1}(x) = P_{ii}(x) \{1 < i \leq k\}
\end{aligned}$$

and $i, j, k \in \mathbb{N}$.

Then we need to show that $S(k+1)$ is also true when $S(k)$ is true in order to show that $S(n)$ is true $\forall n \in \mathbb{N}$.

Now we consider the statement $S(n)$ for $n = k+1$, $k \in \mathbb{N}$, such that corresponding to $S(k+1)$ we have the differential equation

$$(28) \quad y_{(k+1)} + p_{k+11}(x)y_{(k)} + \cdots + p_{k+1k}(x)y_{(1)} + p_{k+1k+1}(x)y = R(x),$$

which represents general $(k+1)^{th}$ -order linear ordinary differential equation.

In the above equation let

$$(29) \quad p_{k+1i}(x) = a_{k+1i}(x) + b_{k+1i}(x), \quad i = 1, 2, \dots, k,$$

then Eqn. (28) gets transformed into

$$\begin{aligned}
(30) \quad & y_{(k+1)} + (a_{k+11}(x)y_{(k)} + b_{k+11}(x)y_{(k)}) + \cdots \\
& \cdots + (a_{k+1k}(x)y_{(1)} + b_{k+1k}(x)y_{(1)}) + p_{k+1k+1}(x)y = R(x)
\end{aligned}$$

$$\begin{aligned}
(31) \quad & \implies (y_{(k+1)} + a_{k+11}(x)y_{(k)}) + (b_{k+11}(x)y_{(k)} + a_{k+12}(x)y_{(k-1)}) \\
& + \cdots + (b_{k+1k}(x)y_{(1)} + p_{k+1k+1}(x)y) = R(x).
\end{aligned}$$

We multiply the above equation by a function $g(x)$, which we call here as Integrating Factor, to obtain

$$\begin{aligned}
(32) \quad & (g(x)y_{(k+1)} + g(x)a_{k+11}(x)y_{(k)}) + \cdots \\
& \cdots + (g(x)b_{k+1k}(x)y_{(1)} + g(x)p_{nn}(x)y) = g(x)R(x).
\end{aligned}$$

If we impose the following conditions on $g(x)$,

$$\begin{aligned}
(33) \quad & g'(x) = g(x)a_{k+11}(x), \\
& (b_{k+11}(x)g(x))' = g(x)a_{k+12}(x), \\
& \vdots \\
& (b_{k+1k-1}(x)g(x))' = g(x)a_{k+1k}(x), \\
& (b_{k+1k}(x)g(x))' = g(x)p_{k+1k+1}(x),
\end{aligned}$$

then Eqn. (32) is transformed into

$$(34) \quad (g(x)y_{(k)})' + (g(x)b_{k+11}(x)y_{(k-1)})' + \cdots + (g(x)b_{k+1k-1}(x)y_{(1)})' + (g(x)b_{k+1k}y)' = g(x)R(x).$$

On integrating the above equation with respect to x , we obtain

$$(35) \quad g(x)y_{(k)} + g(x)b_{k+11}(x)y_{(k-1)} + \cdots + g(x)b_{k+1k-1}(x)y_{(1)} + g(x)b_{k+1k}(x)y = \int g(x)R(x) dx + C$$

$$(36) \quad \implies y_{(k)} + b_{k+11}(x)y_{(k-1)} + \cdots + b_{k+1k-1}(x)y_{(1)} + b_{k+1k}(x)y = \frac{1}{g(x)} \int g(x)R(x) dx + \frac{C}{g(x)}.$$

Let

$$(37) \quad \frac{1}{g(x)} \int g(x)R(x) dx + \frac{C}{g(x)} = Q(x)$$

$$(38) \quad \text{and } b_{k+1i}(x) = p_{ki}(x), i = 1, 2, \dots, k$$

such that $p_{ki}(x)$ represent coefficients of the k^{th} -order linear ordinary differential equation obtained, then Eqn. (36) becomes

$$(39) \quad y_{(k)} + p_{k1}(x)y_{(k-1)} + \cdots + p_{kk-1}(x)y_{(1)} + p_{kk}(x)y = Q(x).$$

Also from the set of equations given in Eqn. (33) we have

$$(40) \quad g(x) = e^{\int a_{k+11}(x) dx},$$

$$(41) \quad p_{ki-1}(x)a_{k+11}(x) + p'_{ki-1}(x) = a_{k+1i}(x), i = 2, 3, \dots, k,$$

$$(42) \quad p_{kk}(x)a_{k+11}(x) + p'_{kk}(x) = p_{k+1k+1}(x).$$

We observe that the differential equation obtained in Eqn. (39) is equivalent in form to Eqn. (21) corresponding to $S(k)$. Therefore, we can use the result corresponding to $S(k)$ in the differential equation obtained

in Eqn. (39) to get

$$\begin{aligned}
(43) \quad y &= \frac{1}{g_k(x)} \int \frac{g_k(x)}{g_{k-1}(x)} \int \dots \int \frac{g_2(x)}{g_1(x)} \int g_1(x) Q(x) \underbrace{dx dx \dots dx}_{k \text{ times}} \\
&+ \frac{C_1}{g_k(x)} \int \frac{g_k(x)}{g_{k-1}(x)} \int \dots \int \frac{g_2(x)}{g_1(x)} \underbrace{dx dx \dots dx}_{(k-1) \text{ times}} \\
&+ \dots + \frac{C_{k-1}}{g_k(x)} \int \frac{g_k(x)}{g_{k-1}(x)} dx + \frac{C_k}{g_k(x)},
\end{aligned}$$

where

$$(44) \quad g_{k-i+1}(x) = e^{\int a_{i1}(x) dx}, \quad \{i = 1, 2, \dots, k\},$$

$$(45) \quad p_{ij}(x) = a_{ij}(x) + p_{i-1j}(x) \quad \{1 \leq j \leq i-1; 1 \leq i \leq k\},$$

$$(46) \quad a_{11}(x) = p_{11}(x),$$

$$(47) \quad p_{i-1j}(x) a_{i1}(x) + p'_{i-1j}(x) = a_{ij+1}(x) \quad \{1 \leq j \leq i-1; 1 \leq i \leq k\},$$

$$(48) \quad p_{i-1i-1}(x) a_{i1}(x) + p'_{i-1i-1}(x) = p_{ii}(x) \quad \{1 < i \leq k\}$$

and $i, j, k \in \mathbb{N}$.

On using Eqn. (37) in Eqn. (43) we get

$$\begin{aligned}
(49) \quad y &= \frac{1}{g_k(x)} \int \frac{g_k(x)}{g_{k-1}(x)} \int \dots \int \frac{g_1(x)}{g(x)} \int g(x) R(x) \underbrace{dx dx \dots dx}_{(k+1) \text{ times}} \\
&+ \frac{C}{g_k(x)} \int \frac{g_k(x)}{g_{k-1}(x)} \int \dots \int \frac{g_1(x)}{g(x)} \underbrace{dx dx \dots dx}_{k \text{ times}} \\
&+ \dots + \frac{C_{k-1}}{g_k(x)} \int \frac{g_k(x)}{g_{k-1}(x)} dx + \frac{C_k}{g_k(x)}.
\end{aligned}$$

Let

$$(50) \quad G_1(x) = g(x), G_2(x) = g_1(x), \dots, G_{k+1}(x) = g_k(x),$$

then the solution expression given by Eqn. (49) becomes

$$\begin{aligned}
(51) \quad y &= \frac{1}{G_{k+1}(x)} \int \frac{G_{k+1}(x)}{G_k(x)} \int \dots \int \frac{G_2(x)}{G_1(x)} \int G_1(x) R(x) \underbrace{dx dx \dots dx}_{(k+1) \text{ times}} \\
&+ \frac{C}{G_{k+1}(x)} \int \frac{G_{k+1}(x)}{G_k(x)} \int \dots \int \frac{G_2(x)}{G_1(x)} \underbrace{dx dx \dots dx}_{k \text{ times}} \\
&+ \dots + \frac{C_{k-1}}{G_{k+1}(x)} \int \frac{G_{k+1}(x)}{G_k(x)} dx + \frac{C_k}{G_{k+1}(x)}.
\end{aligned}$$

Also on combining the equations - Eqn. (29), Eqn. (40), Eqn. (41) and Eqn. (42) with the equations - Eqn. (45), Eqn. (44), Eqn. (47) and Eqn. (48) respectively, we obtain the set of equations required

for determining the parameters for the solution expression given by Eqn. (51) as

$$(52) \quad G_{k-i+2}(x) = e^{\int a_{i1}(x) dx}, \{i = 1, 2, \dots, k+1\},$$

$$(53) \quad p_{ij}(x) = a_{ij}(x) + p_{i-1j}(x) \{1 \leq j \leq i-1; 1 \leq i \leq k+1\},$$

$$(54) \quad a_{11}(x) = p_{11}(x),$$

$$(55) \quad p_{i-1j}(x)a_{i1}(x) + p'_{i-1j}(x) = a_{ij+1}(x) \\ \{1 \leq j \leq i-1; 1 \leq i \leq k+1\},$$

$$(56) \quad p_{i-1i-1}(x)a_{i1}(x) + p'_{i-1i-1}(x) = p_{ii}(x) \{1 < i \leq k+1\}$$

and $i, j, k \in \mathbb{N}$ such that Eqn. (51) gives the solution of the differential equation given by Eqn. (28).

Moreover, the desirable form of the result for the solution of the differential equation given by Eqn. (28), according to $S(k+1)$, can be described as

$$(57) \quad y = \frac{1}{g_{k+1}(x)} \int \frac{g_{k+1}(x)}{g_k(x)} \int \dots \int \frac{g_2(x)}{g_1(x)} \int g_1(x) R(x) \underbrace{dx dx \dots dx}_{(k+1) \text{ times}} \\ + \frac{C_1}{g_{k+1}(x)} \int \frac{g_{k+1}(x)}{g_k(x)} \int \dots \int \frac{g_2(x)}{g_1(x)} \underbrace{dx dx \dots dx}_{k \text{ times}} \\ + \dots + \frac{C_k}{g_{k+1}(x)} \int \frac{g_{k+1}(x)}{g_k(x)} dx + \frac{C_{k+1}}{g_{k+1}(x)},$$

where

$$(58) \quad g_{k-i+2}(x) = e^{\int a_{i1}(x) dx}, \{i = 1, 2, \dots, k+1\},$$

$$(59) \quad p_{ij}(x) = a_{ij}(x) + p_{i-1j}(x) \{1 \leq j \leq i-1; 1 \leq i \leq k+1\},$$

$$(60) \quad a_{11}(x) = p_{11}(x),$$

$$(61) \quad p_{i-1j}(x)a_{i1}(x) + p'_{i-1j}(x) = a_{ij+1}(x) \\ \{1 \leq j \leq i-1; 1 \leq i \leq k+1\},$$

$$(62) \quad p_{i-1i-1}(x)a_{i1}(x) + p'_{i-1i-1}(x) = p_{ii}(x) \{1 < i \leq k+1\}$$

and $i, j, k \in \mathbb{N}$ such that Eqn. (57) gives the solution of the differential equation given by Eqn. (28).

We observe that the result obtained through Eqn. (51) is equivalent to the result stated in Eqn. (57), which is the desired result according to $S(k+1)$.

Thus, $S(k+1)$ is true such that when $S(k)$ is true, $S(k+1)$ is also true for $k \in \mathbb{N}$.

Hence, we have shown the following

(1) $S(1)$ is true, i.e. Theorem-1 follows for $n = 1$.

(2) $S(k+1)$ is true when $S(k)$ is true for $k \in \mathbb{N}$.

Therefore, by the Principle of Mathematical Induction, $S(n)$ is true $\forall n \in \mathbb{N}$ such that Theorem-1 follows $\forall n \in \mathbb{N}$. $\square$

From the above result, it follows that the general solution of the general n^{th} -order linear ordinary differential equation is given as sum of two parts - the complementary solution part and the particular solution part.

The complementary solution part is the solution of the homogeneous equation corresponding to the concerned differential equation given by Eqn. (1) such that in the solution expression given in Eqn. (3) the part of the solution having arbitrary constants gives the complementary solution part. This can be justified by the fact that by putting $R(x) = 0$ in the solution expression we get only the part of the solution having arbitrary constants since the series of integrals involving $R(x)$ represent integrals without the involvement of constants of integration as they have been already accounted for by the arbitrary constants - $C_1, C_2, \dots, C_n$ such that the series of integrals involving $R(x)$ evaluates to 0 when $R(x) = 0$.

Whereas, the part of the solution having series of n integrals and $R(x)$ represents the particular integral part of the general solution as only this term satisfies the non-homogeneous equation as given in Eqn. (1) when $R(x) \neq 0$.

3. RESULT FOR A SPECIAL CASE

In this section we describe the result for a special case based on Theorem-1 which gives the complete solution of general n^{th} -order linear ordinary differential equation when $n - 1$ out of n complementary solutions are known. The result is presented in form of a corollary of Theorem-1 which is stated here along with the proof as follows

Corollary 1. *If $n - 1$ out of n complementary solutions of the differential equation,*

$$(63) \quad y_{(n)} + P_{n1}(x)y_{(n-1)} + \dots + P_{nn-1}(x)y_{(1)} + P_{nn}(x)y = R(x),$$

are known then its solution can be determined completely such that the solution is given as

$$(64) \quad \begin{aligned} y = & \frac{1}{g_n(x)} \int \frac{g_n(x)}{g_{n-1}(x)} \int \dots \int \frac{g_2(x)}{g_1(x)} \int g_1(x) R(x) \underbrace{dx dx \dots dx}_{n \text{ times}} \\ & + \frac{C_1}{g_n(x)} \int \frac{g_n(x)}{g_{n-1}(x)} \int \dots \int \frac{g_2(x)}{g_1(x)} \underbrace{dx dx \dots dx}_{(n-1) \text{ times}} \\ & + \dots + \frac{C_{n-1}}{g_n(x)} \int \frac{g_n(x)}{g_{n-1}(x)} dx + \frac{C_n}{g_n(x)}, \end{aligned}$$

and the Integrating Factors $-g_i(x)$, $i = 1, 2, \dots, n$ are given as

$$(65) \quad \frac{1}{g_{n-k}(x)} = \frac{1}{g_n(x)} \prod_{i=1}^k F(i, i), \quad k = 1, 2, \dots, n-2,$$

$$(66) \quad \frac{1}{g_n(x)} = f_n(x),$$

$$(67) \quad g_1(x) = e^{\int a_{n1}(x) dx},$$

where $F(k, j)$ is defined recursively as

$$(68) \quad F(k, j) = \frac{d}{dx} \left(\frac{F(k, j-1)}{F(k-1, j-1)} \right)$$

$$(69) \quad \text{and } F(k, 1) = \frac{d}{dx} (g_n(x) f_{n-k}(x)),$$

$f_i(x)$, $i = 2, 3, \dots, n$, represent $n-1$ known complementary solutions, $a_{n1}(x)$ is defined as

$$(70) \quad a_{n1}(x) = P_{n1}(x) - (a_{n-11}(x) + \dots + a_{21}(x) + P_{11}(x))$$

such that the parameters required for determining $a_{n1}(x)$ are given as

$$(71) \quad a_{j1}(x) = \frac{g'_{n-j+1}(x)}{g_{n-j+1}(x)}, \quad j = 2, 3, \dots, n-1,$$

$$(72) \quad \text{and } a_{11}(x) = P_{11}(x) = \frac{g'_n(x)}{g_n(x)}.$$

Proof. Consider the general n^{th} -order linear ordinary differential equation given as

$$(73) \quad y_{(n)} + P_{n1}(x)y_{(n-1)} + \dots + P_{nn-1}(x)y_{(1)} + P_{nn}(x)y = R(x),$$

such that it has solution given by Theorem-1 as

$$(74) \quad \begin{aligned} y = & \frac{1}{g_n(x)} \int \frac{g_n(x)}{g_{n-1}(x)} \int \dots \int \frac{g_2(x)}{g_1(x)} \int g_1(x) R(x) \underbrace{dx dx \dots dx}_{n \text{ times}} \\ & + \frac{C_1}{g_n(x)} \int \frac{g_n(x)}{g_{n-1}(x)} \int \dots \int \frac{g_2(x)}{g_1(x)} \underbrace{dx dx \dots dx}_{(n-1) \text{ times}} \\ & + \dots + \frac{C_{n-1}}{g_n(x)} \int \frac{g_n(x)}{g_{n-1}(x)} dx + \frac{C_n}{g_n(x)}, \end{aligned}$$

where

$$(75) \quad g_{n-i+1}(x) = e^{\int a_{i1}(x) dx}, \quad \{i = 1, 2, \dots, n\},$$

$$(76) \quad P_{ij}(x) = a_{ij}(x) + P_{i-1j}(x) \{1 \leq j \leq i-1; 1 \leq i \leq n\},$$

$$(77) \quad a_{11}(x) = P_{11}(x),$$

$$(78) \quad P_{i-1j}(x)a_{i1}(x) + P'_{i-1j}(x) = a_{ij+1}(x) \{1 \leq j \leq i-1; 1 \leq i \leq n\},$$

$$(79) \quad P_{i-1i-1}(x)a_{i1}(x) + P'_{i-1i-1}(x) = P_{ii}(x) \{1 < i \leq n\}$$

and $i, j, n \in \mathbb{N}$.

Let $f_2(x), f_3(x), \dots, f_n(x)$ represent $n - 1$ (out of n) known complementary solutions corresponding to the differential equation given by Eqn. (73) such that we consider the following set of equations for determining the Integrating Factors - $g_i(x)$, $i = 2, 3, \dots, n$ where we put terms of complementary solution as given in the solution expression equal to the solutions given such that we have

$$\begin{aligned}
 & \frac{1}{g_n(x)} = f_n(x), \\
 & \frac{1}{g_n(x)} \int \frac{g_n(x)}{g_{n-1}(x)} dx = f_{n-1}(x), \\
 & \vdots \\
 & \frac{1}{g_n(x)} \int \frac{g_n(x)}{g_{n-1}(x)} \int \dots \int \frac{g_3(x)}{g_2(x)} \underbrace{dx dx \dots dx}_{(n-2) \text{ times}} = f_2(x).
 \end{aligned}
 \tag{80}$$

From the above set of equations we get

$$\begin{aligned}
 & g_n(x) = \frac{1}{f_n(x)}, \\
 & \frac{1}{g_n(x)} \int \frac{g_n(x)}{g_{n-1}(x)} dx = f_{n-1}(x), \\
 & \implies \frac{g_n(x)}{g_{n-1}(x)} = \frac{d}{dx} (g_n(x) f_{n-1}(x)), \\
 & \implies \frac{1}{g_{n-1}(x)} = \frac{1}{g_n(x)} \frac{d}{dx} (g_n(x) f_{n-1}(x)), \\
 & \frac{1}{g_n(x)} \int \frac{g_n(x)}{g_{n-1}(x)} \int \frac{g_{n-1}(x)}{g_{n-2}(x)} dx dx = f_{n-2}(x), \\
 & \implies \frac{g_n(x)}{g_{n-1}(x)} \int \frac{g_{n-1}(x)}{g_{n-2}(x)} dx = \frac{d}{dx} (g_n(x) f_{n-2}(x)), \\
 & \implies \frac{g_{n-1}(x)}{g_{n-2}(x)} = \frac{d}{dx} \left(\frac{\frac{d}{dx} (g_n(x) f_{n-2}(x))}{\frac{d}{dx} (g_n(x) f_{n-1}(x))} \right).
 \end{aligned}
 \tag{81}$$

(By using Eqn. (82))

Based on the above observations, we define a recursive function $F(k, j)$ as

$$F(k, j) = \frac{d}{dx} \left(\frac{F(k, j-1)}{F(k-1, j-1)} \right)
 \tag{84}$$

$$\text{and } F(a, 1) = \frac{d}{dx} (g_n(x) f_{n-a}(x)).
 \tag{85}$$

Then it follows that

$$(86) \quad \frac{1}{g_n(x)} = f_n(x),$$

$$(87) \quad \frac{1}{g_{n-1}(x)} = \frac{1}{g_n(x)} F(1, 1),$$

$$(88) \quad \frac{1}{g_{n-2}(x)} = \frac{1}{g_n(x)} F(1, 1) F(2, 2),$$

$$(89) \quad \text{and similarly } \frac{1}{g_{n-3}(x)} = \frac{1}{g_n(x)} F(1, 1) F(2, 2) F(3, 3),$$

$\vdots$

$$(90) \quad \frac{1}{g_{n-i}(x)} = \frac{1}{g_n(x)} F(1, 1) F(2, 2) \cdots F(i, i),$$

such that we obtain

$$(91) \quad \frac{1}{g_{n-i}(x)} = \frac{1}{g_n(x)} \prod_{j=1}^i F(j, j), \quad i = 1, 2, \dots, n-2$$

$$(92) \quad \frac{1}{g_n(x)} = f_n(x),$$

$$(93) \quad \text{and } \frac{1}{g_{n-i}(x)} = \frac{1}{g_{n-i-1}(x)} F(i, i), \quad i = 1, 2, \dots, n-2,$$

where these equations give the expression for the Integrating Factors - $g_n(x), g_{n-1}(x), \dots, g_2(x)$.

Also from Eqn. (75) we obtain

$$(94) \quad \begin{aligned} g_{n-k+1}(x) &= e^{\int a_{k1}(x) dx} \\ \implies a_{k1}(x) &= \frac{g'_{n-k+1}(x)}{g_{n-k+1}(x)}, \quad k = 1, 2, \dots, n-1, \end{aligned}$$

such that the above equation gives the expression for the functions $a_{k1}(x)$, $k = 1, 2, \dots, n-1$ in terms of the Integrating Factors given by $g_{n-k+1}(x)$, $k = 1, 2, \dots, n-1$ where the expression of each Integrating Factor is given in terms of the given complementary solutions by the equations - Eqn. (91), Eqn. (92) and Eqn. (93).

From Eqn. (76), for $j = 1, i = n$, we get

$$(95) \quad \begin{aligned} P_{n1}(x) &= a_{n1}(x) + P_{n-11}(x) \\ &= a_{n1}(x) + (a_{n-11}(x) + P_{n-21}(x)) \\ &= a_{n1}(x) + a_{n-11}(x) + (a_{n-21}(x) + P_{n-31}(x)) \\ &= \cdots = a_{n1}(x) + a_{n-11}(x) + \cdots + a_{21}(x) + a_{11}(x) \end{aligned}$$

$$(96) \quad \implies a_{n1}(x) = P_{n1}(x) - \sum_{i=1}^{n-1} a_{i1}(x),$$

such that each $a_{i1}(x)$ is given by Eqn. (94) for $i = 1, 2, \dots, n-1$ and $P_{n1}(x)$ is the coefficient of $y_{(n-1)}$ as given in the differential equation given by Eqn. (73). Also, from Eqn. (77) we have $a_{11}(x) = P_{11}(x)$ such that the above equation can be equivalently expressed as

$$(97) \quad a_{n1}(x) = P_{n1}(x) - \sum_{i=2}^{n-1} a_{i1}(x) - P_{11}(x).$$

Thus, Eqn. (96) and Eqn. (97) give the expression for the function $a_{n1}(x)$ such that the remaining Integrating Factor $g_1(x)$ can be determined by using the relation given in Eqn. (75) as

$$(98) \quad g_1(x) = e^{\int a_{n1}(x) dx},$$

where $a_{n1}(x)$ is as given by Eqn. (96) (and also by Eqn. (97)).

Hence, we have determined all the Integrating Factors, given by the equations - Eqn. (91), Eqn. (92), Eqn. (93) and Eqn. (98), by making use of the $n-1$ complementary solutions, $f_2(x), f_3(x), \dots, f_n(x)$, of the given differential equation (Eqn. (73)).

Therefore, we conclude by stating that when $n-1$ out of n complementary solutions of the differential equation given by Eqn. (73) as

$$y_{(n)} + P_{n1}(x)y_{(n-1)} + \dots + P_{nn-1}(x)y_{(1)} + P_{nn}(x)y = R(x)$$

are known where the solution is given by Eqn. (74) as

$$\begin{aligned} y = & \frac{1}{g_n(x)} \int \frac{g_n(x)}{g_{n-1}(x)} \int \dots \int \frac{g_2(x)}{g_1(x)} \int g_1(x) R(x) \underbrace{dx dx \dots dx}_{n \text{ times}} \\ & + \frac{C_1}{g_n(x)} \int \frac{g_n(x)}{g_{n-1}(x)} \int \dots \int \frac{g_2(x)}{g_1(x)} \underbrace{dx dx \dots dx}_{(n-1) \text{ times}} \\ & + \dots + \frac{C_{n-1}}{g_n(x)} \int \frac{g_n(x)}{g_{n-1}(x)} dx + \frac{C_n}{g_n(x)}, \end{aligned}$$

then all of the Integrating Factors can be determined as

$$(99) \quad g_n(x) = \frac{1}{f_n(x)}, \text{ (From Eqn. (92))}$$

$$(100) \quad g_{n-i}(x) = \frac{g_n(x)}{\prod_{j=1}^i F(j, i)}, \quad i = 1, 2, \dots, n-2, \text{ (From Eqn. (91))}$$

$$(101) \quad g_1(x) = e^{\int a_{n1}(x) dx}, \text{ (From Eqn. (98))}$$

where $F(j, k)$ is defined recursively as

$$(102) \quad F(j, k) = \frac{d}{dx} \left(\frac{F(j, k-1)}{F(j-1, k-1)} \right) \text{ (From Eqn. (84))}$$

$$(103) \quad \text{and } F(a, 1) = \frac{d}{dx} (g_n(x) f_{n-a}(x)) \text{ (From Eqn. (85))},$$

$f_2(x), f_3(x), \dots, f_n(x)$ represent $n-1$ (out of n) known complementary solutions of the differential equation given by Eqn. (73), $a_{n1}(x)$ is given as

$$(104) \quad a_{n1}(x) = P_{n1}(x) - \sum_{i=1}^{n-1} a_{i1}(x), \text{ (From Eqn. (96))}$$

each $a_{i1}(x)$ for $i = 1, 2, \dots, n-1$ is given as

$$(105) \quad a_{k1}(x) = \frac{g'_{n-k+1}(x)}{g_{n-k+1}(x)}, \quad k = 1, 2, \dots, n-1 \text{ (From Eqn. (94))}$$

and $P_{n1}(x)$ is the coefficient of $y_{(n-1)}$ as given in the differential equation given by Eqn. (73). Then by using the obtained Integrating Factors in the solution expression (given by Eqn. (74)) of the concerned differential equation (given by Eqn. (73)), the complete solution of the concerned differential equation can be obtained. $\square$

REFERENCES

- [1] Edward J. Norminton Erwin Kreyszig, Herbert Kreyszig. *Advanced Engineering Mathematics*, chapter 3, pages 105–106. John Wiley And Sons, Inc., 2010.
  - [2] Edward J. Norminton Erwin Kreyszig, Herbert Kreyszig. *Advanced Engineering Mathematics*, chapter 3, pages 116–117. John Wiley And Sons, Inc., 2010.
  - [3] Edward J. Norminton Erwin Kreyszig, Herbert Kreyszig. *Advanced Engineering Mathematics*, chapter 3, pages 111–112. John Wiley And Sons, Inc., 2010.
  - [4] S.R.K. Iyengar R.K. Jain. *Advanced Engineering Mathematics*, chapter 4, pages 261–275. Alpha Science International Ltd., 2002.
- Email address:* rajnishkrjha.2020@gmail.com