Over-expression of BLK in metastatic human breast cancer.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY USA

Metastasis to the brain is a clinical problem in patients with breast cancer¹⁻³. We mined published microarray data^{4,5} to compare primary and metastatic tumor transcriptomes for the discovery of genes associated with metastasis to the lymph nodes and the brain in humans with metastatic breast cancer. We found that the B-lymphocyte kinase, encoded by BLK, was among the genes whose expression was most different in the lymph node metastases of patients with metastatic breast cancer as compared to primary tumors of the breast. BLK mRNA was present at increased quantities in lymph node metastases as compared to primary tumors of the breast. Importantly, expression of BLK in primary tumors was significantly correlated with patient recurrence-free survival. Modulation of BLK expression may be relevant to the biology by which tumor cells metastasize from the breast to the brain in humans with metastatic breast cancer. Inhibition of BLK represents a potential therapeutic approach in halting progression of lymph node metastatic breast cancer.

Keywords: breast cancer, metastasis, brain metastases, central nervous system metastases, the B-lymphocyte kinase, BLK, systems biology of breast cancer, targeted therapeutics in breast cancer.

One report described a 34% incidence of central nervous system metastases in patients treated with trastuzumab for breast cancer². More recently, the NEfERT-T clinical trial⁶ which compared administration of either neratinib or trastuzumab in conjunction with paclitaxel demonstrated that in a randomized, controlled setting, in breast cancer patients treated with neratinib, not only was the incidence of central nervous system recurrence significantly lower, the time to central nervous system metastasis was significantly delayed as compared to patients administered trastuzumab⁶. The alarmingly high rate of central nervous system metastasis described, as well as data, both anecdotal² and from a randomized, controlled setting⁶ illustrating that treatment with trastuzumab may be associated with these events demands an enhanced understanding of the transcriptional makeup of brain metastatic tissues to support identification of therapeutic targets, whether they are treatment related or not. We performed a global comparative analysis of lymph node and brain metastatic tumor tissues in patients with metastatic breast cancer, comparing to primary tumors and normal breast tissues, respectively^{4,5}. We discovered transcriptome-wide differential expression of the gene encoding the B-lymphocyte kinase, BLK, in lymph node and brain metastatic tissues of patients with metastatic breast cancer.

Methods

We used datasets GSE10893⁴ and GSE52604⁵ for this global differential gene expression analysis of brain metastatic breast cancer in conjunction with GEO2R. GSE10893 was generated using Agilent-011521 Human 1A Microarray G4110A technology with $n=71$ primary breast tumors and $n=7$ lymph node metastases from patients with breast cancer; analysis was performed using platform GPL887. GSE52604 was generated using Agilent-014850 Whole Human Genome Microarray 4x44K G4112F with $n=10$ normal breast tissues and $n=35$ brain metastases from patients with breast cancer; analysis was performed using platform GPL6480. The Benjamini and Hochberg method of p -value adjustment was used for ranking of differential expression but raw p -values were used to assess statistical significance of global differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. A statistical test was performed to evaluate whether BLK gene expression was significantly different between primary tumors of the breast and lymph node metastases in humans with breast cancer using a two-tailed t-test. For Kaplan-Meier survival analysis, we used the Kaplan-Meier plotter online tool⁷ for correlation of BLK mRNA expression levels with recurrence-free survival in $n=3951$ breast cancer patients.

Results

We performed global comparative transcriptome analysis of metastatic tumor tissues of patients with metastatic breast cancer using published microarray data^{4,5} to describe the transcriptional landscape of metastasis in human breast cancer in an unbiased fashion and for the discovery of novel therapeutic targets.

BLK is differentially expressed in the lymph node metastases of patients with metastatic breast cancer.

Through blind, systems-level analysis of published microarray data⁴, we identified the

B-lymphocyte kinase, encoded by BLK, as a differentially expressed gene in the lymph node metastatic tissues of humans with breast cancer (Table 1). When sorting each of the genes expressed in lymph node metastases based on significance of difference as compared to primary tumors of the breast in patients with breast cancer, BLK ranked 2 out of 18034 total transcripts (Chart 1), equating to 99.99% differential expression. Differential expression of BLK in the lymph node metastases of patients with metastatic breast cancer was statistically significant (Chart 1; $p=7.29E-08$).

Differential gene expression analyses that utilize primary and metastatic tumor tissues to discover genes associated with metastasis can be challenging due to the relatively heterogeneous cellular composition and constitution of the tumor as compared to normal, untransformed (benign) tissues. Thus, to attempt to validate differential transcriptome-wide differential expression of BLK in human metastatic breast cancer, we queried a second microarray dataset⁵, here comparing normal breast tissues to brain metastases. Again, we identified BLK as a differentially expressed gene in the metastatic tissues of patients with breast cancer (Chart 2). When sorting each of the genes expressed in brain metastases based on significance of difference as compared to normal breast tissues, BLK ranked 5674 out of 41093 total transcripts (Chart 2), equating to 86.2% differential expression. Differential expression of BLK in the brain metastases of patients with metastatic breast cancer was statistically significant (Chart 2; $p=3.95E-04$). Thus, differential expression of BLK, transcriptome-wide, in the metastatic tissues of women with metastatic breast cancer was conserved across two independent microarray datasets, both when compared to benign and transformed breast tissues.

BLK is expressed at higher levels in the lymph node metastases of patients with metastatic breast cancer.

We obtained exact mRNA expression levels for BLK, in primary tumors of the breast and in lymph node metastases of patients with metastatic breast cancer to determine direction and statistical significance of change in BLK expression in lymph node metastatic tissues. BLK was expressed at higher levels in the lymph node metastases of patients with breast cancer as compared to primary tumors of the breast, and this difference was statistically significant (Figure 1; $p<0.00001$).

BLK expression is significantly correlated with survival outcomes in human breast cancer.

We performed Kaplan-Meier survival analysis⁸ in $n=3951$ breast cancer patients in total, to evaluate whether BLK tumor expression was correlated with survival outcomes in breast cancer. We observed a statistically significant correlation between primary tumor expression of BLK and recurrence-free survival (RFS) in patients with breast cancer (Figure 2). Patients whose primary tumors expressed low levels of BLK possessed median RFS of 185.16 months, while patients whose tumors expressed high levels of BLK possessed median OS of 216.66 months. This difference in RFS based on BLK tumor expression in patients with breast cancer was statistically significant (Figure 2, Chart 3; logrank p -value: $1.2e-10$; hazard ratio: 0.7 (0.63-0.78)).

Thus, by mining published microarray data^{4,5} in an unbiased and systematic fashion, we identified the B-lymphocyte kinase, encoded by BLK, as among the genes whose expression was most different, transcriptome-wide, in the lymph node and brain metastases of patients with breast cancer, when compared to primary tumors of the breast and to normal breast tissues, respectively; we observed significantly increased expression of BLK in lymph node metastases as compared to primary tumors of the breast. Further, we found a significant correlation between BLK expression and patient survival outcomes, as recurrence-free survival was significantly greater in patients whose primary tumors expressed higher levels of BLK as compared to patients whose primary tumors expressed lower levels of BLK.

Discussion

We provided evidence here that the B-lymphocyte kinase, encoded by BLK, is among the genes whose expression is most different in the lymph node and brain metastases of patients with metastatic breast cancer, that BLK mRNA is present at significantly increased quantities in lymph node metastatic tissues as compared to primary tumors of the breast, and that primary tumor BLK expression is significantly correlated with patient survival outcomes in human breast cancer. Evaluation of the effects of genetic depletion of BLK in mouse models of metastatic breast cancer on metastasis to the central nervous system is merited. While we found that higher primary tumor expression of BLK was correlated with enhanced patient survival, patient survival outcomes based on primary tumor expression do not necessarily correlate with phenotypic consequence of inhibition. Moreover, the survival data we presented here is based on expression of BLK in primary tumors, not the lymph node metastatic tissues, in which we found significantly increased expression of BLK. An inhibitor of BLK, dasatinib, is available for use in the clinic⁸. Modulation of BLK expression may be relevant to the processes by which breast cancer cells exit the breast, enter the vasculature and/or lymphatics, reside in the lymph nodes, evade immune clearance, breach the blood-brain barrier and colonize the brain. Inhibition of BLK represents a potential therapeutic approach in halting progression of lymph node metastatic breast cancer.

References

1. Lin, N.U., Amiri-Kordestani, L., Palmieri, D., Liewehr, D.J. and Steeg, P.S., 2013. CNS metastases in breast cancer: old challenge, new frontiers.
2. Bendell, J.C., Domchek, S.M., Burstein, H.J., Harris, L., Younger, J., Kuter, I., Bunnell, C., Rue, M., Gelman, R. and Winer, E., 2003. Central nervous system metastases in women who receive trastuzumab-based therapy for metastatic breast carcinoma. *Cancer*, 97(12), pp.2972-2977.
3. Tsukada, Y., Fouad, A., Pickren, J.W. and Lane, W.W., 1983. Central nervous system metastasis from breast carcinoma autopsy study. *Cancer*, 52(12), pp.2349-2354.
4. Weigman, V.J., Chao, H.H., Shabalin, A.A., He, X., Parker, J.S., Nordgard, S.H., Grushko, T., Huo, D., Nwachukwu, C., Nobel, A. and Kristensen, V.N., 2012. Basal-like Breast cancer DNA copy number losses identify genes involved in genomic instability, response to therapy, and patient survival. *Breast cancer research and treatment*, 133(3), pp.865-880.
5. Salhia, B., Kiefer, J., Ross, J.T., Metapally, R., Martinez, R.A., Johnson, K.N., DiPerna, D.M., Paquette, K.M., Jung, S., Nasser, S. and Wallstrom, G., 2014. Integrated genomic and epigenomic analysis of breast cancer brain metastasis. *Plos one*, 9(1), p.e85448.
6. Awada, A., Colomer, R., Inoue, K., Bondarenko, I., Badwe, R.A., Demetriou, G., Lee, S.C., Mehta, A.O., Kim, S.B., Bachelot, T. and Goswami, C., 2016. Neratinib plus paclitaxel vs trastuzumab plus paclitaxel in previously untreated metastatic ERBB2-positive breast cancer: the NEfERT-T randomized clinical trial. *JAMA oncology*, 2(12), pp.1557-1564.
7. Györfy, B., Lanczky, A., Eklund, A.C., Denkert, C., Budczies, J., Li, Q. and Szallasi, Z., 2010. An online survival analysis tool to rapidly assess the effect of 22,277 genes on breast cancer prognosis using microarray data of 1,809 patients. *Breast cancer research and treatment*, 123(3), pp.725-731.
8. Petersen, D.L., Krejsgaard, T., Berthelsen, J., Fredholm, S., Willerslev-Olsen, A., Sibbesen, N.A., Bonefeld, C.M., Andersen, M.H., Francavilla, C., Olsen, J.V. and Hu, T., 2014. B-lymphoid tyrosine kinase (Blk) is an oncogene and a potential target for therapy with dasatinib in cutaneous T-cell lymphoma (CTCL). *Leukemia*, 28(10), pp.2109-2112.

Rank: 2
Probe ID: 15034
p-value: 7.29E-08
t: -5.9394347
B: 7.3854
Gene: BLK
Gene name: B-lymphocyte kinase

Chart 1: BLK is differentially expressed in lymph node metastatic breast cancer when comparing lymph node metastases to primary tumors of the breast.

The rank of global differential expression, the probe/transcript ID, the *p*-value with respect to differential expression transcriptome-wide, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the groups compared, the gene and gene name are listed in this chart.

Rank: 5674
probe ID: A_23_P31725
p-value: 3.95E-04
t: 3.8168128
B: -0.351365
Gene: BLK
Gene name: B-lymphocyte kinase

Chart 2: BLK is differentially expressed in brain metastatic breast cancer when comparing brain metastases to normal breast tissues.

The rank of global differential expression, probe/transcript ID, the *p*-value with respect to differential expression transcriptome-wide, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the groups compared, the gene and gene name are listed in this chart.

Figure 1: BLK is expressed at significantly higher levels in the lymph node metastases of patients with metastatic breast cancer when compared to primary tumors of the breast.

The mRNA expression level of BLK in primary tumors of the breast (left) and in lymph node metastases of women with metastatic breast cancer (right) is graphically depicted; the result of a statistical test evaluating significance of difference in BLK expression between primary tumors of the breast and lymph node metastases, a p -value, is $p < 0.0001$.

Figure 2: Significant correlation between BLK primary tumor expression and recurrence-free survival in patients with breast cancer.

Depicted in this Kaplan-Meier plot is the probability of recurrence-free survival for $n=3951$ total patients stratified into two groups, based on low or high expression of BLK in patient primary tumors. The log rank p -value denoting statistical significance of difference in recurrence-free survival when comparing the two groups, as well as hazard ratio for this comparison is listed above. Listed below is the number of patients at risk (number of patients alive) per interval, after stratification based on BLK expression; in the first interval, number at risk is number of patients alive; in each subsequent interval, number at risk is the number at risk less those who have expired or are censored.

Low BLK expression: 185.16 months
High BLK expression: 216.66 months

Chart 3: Median recurrence-free survival is inferior in patients with low primary tumor expression of BLK.

The median OS (recurrence-free survival) of patients with low primary tumor expression of BLK and high primary tumor expression of BLK is listed in this chart, for $n=3951$ breast cancer patients.