A pseudogene of WDR5 is differentially expressed in brain metastatic breast cancer.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY USA

Metastasis to the brain is a clinical problem in patients with breast cancer¹⁻³. We mined published microarray data^{4,5} to compare primary and metastatic tumor transcriptomes for the discovery of genes associated with brain metastasis in humans with metastatic breast cancer. We found that a non-coding RNA transcribed from the WDR5 locus (WDR5p), a pseudogene transcript, was among the transcripts whose quantity was most significantly different in the brain metastases of patients with metastatic breast cancer. WDR5p transcript was present at increased quantities in brain metastatic tissues as compared to primary tumors of the breast. Modulation of WDR5p expression may be relevant to the biology by which tumor cells metastasize from the breast to the brain in humans with metastatic breast cancer.

Keywords: breast cancer, metastasis, brain metastasis, central nervous system metastasis, WDR5 pseudogene, WDR5p, LOC401127, systems biology of breast cancer, targeted therapeutics in breast cancer.

One report described a 34% incidence of central nervous system metastases in patients treated with trastuzumab for breast cancer². More recently, the NEfERT-T clinical trial⁶ which compared administration of either neratinib or trastuzumab in conjunction with paclitaxel demonstrated that in a randomized, controlled setting, in breast cancer patients treated with neratinib, not only was the incidence of central nervous system recurrence significantly lower, the time to central nervous system metastasis was significantly delayed as compared to patients administered trastuzumab⁶. The alarmingly high rate of central nervous system metastasis described, as well as data, both anecdotal² and from a randomized, controlled setting⁶ illustrating that treatment with trastuzumab may be associated with these events demands an enhanced understanding of the transcriptional makeup of brain metastatic tissues to support identification of therapeutic targets, whether they are treatment related or not. We performed a global comparative analysis of primary and metastatic tumors in patients with brain metastatic breast cancer^{4,5}. We discovered differential and increased expression of a pseudogene transcribed at the WDR5 locus in brain metastatic tissues of patients with metastatic breast cancer.

Methods

We used datasets GSE10893⁴ and GSE52604⁵ for this global differential gene expression analysis of brain metastatic breast cancer in conjunction with GEO2R. GSE10893 was generated using Agilent-011521 Human 1A Microarray G4110A technology with $n=11$ primary breast tumors and $n=3$ brain metastases from patients with breast cancer; analysis was performed using platform GPL885. GSE52604 was generated using Agilent-014850 Whole Human Genome Microarray 4x44K G4112F with $n=10$ normal breast tissues and $n=35$ brain metastases from patients with breast cancer; analysis was performed using platform GPL6480. The Benjamini and Hochberg method of p -value adjustment was used for ranking of differential expression but raw p -values were used to assess statistical significance of global differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. A statistical test was performed to evaluate whether WDR5p gene expression was significantly different between primary tumors of the breast and brain metastases in humans with breast cancer using a two-tailed t-test.

Results

We performed global comparative transcriptome analysis of metastatic and primary tumor tissues of patients with metastatic breast, as well as normal tissues of the breast using published microarray data^{4,5} to describe the transcriptional landscape of brain metastasis in human breast cancer in an unbiased fashion and for the discovery of novel therapeutic targets.

WDR5p is differentially expressed in the brain metastases of patients with brain metastatic breast cancer.

Through blind, systems-level analysis of published microarray data⁴, we identified a pseudogene produced at the WDR5 locus (LOC401127), which we term WDR5p, as a differentially expressed transcript in the brain metastatic tissues of humans with breast cancer (Table 1). When sorting each of the genes expressed in brain metastases based on significance of difference as compared to primary tumors of the breast in patients with breast cancer, WDR5p ranked 148 out of 17418 total transcripts (Chart 1), equating to 99.2% differential expression.

Differential expression of WDR5p in the brain metastases of patients with metastatic breast cancer was statistically significant (Chart 1; $p=5.56E-04$).

WDR5p is differentially expressed in the lymph node metastases of patients with breast cancer.

Differential gene expression analyses that utilize primary and metastatic tumor tissues to discover genes associated with metastasis can be challenging due to the relatively heterogeneous cellular composition and constitution of the tumor as compared to normal, untransformed (benign) tissues. Thus, to attempt to validate differential transcriptome-wide differential expression of WDR5p in brain metastases in human metastatic breast cancer, we queried a second microarray dataset⁵, here comparing normal breast tissues rather than primary tumors of the breast to brain metastases. Again, we identified WDR5p as a differentially expressed gene in the brain metastatic tissues of patients with breast cancer (Chart 2). When sorting each of the genes expressed in brain metastases based on significance of difference as compared to normal breast tissues, WDR5p ranked 1024 out of 41093 total transcripts (Chart 2), equating to 97.5% differential expression. Differential expression of WDR5p in the brain metastases of patients with metastatic breast cancer was statistically significant (Chart 2; $p=1.94e-07$). Thus, differential expression of WDR5p, transcriptome-wide, in the brain metastases of women with metastatic breast cancer was conserved across two independent microarray datasets, both when compared to benign and transformed breast tissues. This suggests that WDR5p differential expression in human brain metastatic breast cancer is a *bona fide* biological process.

WDR5p is expressed at higher levels in the brain metastases of patients with metastatic breast cancer.

We obtained exact mRNA expression levels for WDR5p, in primary tumors of the breast and in brain metastasis of patients with brain metastatic breast cancer to determine direction and statistical significance of change in WDR5p expression in brain metastatic tissues. We observed significantly higher expression of WDR5p in the brain metastases of patients with breast cancer as compared to primary tumors of the breast: WDR5p was expressed at -0.26 ± 0.38 arbitrary units (AU) in primary tumors of the breast, while it was expressed at 0.94 ± 0.57 AU in brain metastatic tissues (Figure 1). The difference in WDR5p mRNA levels between primary tumors of the breast and brain metastatic tissues was statistically significant (Figure 1; $p=0.000916$).

Thus, by mining published microarray data^{4,5} in an unbiased and systematic fashion, we identified an open reading frame, encoded by WDR5p, as among the genes whose expression was most different, transcriptome-wide, in the brain metastases of patients with breast cancer, when compared to primary tumors of the breast and to normal breast tissues; we observed significantly increased expression of WDR5p in brain metastases as compared to primary tumors of the breast.

Discussion

We provided evidence here that a pseudogene transcribed from the WDR5 locus, LOC401127, which we have termed WDR5p, is among the genes whose expression is most different in the brain metastases of patients with metastatic breast cancer and that WDR5p mRNA is present at significantly increased quantity in brain metastatic tissues as compared to primary

1 tumors of the breast. Evaluation of the effects of genetic depletion of WDR5p in mouse models
2 of metastatic breast cancer on metastasis to the central nervous system is merited. Modulation of
3 WDR5p expression may be relevant to the processes by which breast cancer cells exit the breast,
4 enter the vasculature and/or lymphatics, reside in the lymph nodes, evade immune clearance,
5 breach the blood-brain barrier and colonize the brain. These data demonstrate that in humans, a
6 WDR5 non-coding RNA transcript with unknown function is significantly up-regulated in brain
7 metastatic breast cancer.
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28

References

1. Lin, N.U., Amiri-Kordestani, L., Palmieri, D., Liewehr, D.J. and Steeg, P.S., 2013. CNS metastases in breast cancer: old challenge, new frontiers.
2. Bendell, J.C., Domchek, S.M., Burstein, H.J., Harris, L., Younger, J., Kuter, I., Bunnell, C., Rue, M., Gelman, R. and Winer, E., 2003. Central nervous system metastases in women who receive trastuzumab-based therapy for metastatic breast carcinoma. *Cancer*, 97(12), pp.2972-2977.
3. Tsukada, Y., Fouad, A., Pickren, J.W. and Lane, W.W., 1983. Central nervous system metastasis from breast carcinoma autopsy study. *Cancer*, 52(12), pp.2349-2354.
4. Weigman, V.J., Chao, H.H., Shabalin, A.A., He, X., Parker, J.S., Nordgard, S.H., Grushko, T., Huo, D., Nwachukwu, C., Nobel, A. and Kristensen, V.N., 2012. Basal-like Breast cancer DNA copy number losses identify genes involved in genomic instability, response to therapy, and patient survival. *Breast cancer research and treatment*, 133(3), pp.865-880.
5. Sinn, B.V., Fu, C., Lau, R., Litton, J., Tsai, T.H., Murthy, R., Tam, A., Andreopoulou, E., Gong, Y., Murthy, R. and Gould, R., 2019. SET ER/PR: a robust 18-gene predictor for sensitivity to endocrine therapy for metastatic breast cancer. *NPJ breast cancer*, 5(1), pp.1-8.
6. Awada, A., Colomer, R., Inoue, K., Bondarenko, I., Badwe, R.A., Demetriou, G., Lee, S.C., Mehta, A.O., Kim, S.B., Bachelot, T. and Goswami, C., 2016. Neratinib plus paclitaxel vs trastuzumab plus paclitaxel in previously untreated metastatic ERBB2-positive breast cancer: the NEfERT-T randomized clinical trial. *JAMA oncology*, 2(12), pp.1557-1564.
7. Györfy, B., Lanczky, A., Eklund, A.C., Denkert, C., Budczies, J., Li, Q. and Szallasi, Z., 2010. An online survival analysis tool to rapidly assess the effect of 22,277 genes on breast cancer prognosis using microarray data of 1,809 patients. *Breast cancer research and treatment*, 123(3), pp.725-731.

Rank: 148
Probe ID: 8115
p-value: 5.56E-04
t: -4.38
B: -0.026
Gene: LOC401127
Gene name: hypothetical LOC401127

Chart 1: WDR5p, also known as LOC401127, is differentially expressed in brain metastatic breast cancer when comparing brain metastases to primary tumors of the breast.

The rank of global differential expression, the probe/transcript ID, the *p*-value with respect to differential expression transcriptome-wide, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the groups compared, the gene and gene name are listed in this chart.

Rank:1024
probe ID: A_23_P41267
p-value: 1.94e-07
t: -6.0947807
B: 6.981954
Gene: LOC401127
Gene name: WD repeat domain 5 pseudogene

Chart 2: WDR5p is differentially expressed in the brain metastases of patients with breast cancer when comparing brain metastases to normal breast tissues.

The rank of global differential expression, probe/transcript ID, the *p*-value with respect to differential expression transcriptome-wide, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the groups compared, the gene and gene name are listed in this chart.

WDR5p (LOC401127)

Figure 1: WDR5p is expressed at significantly higher levels in the brain metastases of patients with metastatic breast cancer when compared to primary tumors of the breast.

The mRNA expression level of WDR5p in primary tumors of the breast (left) and in brain metastases of women with metastatic breast cancer (right) is graphically depicted; the result of a statistical test evaluating significance of difference in WDR5p expression between primary tumors and brain metastases is $p=0.000916$.