BEAMSHAPES: A PYTHON PACKAGE TO GENERATE DIRECTIVITY PATTERNS FOR VARIOUS SOUND SOURCE MODELS

A PREPRINT

Thejasvi Beleyur *

thejasvib@gmail.com

August 5, 2021

Abstract

beamshapes is an open-source Python package that implements various directivity patterns for sound sources. While there is an abundance of published directivity patterns in the literature - their computational implementations often remain as in-house scripts in proprietary languages. **beamshapes** overcomes this gap, and provides acousticians and bioacousticians easily accessible implementations of sound source directivities.

Keywords acoustics · bioacoustics · directivity · sound radiation · Python · echolocation

1 Summary

Sound sources such as human beings or loudspeakers often exhibit a ‘directionality’ in how loud they sound at different angles. A listener or microphone placed at various angles at a fixed radius will often pick up sometimes drastically different sound levels. The same sound source may however sometimes produce omnidirectional sound fields. The directionality of a sound source can be modelled as a combination of the frequency of the emitted sound and the geometry of the vibrating and non-vibrating parts of the sound source itself. The resulting pattern of sound radiation with angular location is called the *directivity* of the source (Beranek and Mellow 2012).

Directivity (D) describes the relative sound levels at a given angle $D(\theta)$ with relation to the level on-axis ($D(0)$, and thus *directivity* = $\frac{D(\theta)}{D(0)}$). Directivity functions exist for a wide variety of sound sources that can be modelled analytically. A well known example of a directivity function is of the piston in an infinite baffle. The piston is a circular surface of radius a , vibrating back and forth about a hole in an infinite wall or baffle. The directivity is described by $\frac{2J_1(ka \sin \theta)}{ka \sin \theta}$, where J_1 is the Bessel’s function of the first kind, k is the wavenumber, where $k = \frac{2\pi f}{c}$, where f is the frequency of the sound, and c is the speed of sound. The angle of the receiver is θ , which varies from from 0- 2π radians in the azimuth.

Directivity functions can be used in a two-fold manner 1) to deliberately engineer devices to suit particular specifications, eg. loudspeaker sound fields (Beranek and Mellow 2012) and 2) to infer parameters of a sound source itself having assumed a relevant model, eg. estimating direction of call emission (Guarato et al. 2011) and mouth aperture of bat echolocation calls (Jakobsen, Ratcliffe, and Surlykke 2013; Kounitsky et al. 2015).

*1) Centre for the Advanced Study of Collective Behaviour, Konstanz 2) Acoustic and Functional Ecology, Max Planck Institute for Ornithology, Seewiesen 3) Max Planck Institute for Animal Behavior, Radolfzell

Figure 1: Directivity patterns ($\frac{D_\theta}{D_0}$) of the currently implemented sound sources for a common set of ka values for comparison, where k is the wavenumber ($\frac{2\pi f}{c}$) and a is the piston radius - or its equivalent. The directivity pattern shows the ratio between the off-axis sound level at (θ°) to the on-axis level at (0°) in decibels. A) piston in an infinite baffle B) piston in a sphere with the half-angle aperture $\alpha = 30^\circ$, and where the piston radius $a = R \sin \alpha$ C) oscillating cap of a sphere with $\alpha = 30^\circ$, and the equivalent piston radius is $a = R \sin \alpha$ D) vibrating point on a sphere, here the $kR = ka$ for comparison with the other models.

2 Statement of need

A host of published directivity functions exist in the literature, but it is my experience that their computational implementations remain as inhouse scripts that are often in proprietary language platforms. To my knowledge there are no computational implementations of sound source directivities that are open-source and developed using modern software practices such as version-control and unit-testing. In this paper I present **beamshapes**, a Python package that implements directivity patterns for sound source models. As of this publication, **beamshapes** version 0.2.0 implements four sound sources 1) piston in an infinite baffle, 2) point source of a sphere 3) oscillating cap of a sphere and 4) piston in a sphere.

Computational implementations of directivity functions often require long run-times due to the intensive numerical routines and arbitrary precision math involved. Long run-times hinder scientific projects in reducing the number of models and parameter space that can be explored. **beamshapes** boasts parallelised code to generate significant speed-ups in run-times.

The availability of openly-available directivity functions will hopefully stir the acoustics, and specifically the bio-acoustics community to rigorously test and compare sound radiation with models. The availability of multiple implementations allows comparison of data with multiple models. Until recently, computational power has perhaps been a limiting factor to calculating directivity functions. Models with easily calculable outputs have thus been favoured (eg. piston in an infinite baffle), especially in the field of bioacoustics (Strother and Mogus 1970; Mogenssen and Møhl 1979).

Despite the recent availability of computational power, older, simpler models with limited biological relevance continue to dominate the field of bio-acoustics. For instance, the piston in an infinite baffle only predicts the beam-shape for a $\pm 90^\circ$ range off-axis, and assumes front-back symmetry (Figure 1 A). This is unrealistic for most vocalising animals, especially echolocating bats and odontocetes. However, the piston in an infinite baffle continues to be a standard reference model for multiple studies ranging over the past decade (Jakobsen, Ratcliffe, and Surlykke 2013; Kounitsky et al. 2015; Macaulay et al. 2020). The piston in a sphere, for instance recreates many of the highly directional central and side lobes and that are seen in bats, while also predicting sound radiation behind the source (Figure 1 B). In contrast to the echolocation literature, the bird (Witkin 1977; Larsen and Dabelsteen 1990; Brumm 2002; Patricelli, Dantzker, and Bradbury 2007, 2008; Yorzinski and Patricelli 2010) and frog call literature (Gerhardt 1975; Rodriguez and Hödl 2020) for instance has been dominated by quantitative characterisations of sound radiation with no attempts at directly comparing measurements to model predictions. Using model-based directivity patterns to infer source parameters allows the discovery of common parameter spaces that birds occupy, and facilitates cross-species comparisons. The oscillating cap of a sphere and vibrating point on a sphere (Figure 1 C,D) are two other models of potential relevance to bioacousticians attempting to describe the sound radiation of bird and frog calls for instance.

Future releases of **beamshapes** are scheduled to include directivity patterns for additional models of interest such as rectangular cap of a sphere or piston in a closed finite baffle.

3 Software packages used in this work

beamshapes relies on the Python open-source ecosystem and is built on the numpy, scipy, sympy, mpmath and flint libraries (Harris et al. 2020; Virtanen et al. 2020; Meurer et al. 2017; Johansson and others 2021; Hart, Johansson, and Pancratz 2011).

4 Package repository

beamshapes can be currently accessed at https://github.com/thejasvibr/bat_beamshapes.git and the documentation with examples are hosted at <https://beamshapes.readthedocs.io/en/latest/>.

5 Acknowledgements

TB thanks Gaurav Dhariwal for his continual math advice and inputs, and Tim Mellow for strong support through sharing his Mathematica code and timely clarifications. TB thanks Lasse Jakobsen and Holger Goerlitz for productive discussions leading to this project. This work was executed through a combination of TB’s private time, a DAAD stipend, the IMPRS for Organismal Biology, and then finally by a CASCB Medium grant.

References

- Beranek, Leo Leroy, and Tim Mellow. 2012. *Acoustics: Sound Fields and Transducers*. Academic Press.
- Brumm, Henrik. 2002. “Sound Radiation Patterns in Nightingale (*Luscinia Megarhynchos*) Songs.” *Journal für Ornithologie* 143 (4): 468–71.
- Gerhardt, H Carl. 1975. “Sound Pressure Levels and Radiation Patterns of the Vocalizations of Some North American Frogs and Toads.” *Journal of Comparative Physiology* 102 (1): 1–12.
- Guarato, Francesco, Lasse Jakobsen, Dieter Vanderelst, Annemarie Surlykke, and John Hallam. 2011. “A Method for Estimating the Orientation of a Directional Sound Source from Source Directivity and Multi-Microphone Recordings: Principles and Application.” *The Journal of the Acoustical Society of America* 129 (2): 1046–58.
- Harris, Charles R., K. Jarrod Millman, Stéfan J van der Walt, Ralf Gommers, Pauli Virtanen, David Cournapeau, Eric Wieser, et al. 2020. “Array Programming with NumPy.” *Nature* 585: 357–62. <https://doi.org/10.1038/s41586-020-2649-2>.
- Hart, William, Fredrik Johansson, and Sebastian Pancratz. 2011. “FLINT—Fast Library for Number Theory.”

- Jakobsen, Lasse, John M Ratcliffe, and Annemarie Surlykke. 2013. “Convergent Acoustic Field of View in Echolocating Bats.” *Nature* 493 (7430): 93–96.
- Johansson, Fredrik, and others. 2021. *Mpmath: A Python Library for Arbitrary-Precision Floating-Point Arithmetic (Version 1.2.1)*.
- Kounitsky, Pavel, Jens Rydell, Eran Amichai, Arjan Boonman, Ofri Eitan, Anthony J Weiss, and Yossi Yovel. 2015. “Bats Adjust Their Mouth Gape to Zoom Their Biosonar Field of View.” *Proceedings of the National Academy of Sciences* 112 (21): 6724–29.
- Larsen, Ole Næsbye, and Torben Dabelsteen. 1990. “Directionality of Blackbird Vocalization. Implications for Vocal Communication and Its Further Study.” *Ornis Scandinavica*, 37–45.
- Macaulay, Jamie DJ, Chloe E Malinka, Douglas Gillespie, and Peter T Madsen. 2020. “High Resolution Three-Dimensional Beam Radiation Pattern of Harbour Porpoise Clicks with Implications for Passive Acoustic Monitoring.” *The Journal of the Acoustical Society of America* 147 (6): 4175–88.
- Meurer, Aaron, Christopher P Smith, Mateusz Paprocki, Ondrej ertk, Sergey B Kirpichev, Matthew Rocklin, AMiT Kumar, et al. 2017. “SymPy: Symbolic Computing in Python.” *PeerJ Computer Science* 3: e103.
- Mogensen, Finn, and Bertel Møhl. 1979. “Sound Radiation Patterns in the Frequency Domain of Cries from a Vespertilionid Bat.” *Journal of Comparative Physiology* 134 (2): 165–71.
- Patricelli, Gail L, Marc S Dantzker, and Jack W Bradbury. 2007. “Differences in Acoustic Directionality Among Vocalizations of the Male Red-Winged Blackbird (*Agelaius Pheoniceus*) Are Related to Function in Communication.” *Behavioral Ecology and Sociobiology* 61 (7): 1099–1110.
- . 2008. “Acoustic Directionality of Red-Winged Blackbird (*Agelaius Pheoniceus*) Song Relates to Amplitude and Singing Behaviours.” *Animal Behaviour* 76 (4): 1389–1401.
- Rodrguez, Camilo, and Walter Hödl. 2020. “Sound Radiation Pattern of the Advertisement Call of the Highly Territorial Poison Frog *Allobates Femoralis*.” *Behavioural Processes* 170: 103996.
- Strother, GK, and M Mogus. 1970. “Acoustical Beam Patterns for Bats: Some Theoretical Considerations.” *The Journal of the Acoustical Society of America* 48 (6B): 1430–32.
- Virtanen, Pauli, Ralf Gommers, Travis E. Oliphant, Matt Haberland, Tyler Reddy, David Cournapeau, Evgeni Burovski, et al. 2020. “SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python.” *Nature Methods* 17: 261–72. <https://doi.org/10.1038/s41592-019-0686-2>.
- Witkin, Steve R. 1977. “The Importance of Directional Sound Radiation in Avian Vocalization.” *The Condor* 79 (4): 490–93.
- Yorzinski, Jessica L, and Gail L Patricelli. 2010. “Birds Adjust Acoustic Directionality to Beam Their Antipredator Calls to Predators and Conspecifics.” *Proceedings of the Royal Society B: Biological Sciences* 277 (1683): 923–32.