IL2R γ is differentially expressed in the lymph node metastases of patients with breast cancer.

Shahan Mamoor, MS¹

¹shahanmamoor@gmail.com

East Islip, NY USA

Metastasis to the brain is a clinical problem in patients with breast cancer¹⁻³. Between the breast and the brain reside the secondary lymphoid organ, the lymph nodes. We mined published microarray data^{4,5} to compare primary and metastatic tumor transcriptomes for the discovery of genes associated with metastasis to the lymph nodes in humans with metastatic breast cancer. We found that the gamma subunit of the IL-2 receptor, IL2RG, was among the genes whose expression was most different in the lymph node metastases of patients with metastatic breast cancer as compared to primary tumors of the breast. IL2RG mRNA was present at increased quantities in lymph node metastases as compared to primary tumors of the breast. Importantly, expression of IL2RG in primary tumors of the breast was correlated with patient overall survival, in lymph node positive patients but not in lymph node negative patients. Modulation of IL2R γ expression may be relevant to the biology by which tumor cells metastasize from the breast to the lymph nodes and the brain in humans with metastatic breast cancer. These data prompt questioning of the use of NSG mice, deficient in IL2R γ , for the study of human breast cancer metastasis⁶.

Keywords: breast cancer, metastasis, brain metastasis, central nervous system metastasis, lymph node metastasis, metastatic breast cancer, the gamma subunit of the IL-2 receptor, IL2RG, systems biology of breast cancer, targeted therapeutics in breast cancer.

One report described a 34% incidence of central nervous system metastases in patients treated with trastuzumab for breast cancer². More recently, the NEfERT-T clinical trial⁶ which compared administration of either neratinib or trastuzumab in conjunction with paclitaxel demonstrated that in a randomized, controlled setting, in breast cancer patients treated with neratinib, not only was the incidence of central nervous system recurrence significantly lower, the time to central nervous system metastasis was significantly delayed as compared to patients administered trastuzumab⁷. The alarmingly high rate of central nervous system metastasis described, as well as data, both anecdotal² and from a randomized, controlled setting⁷ illustrating that treatment with trastuzumab may be associated with these events demands an enhanced understanding of the transcriptional makeup of brain and lymph node metastatic tissues to support identification of therapeutic targets, whether they are treatment related or not. The lymph nodes reside between the breast and the brain. We performed a global comparative analysis of lymph node metastatic tumor tissues in patients with metastatic breast cancer, compared to primary tumors and normal breast tissues^{4,5}. We discovered transcriptome-wide differential expression of the gene encoding the gamma subunit of the IL-2 receptor, IL2RG, in lymph node metastatic tissues of patients with metastatic breast cancer.

Methods

We used datasets GSE10893⁴ and GSE124648⁵ for this global differential gene expression analysis of brain metastatic breast cancer in conjunction with GEO2R. GSE10893 was generated using Agilent-011521 Human 1A Microarray G4110A technology with $n=71$ primary breast tumors and $n=7$ lymph node metastases from patients with breast cancer; analysis was performed using platform GPL887. GSE124648 was generated using Affymetrix Human Genome U133A array technology with $n=10$ normal breast tissues and $n=44$ lymph node metastases from patients with breast cancer; analysis was performed using platform GPL96. The Benjamini and Hochberg method of p -value adjustment was used for ranking of differential expression but raw p -values were used to assess statistical significance of global differential expression. Log-transformation of data was auto-detected, and the NCBI generated category of platform annotation was used. A statistical test was performed to evaluate whether IL2RG gene expression was significantly different between primary tumors of the breast and lymph node metastases in humans with breast cancer using a two-tailed t-test. For Kaplan-Meier survival analysis, we used the Kaplan-Meier plotter online tool⁸ for correlation of IL2RG mRNA expression levels with overall survival in $n=1402$ breast cancer patients, in $n=313$ lymph node positive patients, and $n=594$ lymph node negative patients.

Results

We performed global comparative transcriptome analysis of metastatic tumor tissues of patients with metastatic breast cancer using published microarray data^{4,5} to describe the transcriptional landscape of metastasis in human breast cancer in an unbiased fashion and for the discovery of novel therapeutic targets.

IL2RG is differentially expressed in the lymph node metastases of patients with metastatic breast cancer.

Through blind, systems-level analysis of published microarray data⁴, we identified the gamma subunit of the IL-2 receptor, encoded by IL2RG, as a differentially expressed gene in the lymph node metastatic tissues of humans with breast cancer (Table 1). When sorting each of the genes expressed in lymph node metastases based on significance of difference as compared to primary tumors of the breast in patients with breast cancer, IL2RG ranked 184 out of 18034 total transcripts (Chart 1), equating to 99.0% differential expression. Differential expression of IL2RG in the lymph node metastases of patients with metastatic breast cancer was statistically significant (Chart 1; $p=2.09\text{e-}03$).

We queried a second microarray dataset⁵ to attempt to validate transcriptome-wide differential expression of IL2RG in the lymph node metastases of patients with metastatic breast cancer. We again observed IL2RG to be among the genes expressed most quantitatively differently in the lymph node metastases of patients with metastatic breast cancer as compared to primary tumors of the breast. When sorting each of the genes expressed in lymph node metastases based on significance of difference as compared to normal breast tissues, IL2RG ranked 6806 out of 22283 total transcripts (Chart 2), equating to 69.5% differential expression. Differential expression of IL2RG in the lymph node metastases of patients with metastatic breast cancer was statistically significant (Chart 2; $p=2.42e-02$).

IL2RG is expressed at higher levels in the lymph node metastases of patients with metastatic breast cancer.

We obtained exact mRNA expression levels for IL2RG, in primary tumors of the breast and in lymph node metastases of patients with metastatic breast cancer to determine direction and statistical significance of change in IL2RG expression in lymph node metastatic tissues. We observed significantly increased expression of IL2RG in the lymph node metastases of patients with breast cancer as compared to primary tumors of the breast: IL2RG was expressed at 0.19 ± 0.73 arbitrary units (AU) in primary tumors of the breast, while it was expressed at 1.16 ± 1.16 AU in lymph node metastatic tissues (Figure 1). The difference in IL2RG mRNA levels between primary tumors of the breast and lymph node metastases was statistically significant (Figure 1; $p=0.002415$).

IL2RG expression is significantly correlated with survival outcomes in human breast cancer.

We performed Kaplan-Meier survival analysis⁸ in $n=1402$ breast cancer patients in total, to evaluate whether IL2RG tumor expression was correlated with survival outcomes in breast cancer. We observed a statistically significant correlation between primary tumor expression of IL2RG and overall survival (OS) in patients with breast cancer (Figure 2). Patients whose primary tumors expressed low levels of IL2RG possessed median OS of 80.64 months, while patients whose tumors expressed high levels of IL2RG possessed median OS of 138 months. This difference in OS based on IL2RG tumor expression in patients with breast cancer was statistically significant (Figure 2, Chart 3; logrank p -value: 0.0039; (hazard ratio: 0.73 (0.59-0.9))).

We also performed Kaplan-Meier survival analysis⁸ in $n=313$ lymph node positive breast cancer patients and $n=594$ lymph node negative breast cancer patients, to evaluate whether lymph node status influenced correlation of IL2RG tumor expression with survival outcomes in human breast cancer. We observed a correlation between primary tumor expression of IL2RG and overall survival (OS) in lymph node positive breast cancer (Figure 2). Lymph node positive patients whose primary tumors expressed low levels of IL2RG possessed median OS of 38.57 months, while lymph node positive patients whose tumors expressed high levels of IL2RG possessed median OS of 55 months. This difference in OS based on IL2RG tumor expression in lymph node positive patients with breast cancer was statistically significant (Figure 2, Chart 3; logrank p -value: 0.045 (hazard ratio: 0.67 (0.45-0.99))). Thus, overall survival based on IL2RG primary tumor expression was markedly shorter in lymph node positive patients than in the general breast cancer patient population. IL2RG primary tumor expression was not significantly correlated with overall survival outcomes in lymph node negative patients (Figure 2, Chart 3; logrank p -value: 0.49; hazard ratio: 0.88 (0.61-1.27)).

Thus, by mining published microarray data^{4,5} in an unbiased and systematic fashion, we identified IL2RG as among the genes whose expression was most different, transcriptome-wide, in the lymph node metastases of patients with breast cancer, when compared to primary tumors; we observed significantly increased expression of IL2RG in lymph node metastases as compared to primary tumors of the breast.

Further, we found a significant correlation between IL2RG expression and patient survival outcomes, as overall survival was significantly greater in patients whose primary tumors expressed higher levels of IL2RG as compared to patients whose primary tumors expressed lower levels of IL2RG, and this correlation was significant in lymph node positive patients but not in lymph node negative patients.

Discussion

We provided evidence here that the gamma subunit of the IL-2 receptor, IL2RG, is among the genes whose expression is most different in the lymph node metastases of patients with metastatic breast cancer as compared to primary tumors of the breast, that IL2RG mRNA is present at significantly increased quantities in lymph node metastatic tissues as compared to primary tumors of the breast, and that primary tumor IL2RG expression is significantly correlated with patient survival outcomes in human breast cancer, in all patients and in lymph node positive patients but not in lymph node negative patients. Evaluation of the effects of genetic depletion of IL2RG in mouse models of metastatic breast cancer on metastasis to the central nervous system is merited. Modulation of IL2RG expression may be relevant to the processes by which breast cancer cells exit the breast, enter the vasculature and/or lymphatics, reside in the lymph nodes, evade immune clearance, breach the blood-brain barrier and colonize the brain. These data suggest that *in vivo* analysis of breast cancer lymph node metastasis in NSG (NOD-SCID- IL2R γ null) mice may be difficult to interpret⁶.

References

1. Lin, N.U., Amiri-Kordestani, L., Palmieri, D., Liewehr, D.J. and Steeg, P.S., 2013. CNS metastases in breast cancer: old challenge, new frontiers.
2. Bendell, J.C., Domchek, S.M., Burstein, H.J., Harris, L., Younger, J., Kuter, I., Bunnell, C., Rue, M., Gelman, R. and Winer, E., 2003. Central nervous system metastases in women who receive trastuzumab-based therapy for metastatic breast carcinoma. *Cancer*, 97(12), pp.2972-2977.
3. Tsukada, Y., Fouad, A., Pickren, J.W. and Lane, W.W., 1983. Central nervous system metastasis from breast carcinoma autopsy study. *Cancer*, 52(12), pp.2349-2354.
4. Weigman, V.J., Chao, H.H., Shabalin, A.A., He, X., Parker, J.S., Nordgard, S.H., Grushko, T., Huo, D., Nwachukwu, C., Nobel, A. and Kristensen, V.N., 2012. Basal-like Breast cancer DNA copy number losses identify genes involved in genomic instability, response to therapy, and patient survival. *Breast cancer research and treatment*, 133(3), pp.865-880.
5. Sinn, B.V., Fu, C., Lau, R., Litton, J., Tsai, T.H., Murthy, R., Tam, A., Andreopoulou, E., Gong, Y., Murthy, R. and Gould, R., 2019. SET ER/PR: a robust 18-gene predictor for sensitivity to endocrine therapy for metastatic breast cancer. *NPJ breast cancer*, 5(1), pp.1-8.
6. Puchalapalli, M., Zeng, X., Mu, L., Anderson, A., Hix Glickman, L., Zhang, M., Sayyad, M.R., Mosticone Wangenstein, S., Clevenger, C.V. and Koblinski, J.E., 2016. NSG mice provide a better spontaneous model of breast cancer metastasis than athymic (nude) mice. *PLoS One*, 11(9), p.e0163521.
7. Awada, A., Colomer, R., Inoue, K., Bondarenko, I., Badwe, R.A., Demetriou, G., Lee, S.C., Mehta, A.O., Kim, S.B., Bachelot, T. and Goswami, C., 2016. Neratinib plus paclitaxel vs trastuzumab plus paclitaxel in previously untreated metastatic ERBB2-positive breast cancer: the NEfERT-T randomized clinical trial. *JAMA oncology*, 2(12), pp.1557-1564.
8. Györfy, B., Lanczky, A., Eklund, A.C., Denkert, C., Budczies, J., Li, Q. and Szallasi, Z., 2010. An online survival analysis tool to rapidly assess the effect of 22,277 genes on breast cancer prognosis using microarray data of 1,809 patients. *Breast cancer research and treatment*, 123(3), pp.725-731.

Rank: 184
Probe ID: 18558
p-value: 2.09e-03
t: -3.1817845
B: -1.4344
Gene: IL2RG
Gene name: interleukin 2 receptor subunit gamma

Chart 1: IL2RG is differentially expressed in lymph node metastatic breast cancer when comparing lymph node metastases to primary tumors of the breast.

The rank of global differential expression, the probe/transcript ID, the *p*-value with respect to differential expression transcriptome-wide, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the groups compared, the gene and gene name are listed in this chart.

Rank: 6806
Probe ID: 204116_at
p-value: 2.42e-02
t: -2.317223
B: -4.383545
Gene: IL2RG
Gene name: interleukin 2 receptor subunit gamma

Chart 2: IL2RG is differentially expressed in lymph node metastatic breast cancer when comparing lymph node metastases to normal breast tissues.

The rank of global differential expression, the probe/transcript ID, the *p*-value with respect to differential expression transcriptome-wide, *t*, a moderated *t*-statistic, *B*, the log-odds of differential expression between the groups compared, the gene and gene name are listed in this chart.

Figure 1: IL2RG is expressed at significantly higher levels in the lymph node metastases of patients with metastatic breast cancer when compared to primary tumors of the breast.

The mRNA expression level of IL2RG in primary tumors of the breast (left) and in lymph node metastases of women with metastatic breast cancer (right) is graphically depicted; the result of a statistical test evaluating significance of difference in IL2RG expression between primary tumors of the breast and lymph node metastases, a *p*-value, is *p*=0.002415.

Figure 2: IL2RG primary tumor expression is significantly correlated with overall survival in all patients with breast cancer and in lymph node positive patients but not in lymph node negative patients.

Depicted in this Kaplan-Meier plot (**above**) is the probability of overall survival for $n=1402$ total patients stratified into two groups, based on low or high expression of IL2RG in patient primary tumors, in the lower survival quartile. The log rank p -value denoting statistical significance of difference in overall survival when comparing the two groups, as well as hazard ratio for this comparison is listed above. Listed below is the number of patients at risk (number of patients alive) per interval, after stratification based on IL2RG expression; in the first interval, number at risk is number of patients alive; in each subsequent interval, number at risk is the number at risk less those who have expired or are censored. Below, Kaplan-Meier plots for probability of overall survival in $n=313$ lymph node positive patients (**left**) and $n=594$ lymph node negative patients (**right**).

All patients

Low IL2RG expression: 80.64 months

High IL2RG expression: 138 months

Lymph node positive patients

Low IL2RG expression: 38.57 months

High IL2RG expression: 55 months

Chart 3: Median overall survival is inferior in patients with low primary tumor expression of IL2RG.

The median OS (overall survival) of patients with low primary tumor expression of IL2RG and high primary tumor expression of IL2RG is listed in this chart, for $n=1402$ breast cancer patients and for $n=313$ lymph node positive patients.